

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12662 A2

(51) International Patent Classification⁷: **C07K 14/00**

(21) International Application Number: **PCT/US00/22315**

(22) International Filing Date: 14 August 2000 (14.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/149,641 17 August 1999 (17.08.1999) US
60/164,203 9 November 1999 (09.11.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:
US 60/149,641 (CIP)
Filed on 17 August 1999 (17.08.1999)
US 60/164,203 (CIP)
Filed on 9 November 1999 (09.11.1999)

(71) Applicant (for all designated States except US): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LAL, Preeti** [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **TANG, Y., Tom** [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **BANDMAN, Olga** [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). **BURFORD, Neil** [GB/US]; 1308 4th Avenue, San

Francisco, CA 94122 (US). **AZIMZAI, Yalda** [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **LU, Dyung, Aina, M.** [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). **PATTERSON, Chandra** [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **MEMBRANE ASSOCIATED PROTEINS**

(57) Abstract: The invention provides human membrane associated proteins (MEMAP) and polynucleotides which identify and encode MEMAP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of MEMAP.

WO 01/12662 A2



MEMBRANE ASSOCIATED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of membrane associated proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cell proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders.

BACKGROUND OF THE INVENTION

Eukaryotic cells are surrounded by plasma membranes which enclose the cell and maintain an environment inside the cell that is distinct from its surroundings. In addition, eukaryotic organisms are distinct from prokaryotes in possessing many intracellular organelle and vesicle structures. Many of the metabolic reactions which distinguish eukaryotic biochemistry from prokaryotic biochemistry take place within these structures. The plasma membrane and the membranes surrounding organelles and vesicles are composed of phosphoglycerides, fatty acids, cholesterol, phospholipids, glycolipids, proteoglycans, and proteins. These components confer identity and functionality to the membranes with which they associate.

Integral Membrane Proteins

The majority of known integral membrane proteins are transmembrane proteins (TM) which are characterized by an extracellular, a transmembrane, and an intracellular domain. TM domains are typically comprised of 15 to 25 hydrophobic amino acids which are predicted to adopt an α -helical conformation. TM proteins are classified as bitopic (Types I and II) and polytopic (Types III and IV) (Singer, S.J. (1990) Annu. Rev. Cell Biol. 6:247-96). Bitopic proteins span the membrane once while polytopic proteins contain multiple membrane-spanning segments. TM proteins that act as cell-surface receptor proteins involved in signal transduction include growth and differentiation factor receptors, and receptor-interacting proteins such as *Drosophila* pecanex and frizzled proteins, LIV-1 protein, NF2 protein, and GNS1/SUR4 eukaryotic integral membrane proteins. TM proteins also act as transporters of ions or metabolites, such as gap junction channels (connexins) and ion channels, and as cell anchoring proteins, such as lectins, integrins, and fibronectins. TM proteins act as vesicle organelle-forming molecules, such as calveolins, or as cell recognition molecules, such as cluster of differentiation (CD) antigens, glycoproteins, and mucins.

Many membrane proteins (MPs) contain amino acid sequence motifs that target these proteins to specific subcellular sites. Examples of these motifs include PDZ domains, KDEL, RGD, NGR, and GSL sequence motifs, von Willebrand factor A (vWFA) domains, and EGF-like domains. RGD, NGR, and GSL motif-containing peptides have been used as drug delivery agents in cancer

treatments which target tumor vasculature (Arap, W. et al. (1998) Science, 279:377-380). Furthermore, MPs may also contain amino acid sequence motifs, such as the carbohydrate recognition domain (CRD), also known as the C-type lectin domain, that mediate interactions with extracellular or intracellular molecules.

5 Membrane proteins may also interact with and regulate the properties of the membrane lipids. Phospholipid scramblase, a type II plasma membrane protein, mediates calcium dependent movement of phospholipids (PL) between membrane leaflets. Calcium induced remodeling of plasma membrane PL plays a key role in expression of platelet anticoagulant activity and in clearance of injured or apoptotic cells (Zhou Q. et al. (1997) J. Biol. Chem. 272:18240-18244). Scott syndrome, a
10 bleeding disorder, is caused by an inherited deficiency in plasma membrane PL scramblase function (Online Mendelian Inheritance in Man (OMIM) *262890 Platelet Receptor for Factor X, Deficiency of).

Chemical modification of amino acid residue side chains alters the manner in which MPs interact with other molecules, such as phospholipid membranes. Examples of such chemical
15 modifications to amino acid residue side chains are covalent bond formation with glycosaminoglycans, oligosaccharides, phospholipids, acetyl and palmitoyl moieties, ADP-ribose, phosphate, and sulphate groups.

One function of TM proteins is to facilitate cell-cell communication. The slit proteins are extracellular matrix proteins expressed by cells at the ventral midline of the nervous system. Slit
20 proteins are ligands for the repulsive guidance receptor Robo and thus play a role in repulsive axon guidance (Brose, K. et al. (1999) Cell 96:795-806).

In some cases TM proteins serve as transporters or channels in the cell membrane. For example, the mouse transporter protein (MTP) has four transmembrane domains and resides in an intracellular membrane compartment. MTP can mediate transport of nucleosides in vitro. The role
25 of MTP in the cell may therefore be to transfer nucleosides between the cytosol and the lumen of intracellular organelles (Hogue, D. L. (1996) J. Biol. Chem. 271:9801-9808). The human stomatin-like protein (hSLP-1), expressed primarily in the brain, contains an N-terminal domain similar to the erythrocyte internal membrane protein stomatin, as well as a non-specific lipid transfer protein domain at the C-terminus. hSLP-1 is the human homologue of the C. elegans behavioral gene unc-
30 24, which is believed to be involved in lipid transfer between closely apposed membranes (Seidel, G. and Prohaska, R (1998) Gene 225:23-29).

The transmembrane 4 superfamily (TM4SF) or tetraspan family is a multigene family encoding type III integral membrane proteins (Wright, M.D. and Tomlinson, M.G. (1994) Immunol. Today 15:588-594). TM4SF is comprised of membrane proteins which traverse the cell membrane
35 four times. Members of the TM4SF include platelet and endothelial cell membrane proteins,

melanoma-associated antigens, leukocyte surface glycoproteins, colonal carcinoma antigens, tumor-associated antigens, and surface proteins of the schistosome parasites (Jankowski, S.A. (1994) *Oncogene* 9:1205-1211). Members of the TM4SF share about 25-30% amino acid sequence identity with one another.

5 A number of TM4SF members have been implicated in signal transduction, control of cell adhesion, regulation of cell growth and proliferation, including development and oncogenesis, and cell motility, including tumor cell metastasis. Expression of TM4SF proteins is associated with a variety of tumors and the level of expression may be altered when cells are growing or activated.

Tumor antigens are cell surface molecules that are differentially expressed in tumor cells
10 relative to normal cells. Tumor antigens distinguish tumor cells immunologically from normal cells and provide diagnostic and therapeutic targets for human cancers (Takagi, S. et al. (1995) *Int. J. Cancer* 61: 706-715; Liu, E. et al. (1992) *Oncogene* 7: 1027-1032). For example, the biliary glycoprotein-encoding gene is a member of the human carcinoembryonic antigen family, which are important tumor markers for colorectal carcinomas (Hammarstrom, S. (1999) *Semin. Cancer Bio.*
15 9:67-81). Another example is the neuron and testis specific protein Ma1, a marker for paraneoplastic neuronal disorders (Dalmau, J. et al. (1999) *Brain* 122:27-39).

Other types of cell surface antigens include those identified on leukocytic cells of the immune system. These antigens have been identified using systematic, monoclonal antibody (mAb)-based "shot gun" techniques. These techniques have resulted in the production of hundreds of mAbs
20 directed against unknown cell surface leukocytic antigens. These antigens have been grouped into "clusters of differentiation" based on common immunocytochemical localization patterns in various differentiated and undifferentiated leukocytic cell types. Antigens in a given cluster are presumed to identify a single cell surface protein and are assigned a "cluster of differentiation" or "CD" designation. Some of the genes encoding proteins identified by CD antigens have been cloned and
25 verified by standard molecular biology techniques. CD antigens have been characterized as both transmembrane proteins and cell surface proteins anchored to the plasma membrane via covalent attachment to fatty acid-containing glycolipids such as glycosylphosphatidylinositol (GPI). (Reviewed in Barclay, A. N. et al. (1995) The Leucocyte Antigen Facts Book, Academic Press, San Diego, CA, pp. 17-20.)

30 The TM cell surface glycoprotein CD69 is an early activation antigen of T lymphocytes. CD69 is homologous to members of a supergene family of type II integral membrane proteins having C-type lectin domains. Although the precise functions of the CD-69 antigen is not known, evidence suggests that these proteins transmit mitogenic signals across the plasma membrane and are up-regulated in response to lymphocyte activation (Hamann, J. et. al. (1993) *J. Immunol.* 150:4920-
35 4927).

Macrophages are involved in functions including clearance of senescent or apoptotic cells, cytokine production, hemopoiesis, bone resorption, antigen transport, and neuroendocrine regulation. These diverse roles are influenced by specialized macrophage plasma membrane proteins. The murine macrophage restricted C-type lectin is a type II integral membrane protein expressed exclusively in macrophages. The strong expression of this protein in bone marrow suggests a hemopoietic function, while the lectin domain suggests it may be involved in cell-cell recognition (Balch, S. G. et al. (1998) *J. Biol. Chem.* 273:18656-18664).

The surface of red blood cells is populated with characteristic glycoproteins, such as the major sialoglycoproteins glycophorin A and B. Red blood cells lacking either glycophorin A or B are resistant to infection with the malaria parasite *Plasmodium falciparum* (OMIM Entry 111300 Blood Group-MN Locus). White blood cells also possess characteristic surface glycoproteins, such as the plasma cell glycoprotein-1 (PC-1). PC-1 is expressed on the surface of plasma cells, which are terminally differentiated, antibody-secreting B-lymphocytes. The extracellular domain of PC-1 has nucleotide phosphodiesterase (pyrophosphatase) activity (Funakoshi, I. et al. (1992) *Arch. Biochem. Biophys.* 295:180-187). Phosphodiesterase activity is associated with the hydrolytic removal of nucleotide subunits from oligonucleotides. Although the precise physiological role of PC-1 is not clear, increased PC-1 phosphodiesterase activity has been correlated with insulin resistance in patients with noninsulin-dependent diabetes mellitus, with abnormalities of bone mineralization and calcification, and with defects in renal tubule function. In addition, it appears that hPC-1 and mPC-1 are members of a multigene family of transmembrane phosphodiesterases with extracellular active sites. These enzymes may play a role in regulating the concentration of pharmacologically active extracellular compounds such as adenosine or other nucleotide derivatives in a variety of tissues and cell types. (Reviewed in Goding, J. W. et al. (1998) *Immunol. Rev.* 161:11-26.)

Peripheral and Anchored Membrane Proteins

Some membrane proteins are not membrane-spanning but are attached to the plasma membrane via membrane anchors or interactions with integral membrane proteins. Membrane anchors are covalently joined to a protein post-translationally and include such moieties as prenyl, myristyl, and glycosylphosphatidyl inositol (GPI) groups. Membrane localization of peripheral and anchored proteins is important for their function in processes such as receptor-mediated signal transduction. For example, prenylation of Ras is required for its localization to the plasma membrane and for its normal and oncogenic functions in signal transduction.

The pancortins are a group of four glycoproteins which are predominantly expressed in the cerebral cortex of adult rodents. Immunological localization indicates that the pancortins are endoplasmic reticulum anchored proteins. The pancortins share a common sequence in the middle of

their structure, but have alternative sequences at both ends due to differential promoter usage and alternative splicing. Each pancortin appears to be differentially expressed and may perform different functions in the brain (Nagano, T. et al. (1998) Mol. Brain Res. 53:13-23).

The discovery of new membrane associated proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, membrane associated proteins, referred to collectively as "MEMAP" and individually as "MEMAP-1," "MEMAP-2," "MEMAP-3," "MEMAP-4," "MEMAP-5," "MEMAP-6," "MEMAP-7," "MEMAP-8," "MEMAP-9," "MEMAP-10," "MEMAP-11," "MEMAP-12," "MEMAP-13," "MEMAP-14," "MEMAP-15," "MEMAP-16," "MEMAP-17," "MEMAP-18," "MEMAP-19," "MEMAP-20," "MEMAP-21," "MEMAP-22," "MEMAP-23," "MEMAP-24," "MEMAP-25," "MEMAP-26," "MEMAP-27," "MEMAP-28," "MEMAP-29," "MEMAP-30," "MEMAP-31," "MEMAP-32," "MEMAP-33," "MEMAP-34," "MEMAP-35," "MEMAP-36," and "MEMAP-37." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-37.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-37. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:38-74.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid

sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of
5 SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising an amino acid
10 sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of
15 SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a
20 polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino
25 acid sequence selected from the group consisting of SEQ ID NO:1-37.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, b) a naturally occurring polynucleotide sequence having at least
30 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide
35 sequence selected from the group consisting of a) a polynucleotide sequence selected from the group

consisting of SEQ ID NO:38-74, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence

selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting
5 agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment the composition.

10 Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence
15 selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In
20 another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional MEMAP, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an
25 amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a)
30 combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a)
35 an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally

occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:38-74, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of the above polynucleotide sequence; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding MEMAP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of MEMAP.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding MEMAP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the

invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

“MEMAP” refers to the amino acid sequences of substantially purified MEMAP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term “agonist” refers to a molecule which intensifies or mimics the biological activity of MEMAP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of MEMAP either by directly interacting with MEMAP or by acting on components of the biological pathway in which MEMAP participates.

An “allelic variant” is an alternative form of the gene encoding MEMAP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding MEMAP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as MEMAP or a polypeptide with at least one functional characteristic of MEMAP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding MEMAP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding MEMAP. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent MEMAP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of MEMAP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to a sequence of a naturally occurring

protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence.

Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of MEMAP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of MEMAP either by directly interacting with MEMAP or by acting on components of the biological pathway in which MEMAP participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind MEMAP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the

designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic MEMAP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding MEMAP or fragments of MEMAP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (PE Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

Original Residue	Conservative Substitution
Ala	Gly, Ser
Arg	His, Lys
Asn	Asp, Gln, His
Asp	Asn, Glu
Cys	Ala, Ser
Gln	Asn, Glu, His
Glu	Asp, Gln, His

	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
5	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
10	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of MEMAP or the polynucleotide encoding MEMAP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the

present embodiments.

A fragment of SEQ ID NO:38-74 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:38-74, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:38-74 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:38-74 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:38-74 and the region of SEQ ID NO:38-74 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-37 is encoded by a fragment of SEQ ID NO:38-74. A fragment of SEQ ID NO:1-37 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-37. For example, a fragment of SEQ ID NO:1-37 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-37. The precise length of a fragment of SEQ ID NO:1-37 and the region of SEQ ID NO:1-37 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent-identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms

is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at

<http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence

analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>.

The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative

substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity.

Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide

sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of MEMAP which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of MEMAP which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of MEMAP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of MEMAP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an MEMAP may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of MEMAP.

"Probe" refers to nucleic acid sequences encoding MEMAP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

5 "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

10 Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

15 Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs
20 can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to
25 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from
30 megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection
35 programs may also be obtained from their respective sources and modified to meet the user's specific

needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both
5 unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

10 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have
15 been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a
20 vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription,
25 translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

30 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic
35 acids encoding MEMAP, or fragments thereof, or MEMAP itself, may comprise a bodily fluid; an

extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with

a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook, J. et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human membrane associated proteins (MEMAP), the polynucleotides encoding MEMAP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, autoimmune/inflammatory, neurological and

gastrointestinal disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding MEMAP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each MEMAP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each MEMAP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding MEMAP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:38-74 and to distinguish between SEQ ID NO:38-74 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express MEMAP as a fraction of total tissues expressing MEMAP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing MEMAP as a fraction of total tissues expressing MEMAP. Column 5 lists the vectors used to subclone each cDNA library. Of particular note is the expression of SEQ ID NO:41, SEQ ID NO:48, and SEQ ID NO:56 in nervous tissues, of SEQ ID NO:52, SEQ ID NO:65, and SEQ ID NO:74 in gastrointestinal tissues, and of SEQ ID NO:55 in hematopoietic/immune tissues.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding MEMAP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO:38 maps to chromosome 4 within the interval from 77.9 to 86.0 centiMorgans, to chromosome 6 within the interval from 132.7 to 144.4 centiMorgans, and to chromosome 14 within the interval from 89.4 to 103.7 centiMorgans. The interval on chromosome 4 from 77.9 to 86.0 centiMorgans also contains a gene associated with deoxycytidine kinase deficiency. The interval on chromosome 6 from 132.7 to 144.4 centiMorgans also contains genes associated with peroxisomal disorders and leukemia. The interval on chromosome 14 from 89.4 to 103.7 centiMorgans also contains genes associated with spinocerebellar ataxia and protease inhibitor deficiencies. SEQ ID NO:39 maps to chromosome 2 within the interval from 236.2 to 269.5 centiMorgans, and to the X chromosome within the interval from 94.4 to 97.4 centiMorgans. The interval on chromosome 2 from 236.2 to 269.5 centiMorgans also contains genes associated with Crigler-Najjar syndrome, Oguchi disease, and oxaolis I. The interval on the X chromosome from 94.4 to 97.4 centiMorgans also contains genes associated with Charcot-Marie tooth disease, X-linked severe combined immunodeficiency, alpha thalassemia/mental retardation syndrome, Menkes' syndrome, and choroideremia. SEQ ID NO:42 maps to chromosome 1 within the interval from 218.2 to 232.0 centiMorgans. This interval also contains genes associated with familial hypertrophic cardiomyopathy, malignant hyperthermia, and hypokalemic periodic paralysis. SEQ ID NO:44 maps to chromosome 7 within the interval from 136.4 to 145.8 centiMorgans, to chromosome 14 within the interval from 28.0 to 32.9 centiMorgans, and to chromosome 14 within the interval from 71.5 to 73.7 centiMorgans. The interval on chromosome 7 from 136.4 to 145.8 centiMorgans also contains genes associated with diphosphoglycerate mutase deficiency. SEQ ID NO:60 maps to chromosome 7 within the interval from 167.6 to 184.0 centiMorgans, and to chromosome 14 within the interval from 50.0 to 59.0 centiMorgans. SEQ ID NO:63 maps to chromosome 8 within the interval from 101.0 to 125.8 centiMorgans, and to chromosome 8 within the interval from 132.4 to 135.1 centiMorgans. SEQ ID NO:67 maps to chromosome 4 within the interval from 145.3 to 146.4 centiMorgans.

The invention also encompasses MEMAP variants. A preferred MEMAP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the MEMAP amino acid sequence, and which contains at least one functional or structural characteristic of MEMAP.

The invention also encompasses polynucleotides which encode MEMAP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:38-74, which encodes MEMAP. The polynucleotide sequences of SEQ ID NO:38-74, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding MEMAP.

In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding MEMAP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID
5 NO:38-74 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:38-74. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of MEMAP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the
10 genetic code, a multitude of polynucleotide sequences encoding MEMAP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the
15 polynucleotide sequence of naturally occurring MEMAP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode MEMAP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring MEMAP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding
20 MEMAP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding MEMAP and its derivatives without altering the encoded amino acid
25 sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode MEMAP and MEMAP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell
30 systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding MEMAP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:38-74 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and
35 S.L. Berger (1987) Methods Enzymol. 152:399-407; Kimmel, A.R. (1987) Methods Enzymol.

152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (PE Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (PE Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (PE Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding MEMAP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National

Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, PE Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode MEMAP may be cloned in recombinant DNA molecules that direct expression of MEMAP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express MEMAP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter MEMAP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of MEMAP, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of

gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding MEMAP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.) Alternatively, MEMAP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (PE Biosystems). Additionally, the amino acid sequence of MEMAP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active MEMAP, the nucleotide sequences encoding MEMAP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding MEMAP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding MEMAP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding MEMAP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no

additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic.

- 5 The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding MEMAP and appropriate transcriptional and translational
10 control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

- 15 A variety of expression vector/host systems may be utilized to contain and express sequences encoding MEMAP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower
20 mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-
25 1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses,
30 adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.)
35 The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding MEMAP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding MEMAP can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding MEMAP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of MEMAP are needed, e.g. for the production of antibodies, vectors which direct high level expression of MEMAP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of MEMAP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, *supra*; and Scorer, *supra*.)

Plant systems may also be used for expression of MEMAP. Transcription of sequences encoding MEMAP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, *supra*; Broglie, *supra*; and Winter, *supra*.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding MEMAP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses MEMAP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of

DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

5 For long term production of recombinant proteins in mammalian systems, stable expression of MEMAP in cell lines is preferred. For example, sequences encoding MEMAP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in
10 enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These
15 include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk*⁻ and *apr*⁻ cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat*
20 confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins
25 (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is
30 also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding MEMAP is inserted within a marker gene sequence, transformed cells containing sequences encoding MEMAP can be identified by the absence of marker gene function.

Alternatively, a marker gene can be placed in tandem with a sequence encoding MEMAP under the control of a single promoter. Expression of the marker gene in response to induction or selection
35 usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding MEMAP and that express MEMAP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of MEMAP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on MEMAP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding MEMAP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding MEMAP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding MEMAP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode MEMAP may be designed to contain signal sequences which direct secretion of MEMAP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of

the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding MEMAP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric MEMAP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of MEMAP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the MEMAP encoding sequence and the heterologous protein sequence, so that MEMAP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled MEMAP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

MEMAP of the present invention or fragments thereof may be used to screen for compounds that specifically bind to MEMAP. At least one and up to a plurality of test compounds may be screened for specific binding to MEMAP. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of

MEMAP, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which MEMAP binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express MEMAP, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing MEMAP or cell membrane fractions which contain MEMAP are then contacted with a test compound and binding, stimulation, or inhibition of activity of either MEMAP or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with MEMAP, either in solution or affixed to a solid support, and detecting the binding of MEMAP to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

MEMAP of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of MEMAP. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for MEMAP activity, wherein MEMAP is combined with at least one test compound, and the activity of MEMAP in the presence of a test compound is compared with the activity of MEMAP in the absence of the test compound. A change in the activity of MEMAP in the presence of the test compound is indicative of a compound that modulates the activity of MEMAP. Alternatively, a test compound is combined with an in vitro or cell-free system comprising MEMAP under conditions suitable for MEMAP activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of MEMAP may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding MEMAP or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of

interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding MEMAP may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding MEMAP can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding MEMAP is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress MEMAP, e.g., by secreting MEMAP in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of MEMAP and membrane associated proteins. In addition, the expression of MEMAP is closely associated with neurological and gastrointestinal tissues, cancer, cell proliferation, and inflammation/trauma. Therefore, MEMAP appears to play a role in cell proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders. In the treatment of disorders associated with increased MEMAP expression or activity, it is desirable to decrease the expression or activity of MEMAP. In the treatment of disorders associated with decreased MEMAP expression or activity, it is desirable to increase the expression or activity of MEMAP.

Therefore, in one embodiment, MEMAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or

activity of MEMAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including

mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas.

In another embodiment, a vector capable of expressing MEMAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of MEMAP including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified MEMAP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of MEMAP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of MEMAP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of MEMAP including, but not limited to, those listed above.

In a further embodiment, an antagonist of MEMAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of MEMAP. Examples of such disorders include, but are not limited to, those cell proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders described above. In one aspect, an antibody which specifically binds MEMAP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express MEMAP.

In an additional embodiment, a vector expressing the complement of the polynucleotide

encoding MEMAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of MEMAP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of MEMAP may be produced using methods which are generally known in the art. In particular, purified MEMAP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind MEMAP. Antibodies to MEMAP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with MEMAP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to MEMAP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of MEMAP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to MEMAP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce MEMAP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for MEMAP may also be generated. For example, such fragments include, but are not limited to, $F(ab')_2$ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between MEMAP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering MEMAP epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for MEMAP. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of MEMAP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple MEMAP epitopes, represents the average affinity, or avidity, of the antibodies for MEMAP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular MEMAP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in

which the MEMAP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_d ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of MEMAP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of MEMAP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

In another embodiment of the invention, the polynucleotides encoding MEMAP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding MEMAP. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding MEMAP. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding MEMAP may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency

(e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) Science 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) Science 270:475-480; Bordignon, C. et al. (1995) Science 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) Cell 75:207-216; Crystal, R.G. et al. (1995) Hum. Gene Therapy 6:643-666; Crystal, R.G. et al. (1995) Hum. Gene Therapy 6:667-703), thalassamias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) Science 270:404-410; Verma, I.M. and N. Somia (1997) Nature 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) Nature 335:395-396; Poeschla, E. et al. (1996) Proc. Natl. Acad. Sci. USA. 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in MEMAP expression or regulation causes disease, the expression of MEMAP from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in MEMAP are treated by constructing mammalian expression vectors encoding MEMAP and introducing these vectors by mechanical means into MEMAP-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) Annu. Rev. Biochem. 62:191-217; Ivics, Z. (1997) Cell 91:501-510; Boulay, J-L. and H. Récipon (1998) Curr. Opin. Biotechnol. 9:445-450).

Expression vectors that may be effective for the expression of MEMAP include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). MEMAP may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) Proc. Natl. Acad. Sci. USA 89:5547-5551; Gossen, M. et al. (1995) Science 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) Curr. Opin. Biotechnol. 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the

FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, supra), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding MEMAP from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to MEMAP expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding MEMAP under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bonyhadi, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding MEMAP to cells which have one or more genetic abnormalities with respect to the expression of MEMAP. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have

proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) *Annu. Rev. Nutr.* 19:511-544; and Verma, I.M. and N. Somia (1997) *Nature* 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding MEMAP to target cells which have one or more genetic abnormalities with respect to the expression of MEMAP. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing MEMAP to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) *J. Virol.* 73:519-532 and Xu, H. et al. (1994) *Dev. Biol.* 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding MEMAP to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) *Curr. Opin. Biotechnol.* 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for MEMAP into the alphavirus genome in place of the capsid-coding region results in the production of a large number of MEMAP-coding RNAs and the synthesis of high levels of MEMAP in vector transduced cells. While

alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of MEMAP into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding MEMAP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding MEMAP. Such DNA sequences may be incorporated into a wide variety of

vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible
5 modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine,
10 cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding MEMAP. Compounds which may be effective in altering expression of a specific polynucleotide may include,
15 but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased
20 MEMAP expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding MEMAP may be therapeutically useful, and in the treatment of disorders associated with decreased MEMAP expression or activity, a compound which specifically promotes expression of the polynucleotide encoding MEMAP may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in
25 altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a
30 library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding MEMAP is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding MEMAP are assayed by any method commonly known in the art. Typically, the expression of a
35 specific nucleotide is detected by hybridization with a probe having a nucleotide sequence

complementary to the sequence of the polynucleotide encoding MEMAP. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruice, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruice, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of MEMAP, antibodies to MEMAP, and mimetics, agonists, antagonists, or inhibitors of MEMAP.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the

case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising MEMAP or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, MEMAP or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example MEMAP or fragments thereof, antibodies of MEMAP, and agonists, antagonists or inhibitors of MEMAP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the

active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind MEMAP may be used for the diagnosis of disorders characterized by expression of MEMAP, or in assays to monitor patients being treated with MEMAP or agonists, antagonists, or inhibitors of MEMAP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for MEMAP include methods which utilize the antibody and a label to detect MEMAP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring MEMAP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of MEMAP expression. Normal or standard values for MEMAP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to MEMAP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of MEMAP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding MEMAP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of MEMAP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess

expression of MEMAP, and to monitor regulation of MEMAP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding MEMAP or closely related molecules may be used to identify nucleic acid sequences which encode MEMAP. The specificity of the probe, whether
 5 it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding MEMAP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%
 10 sequence identity to any of the MEMAP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:38-74 or from genomic sequences including promoters, enhancers, and introns of the MEMAP gene.

Means for producing specific hybridization probes for DNAs encoding MEMAP include the cloning of polynucleotide sequences encoding MEMAP or MEMAP derivatives into vectors for the
 15 production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding MEMAP may be used for the diagnosis of disorders
 20 associated with expression of MEMAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including
 25 adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS),
 30 Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis,
 35 erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves'

disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis,

peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas. The polynucleotide sequences encoding MEMAP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR
5 technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered MEMAP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding MEMAP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide
10 sequences encoding MEMAP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding MEMAP in the
15 sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of MEMAP, a normal or standard profile for expression is established. This may be accomplished by
20 combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding MEMAP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values
25 obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from
30 successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance
35 of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals

to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding MEMAP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding MEMAP, or a fragment of a polynucleotide complementary to the polynucleotide encoding MEMAP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding MEMAP may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding MEMAP are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of MEMAP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the

polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, antibodies specific for MEMAP, or MEMAP or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and

toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, supra). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The

optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for MEMAP to quantify the levels of MEMAP expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of

protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

5 Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are
10 well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding MEMAP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may
15 be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes
20 (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism
25 (RFLP). (See, e.g., Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding MEMAP on a
30 physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse,
35 may reveal associated markers even if the exact chromosomal locus is not known. This information is

valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, MEMAP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between MEMAP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with MEMAP, or fragments thereof, and washed. Bound MEMAP is then detected by methods well known in the art. Purified MEMAP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding MEMAP specifically compete with a test compound for binding MEMAP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with MEMAP.

In additional embodiments, the nucleotide sequences which encode MEMAP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/149,641 and U.S. Ser. No. 60/164,203 are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1

ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows.

Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (PE Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (PE Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned

sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA.

The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:38-74. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the

product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding MEMAP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Chromosomal Mapping of MEMAP Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:38-74 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:38-74 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Génethon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

The genetic map locations of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:60, SEQ ID NO:63, and SEQ ID NO:67 are described in The Invention as ranges, or intervals, of human chromosomes. More than one map location is reported for SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:60, and SEQ ID NO:63, indicating that previously mapped sequences having similarity, but not complete identity, to SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:60, and SEQ ID NO:63 were assembled into their respective

clusters. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.)

5 The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Diseases associated with the public and Incyte sequences located within the indicated intervals are also reported in the Invention where applicable. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be
10 employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

VI. Extension of MEMAP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:38-74 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this
15 fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would
20 result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art: PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction
25 mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the
30 alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE
35 and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar,

Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems).

In like manner, the polynucleotide sequences of SEQ ID NO:38-74 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

VII. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:38-74 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a

SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

5 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and
10 compared.

VIII. Microarrays

 The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the
15 aforementioned technologies should be uniform and solid with a non-porous surface (Schena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to
20 those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

 Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be
25 selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser
30 desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

35 Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and

poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37 °C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85 °C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

15 Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μg. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 μl of the array element DNA, at an average concentration of 100 ng/μl, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60 °C followed by washes in

0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital

(A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

IX. Complementary Polynucleotides

Sequences complementary to the MEMAP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring MEMAP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of MEMAP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the MEMAP-encoding transcript.

X. Expression of MEMAP

Expression and purification of MEMAP is achieved using bacterial or virus-based expression systems. For expression of MEMAP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express MEMAP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of MEMAP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding MEMAP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases.

Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, MEMAP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from MEMAP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified MEMAP obtained by these methods can be used directly in the assays shown in Examples XI and XV.

XI. Demonstration of MEMAP Activity

MEMAP activity is demonstrated using a generic immunoblotting strategy or through a MEMAP-specific activity assay as outlined below. As a general approach, cell lines or tissues transformed with a vector containing MEMAP coding sequences can be assayed for MEMAP activity by immunoblotting. Transformed cells are denatured in SDS in the presence of β -mercaptoethanol, nucleic acids are removed by ethanol precipitation, and proteins are purified by acetone precipitation. Pellets are resuspended in 20 mM tris buffer at pH 7.5 and incubated with Protein G-Sepharose pre-coated with an antibody specific for MEMAP. After washing, the Sepharose beads are boiled in electrophoresis sample buffer, and the eluted proteins subjected to SDS-PAGE. Proteins are transferred from the SDS-PAGE gel to a membrane for immunoblotting, and the MEMAP activity is assessed by visualizing and quantifying bands on the blot using antibody specific for MEMAP as the primary antibody and ^{125}I -labeled IgG specific for the primary antibody as the secondary antibody.

A specific assay for MEMAP activity measures the expression of MEMAP on the cell surface. cDNA encoding MEMAP is transfected into a mammalian (non-human) cell line. Cell surface proteins are labeled with biotin as described in de la Fuente, M.A., et al. ((1997) Blood 90:2398-2405). Immunoprecipitations are performed using MEMAP-specific antibodies, and immunoprecipitated samples are analyzed using SDS-PAGE and immunoblotting techniques. The ratio of labeled immunoprecipitant to unlabeled immunoprecipitant is proportional to the amount of MEMAP expressed on the cell surface.

In an alternative specific assay, MEMAP transport activity is assayed by measuring uptake of

labeled substrates into Xenopus laevis oocytes. Oocytes at stages V and VI are injected with MEMAP mRNA (10 ng per oocyte) and incubated for 3 days at 18°C in OR2 medium (82.5mM NaCl, 2.5 mM KCl, 1mM CaCl₂, 1mM MgCl₂, 1mM Na₂HPO₄, 5 mM Hepes, 3.8 mM NaOH, 50µg/ml gentamycin, pH 7.8) to allow expression of MEMAP protein. Oocytes are then transferred to standard uptake medium (100mM NaCl, 2 mM KCl, 1mM CaCl₂, 1mM MgCl₂, 10 mM Hepes/Tris pH 7.5). Uptake of various substrates (e.g., amino acids, sugars, drugs, and neurotransmitters) is initiated by adding a ³H substrate to the oocytes. After incubating for 30 minutes, uptake is terminated by washing the oocytes three times in Na⁺-free medium, measuring the incorporated ³H, and comparing with controls. MEMAP activity is proportional to the level of internalized ³H substrate.

XII. Functional Assays

MEMAP function is assessed by expressing the sequences encoding MEMAP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10 µg of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 µg of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of MEMAP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding MEMAP and either CD64 or CD64-GFP.

CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions

of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding MEMAP and other genes of interest can be analyzed by
5 northern analysis or microarray techniques.

XIII. Production of MEMAP Specific Antibodies

MEMAP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

10 Alternatively, the MEMAP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

15 Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (PE Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-
20 MEMAP activity by, for example, binding the peptide or MEMAP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIV. Purification of Naturally Occurring MEMAP Using Specific Antibodies

Naturally occurring or recombinant MEMAP is substantially purified by immunoaffinity chromatography using antibodies specific for MEMAP. An immunoaffinity column is constructed by
25 covalently coupling anti-MEMAP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing MEMAP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of MEMAP (e.g., high ionic strength
30 buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/MEMAP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and MEMAP is collected.

XV. Identification of Molecules Which Interact with MEMAP

MEMAP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter
35 reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate

molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled MEMAP, washed, and any wells with labeled MEMAP complex are assayed. Data obtained using different concentrations of MEMAP are used to calculate values for the number, affinity, and association of MEMAP with the candidate molecules.

5 Alternatively, molecules interacting with MEMAP are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, Nature 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

 MEMAP may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions
10 between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

 Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the
15 invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	38	112301	PITUNOT01	003382R1 (HMC1NOT01), 094523R1 (PITUNOT01), 112301H1 (PITUNOT01), 301778X11 (TESTNOT04), 320551X13 (EOSIHET02), 1368852R1 (SCORNON02), 1800784H1 (COLNNOT27), 2117174X17C1 (BRSTTUT02), 2526345F6 (BRAITUT21), 4333609H1 (KIDCTMT01)
2	39	997947	KIDNTUT01	997947H1 (KIDNTUT01), 997947T6 (KIDNTUT01), 1417936X306D1 (KIDNNOT09), 1672062X307V1 (BLADNOT05), 3738956T6 (MENTNOT01), SCCA01437V1, SCCA05013V1, SCCA01691V1, SCCA02873V1
3	40	1521513	BLADTUT04	1222062H1 (NEUTGMT01), 1521513H1 (BLADTUT04), 1521513T1 (BLADTUT04), 3558522F6 (LUNGNOT31), 3558522T6 (LUNGNOT31)
4	41	1863994	PROSNOT19	265171R6 (HNT2AGT01), 1863994H1 (PROSNOT19), 3750444F6 (UTRSNOT18), 4177677F6 (BRAINOT22), 4697638F6 (BRALNOT01), 4774040F6 (BRAQNOT01), SCEA02960V1
5	42	2071941	ISLTNOT01	286350R1 (EOSIHET02), 491305R1 (HNT2AGT01), 724168R1 (SYNOOAT01), 1466668F1 (PANCUTUT02), 2071941H1 (ISLTNOT01), 2071941X11C1 (ISLTNOT01), 3579445H1 (293TF3T01)
6	43	2172512	ENDCNOT03	2172512H1 (ENDCNOT03), 2544419F6 (UTRSNOT11), 2798626H1 (NPOLNOT01), 3203359H1 (PENCNOT02), g1241299
7	44	2483172	SMCANOT01	217987F1 (STOMNOT01), 1289703F6 (BRAINOT11), 1289703T6 (BRAINOT11), 2211377F6 (SINTFET03), 2483172H1 (SMCANOT01), 2493236H1 (ADRETUT05), 3274006F6 (PROSBPT06)

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
8	45	2656128	THYMNOT04	2654722T6 (THYMNOT04), 2656128H1 (THYMNOT04), 2837168F6 (TLYMNOT03)
9	46	5855841	FIBAUNT02	894553T1 (BRSTNOT05), 1296289F1 (PGANNOT03), 1466541T1 (PANNCTUT02), 2046927F6 (THP1T7T01), 2058873R6 (OVARNOT03), 3800875F6 (SPLNNOT12), 5855841H1 (FIBAUNT02)
10	47	603462	BRSTTUT01	603462H1 (BRSTTUT01), 1487733H1 (UCMCL5T01), 1750451F6 (STOMTUT02), 5182853T6 (LUNGWT03)
11	48	747681	BRAITUT01	747681H1 (BRAITUT01), 752009R1 (BRAITUT01), 1267874F1 (BRAINOT09), 1833308R6 (BRAINON01), 2673538X19F1 (KIDNNOT19), SBCA07003F3, SCDA07521V1, SCDA04982V1, SCDA07275V1
12	49	919469	RATRN02	153337R6 (THP1PLB02), 1525415F6 (UCMCL5T01), 1527804F1 (UCMCL5T01), 1985565R6 (LUNGAST01), 2397811T6 (THP1AZT01), SARB01416F1, SARA03198F1
13	50	977658	BRSTNOT02	977658H1 (BRSTNOT02), 1873689F6 (LEUKNOT02), 2155095F6 (BRAINOT09), 2186432F6 (PROSNOT26), 2204117F6 (SPLNFET02), 2206291F6 (SPLNFET02), 3255048R6 (OVARTUN01), 3501520H1 (ADRENOT11), 3743427H1 (THYMNOT08)
14	51	1004703	BRSTNOT03	742178H1 (PANCNOT04), 1444583F6 (THYRN03), 2068902X15C1 (ISLTNOT01), 2616367F6 (GBLANOT01), SBVA02190V1
15	52	1334051	COLNNOT13	3222815T6 (COLNNON03), SXBC00794V1, SXBC00639V1
16	53	1336728	COLNNOT13	630458R6 (KIDNNOT05), 1336728H1 (COLNNOT13), SXBC00758V1, SXBC01825V1, SXBC01531V1, SXBC01624V1, SXBC00128V1

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
17	54	1452856	PENITUT01	873008R1 (LUNGAST01), 1452856H1 (PENITUT01), 2433573H1 (BRAVUNT02), 2444932F6 (THP1NOT03), 2858295F6 (SININOT03)
18	55	1562471	SPLNNOT04	286237F1 (EOSIHET02), 1562471H1 (SPLNNOT04), 1880730F6 (LEUKNOT03), 3420608F6 (UCMCNOT04), SBWA00968V1, SXBC01387V1, SBWA02301V1
19	56	1618158	BRAITUT12	967563R1 (BRSTNOT05), 1618158H1 (BRAITUT12), 1785271F6 (BRAINOT10), 2074680F6 (ISLTNOT01), 2822196H1 (ADRETUT06)
20	57	1656935	URETTUT01	1656935F6 (URETTUT01), 1656935H1 (URETTUT01), 2827605F6 (TLYMNOT03), 5272146H1 (OVARINOT02), g1482116
21	58	1859305	PROSNOT18	079372F1 (SYNORAB01), 639845R1 (BRSTNOT03), 1859305H1 (PROSNOT18), 3328091F6 (HEAONOT04), 3354812F6 (PROSNOT28), 5510642H1 (BRADDIR01)
22	59	1949083	PITUNOT01	1287161H1 (BRAINOT11), 1949083H1 (PITUNOT01), 1949083R6 (PITUNOT01), 1949083T6 (PITUNOT01), 3814131F6 (TONSNOT03)
23	60	1996357	BRSTTUT03	260527R6 (HNT2RAT01), 260527T6 (HNT2RAT01), 1313441F1 (BLADTUT02), 1442781R1 (THYRNOT03), 1996357H1 (BRSTTUT03), 1996357T6 (BRSTTUT03), 4262451H1 (BSCNDIT02), SAZA00147F1
24	61	2061330	OVARNOT03	2061330H1 (OVARNOT03), 2724233T6 (LUNGUTUT10), 5104031T6 (PROSTUS20)
25	62	2346947	TESTTUT02	2346947F6 (TESTTUT02), 2346947H1 (TESTTUT02), 4051345F6 (SINTNOT18)

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
26	63	2795577	NPOLNOT01	867213R6 (BRAITUT03), 2381770H1 (ISLTNOT01), 2795577CT1 (NPOLNOT01), 2795577H1 (NPOLNOT01)
27	64	3255825	OVARTUN01	3255825CT1 (OVARTUN01), 3255825H1 (OVARTUN01)
28	65	3393430	LUNGNOT28	2187169H1 (PROSNOT26), 3393256H1 (LUNGNOT28), 3393430H1 (LUNGNOT28), 3395774H1 (LUNGNOT28), 4689688H1 (LIVRTUT12), 4895996H1 (LIVRTUT12), 4896461F6 (LIVRTUT12), 4984527F6 (LIVRTUT10), 4992946H1 (LIVRTUT11)
29	66	3490990	EPIGNOT01	1235428F1 (LUNGFET03), 1662973T6 (BRSTNOT09), 2362021H1 (LUNGFET05), 2362021R6 (LUNGFET05), 3490990H1 (EPIGNOT01)
30	67	3635154	LIVRNOT03	027592H1 (SPLNFET01), 3635154H1 (LIVRNOT03), g1012932
31	68	4374347	CONFNOT03	860875X11 (BRAITUT03), 898143R6 (BRSTNOT05), 4374347H1 (CONFNOT03)
32	69	4596747	COLSTUT01	137213R1 (SYNORAB01), 545568R6 (OVARNOT02), 1235402F1 (LUNGFET03), 1268010F1 (BRAINOT09), 1271078F1 (TESTTUT02), 1301951F6 (BRSTNOT07), 1994442R6 (BRSTTUT03), 2343102H1 (TESTTUT02), 3274538F6 (PROSBPT06), 4596747H1 (COLSTUT01)
33	70	5052680	BRSTNOT33	1973688H1 (UCMCL5T01), 3926410F6 (KIDNNOT19), 4501839F6 (BRAVXT02), 5052680F6 (BRSTNOT33), 5052680H1 (BRSTNOT33), 5186780F6 (LUNGTM04)
34	71	5373575	BRAINOT22	262776T6 (HNT2AGT01), 1234057F1 (LUNGFET03), 1741526R6 (HIPONON01), 1871204F6 (SKINBIT01), 2192479F6 (THYRTUT03), 2556849H1 (THYMNOT03), 2722451T6 (LUNGFTUT10), 4114985H1 (UTRSTUT07), 5373575H1 (BRAINOT22)

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
35	72	5524468	LIVRDIR01	4024068F6 (BRAXNOT02), 5524468H1 (LIVRDIR01), SXBC01952V1
36	73	5944279	COLADIT05	1662182H1 (BRSTNOT09), 1698677F6 (BLADTUT05), 1916639R6 (PROSNOT06), 1916639T6 (PROSNOT06), 2298565R6 (BRSTNOT05), 2298565T6 (BRSTNOT05), 2583019F6 (KIDNTUT13), 2870903F6 (THYRNOT10), 3970715H1 (PROSTUT10), 3971695H1 (PROSTUT10), 5944279H1 (COLADIT05)
37	74	6114480	SINITMT04	1579843F6 (DUODNOT01), 1579843T6 (DUODNOT01), 4181024T6 (SINITUT03), 6114480H1 (SINITMT04), SXBC00007V1, SXBC00504V1, SCSA05104V1

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
1	351	S31 T116 S169 T229 T2 S209 T306	N128	Signal peptide: M1-A33	Paraneoplastic neuronal antigen MA1 [Homo sapiens] g4104634	BLAST-GenBank MOTIFS SPSCAN
2	458	T198 S27 S37 T87 S251 S257 S325 S373 S405 S422 T454 T210 S228 S401 Y93	N75 N159 N279 N445	Signal peptide: M1-T24 Glycoprotein signature: C199-L448	Pancortin-3 [Mus musculus] g3218528	BLAST-GenBank MOTIFS SPSCAN HMMER BLAST-PRODOM
3	219	T51 S120 S163 T175 T181 S3 T12 T45 S75 S104 S128	N2 N62 N107	Signal peptide: M1-C42 Transmembrane domain: L32-F49 C-type lectin domain: C80-E206	Murine macrophage C- type lectin [Mus musculus] g5821286	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLIMPS-BLOCKS PROFILES SCAN BLAST-DOMO
4	276	S213 S91 S113 S35 S70 S76 S147 T163 S206		Signal peptide: M1-G31 Transmembrane domain: I184-F201 Cell attachment sequence: R149-D151		BLAST-GenBank MOTIFS HMMER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
5	375	S18 S205 T286 S3 S120 S197 T260 Y85		Transmembrane domains: W139-R158; F173-H191 P232-Q254 Transmembrane protein signature: I95-C369	Transmembrane protein [S. pombe] g1065898	BLAST-GenBank MOTIFS HMMER BLAST-DBO BLAST-PRODOR
6	249	T7 T135 T170 S204 Y154	N18 N92 N147		Phospholipid scramblase [Homo sapiens] g4092081	BLAST-GenBank MOTIFS
7	353	T162 T4 S97 T115 S165 S194 T225 S242 S17 S47 S205	N299	Signal peptide: M1-A33	Paraneoplastic neuronal antigen MA1 [Homo sapiens] g4104634	BLAST-GenBank MOTIFS SPSCAN
8	194	T12 S115 S29 S99 S187	N95 N147	Signal peptide: M1-C50 Transmembrane domain: L38-L56 C-type lectin domain: C75-E194	Lectin-like NK cell receptor L1T1 [Homo sapiens] g6651065	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLIMPS-BLOCKS BLAST-DBO
9	322	S304 S48 S146 S72 T133 S255 S280	N20 N60 N70	Signal peptide: M1-A50		BLAST-GenBank MOTIFS SPSCAN

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
10	335	S125 S140 S183 S222 T252		Transmembrane domains: G71-L94; A255-I283 Gufa transmembrane protein domain: L12-H101; G180-G335 Glycosaminoglycan attachment site: S310-G313	Gufa protein [Thermotoga maritima] g4982315	BLAST-GenBank MOTIFS HMMER BLAST-PRODOM BLAST-DOMO
11	620	S49 S108 T146 S300 T348 T349 S607 S4 S128 S183 S234 T420 S460 S467 S543 Y597	N144 N202 N264 N274 N293 N341 N492 N505 N526 N542	Transmembrane domain: M563-W582 Immunoglobulin domain: G439-A499 Leucine-rich repeat signature: L337-L350 Glycoprotein hormone receptor domain: T40-L198	Slit2 [Rattus norvegicus] g4585574	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-PRINTS BLAST-DOMO

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
12	491	T231 T232 S253 T482 S185 S276	N56 N220 N229	Transmembrane domains: I115-I142; I184-V201 F422-F441 Transmembrane protein domain: L8-Y215; I396-F471	Selectively expressed in embryonic epithelia protein-1 [Mus musculus] g6715148 PB39 [Homo sapiens] g3462515	BLAST-GenBank MOTIFS HMMER BLAST-PRODOM
13	580	S557 S10 T34 S51 T92 T210 S343 T12 S217 T222 S268 S296 T417 T523 S550	N159 N179 N220 N230	Transmembrane domains: F297-F313; I356-I373 L496-I514 Lipases serine active site: L104-A113		MOTIFS HMMER
14	455	T53 T182 S239 S69 S135 S202 T280 S355 S372 Y38	N67 N180 N243	Transmembrane domains: V81-V99; I343-I361 S375-V392; W425-Y442 Glycosaminoglycan attachment site: S162-G165	putative G- protein coupled receptor [Homo sapiens] g6649579	BLAST-GenBank MOTIFS HMMER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
15	277	S265 T66 T225 S268 S273 S30 S49 S61 S152 S193 Y242	N29 N38 N47 N48 N92 N160 N210	Transmembrane domain: K9-F27 Brush border protein domain: Y8-R277 RGD cell attachment sequence: R113-D115	AdRab-A brush border membrane protein [Oryctolagus cuniculus] g1762	BLAST-GenBank MOTIFS HMMER BLAST-PRODOM
16	647	S490 T50 S67 S105 T110 S121 T220 S249 S264 S272 S322 T389 S469 T501 S639 S132 T155 S242 S324 T381 T400 S522 S554	N261	Signal peptide: M1-A22 Transmembrane domains: L328-L347; M406-L424 L559-A578; W618-L638 GufA transmembrane protein domain: E485-L640 Glycosaminoglycan attachment site: S34-G37	LIV-1 protein [Homo sapiens] g1256001	BLAST-GenBank MOTIFS SPSCAN HMMER BLAST-PRODOM
17	406	S29 S215 S236 T69	N23	Transmembrane domains: Q76-V95; W286-S313 M367-I384		MOTIFS HMMER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
18	290	T221 S44 S69 S71 S81 T94 T101 T113 T131 S216 Y284	N88	Signal peptide: M1-A19 Transmembrane domains: P160-M181 Immunoglobulin domain: R33-I110 Transmembrane glycoprotein domain: I22-D116	NK inhibitory receptor [Homo sapiens] g6707799 CMRF-35-H9 leukocyte antigen [Homo sapiens] g4103066	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLAST-PRODOM BLAST-DOMO
19	390	S7 T68 S153 T23 T166 T281 Y20 Y37	N5 N88 N330 N367	Immunoglobulins and MHC proteins signature: T90-P112; F242-V259 Glycoprotein antigen signature: L61-V72; V92-I113		MOTIFS BLIMPS-BLOCKS BLIMPS-PRODOM
20	427	S13 S41 S65 S66 S99 T150 S323 S324 S101 S275 S353 S367 T399 Y71	N106 N148 N171 N233 N312	Mucin glycoprotein precursor domain: V136-P142	Gastric mucin [Sus scrofa] g915208	BLAST-GenBank MOTIFS BLIMPS-PRODOM

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
21	459	T4 S60 S66 S116 T176 S16 T235	N14 N158 N323	Transmembrane domains: F202-V219; I246-L268 W343-L367; P417-P440	six transmembrane epithelial antigen of prostate [Homo sapiens] g6572948	BLAST-GenBank MOTIFS HMMER
22	229	S13 S118 T155 Y24		Transmembrane domains: I93-V111; V132-L150 F164-V182 Transmembrane protein domain: S156-V182		MOTIFS HMMER BLIMPS-PRODOM
23	311	S85 S234 S236 S269 S80 S119 S186 T294	N22	Transmembrane domains: W58-I76; P152-K177 A216-Y232		MOTIFS HMMER
24	92	S47 T54 T12 S70	N62		HERV-E envelope glycoprotein [Homo sapiens] g2587024	BLAST-GenBank MOTIFS
25	258	S34 T33 S148 S243		Transmembrane domains: I39-I57; F86-L106 V122-I140; L190-S210		MOTIFS HMMER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
26	226	S56 S128 T196 T167 Y194	N54 N187 N198	Signal peptide: M1-P50 Transmembrane domains: T23-L43; M72-A89 I101-I124; I158-N178 Transmembrane 4 family signature: A70-I120 Lysosomal-associated transmembrane protein domain: C15-Y223	MTP (mouse transporter protein) [Mus musculus] g1276631	BLAST-GenBank MOTIFS SPSCAN HMMER PROFILESCAN BLAST-PRODOM
27	136	S3 S132		Signal peptide: M1-R53 Transmembrane domains: I10-L28; T26-I50 F70-L89 Transmembrane protein domain: D31-V104		MOTIFS SPSCAN HMMER BLAST-PRODOM

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
28	458	T408 T98 S126 S170 T334	N96 N151 N293 N332	Signal peptide: M1-A20 Transmembrane domain: L10-N30 Membrane glycoprotein signature: L9-V101; L64-Q457 Olfactory ligand binding domain: T67-S452	Potential ligand (odorant) binding protein [Rattus rattus] g57732	BLAST-GenBank MOTIFS SPSCAN HMMER BLAST-PRODOM BLAST-DOMO
29	368	S24 T166 T302 S12 S134 Y307	N17		Fuzzy (TM protein involved in tissue polarity) [Drosophila melanogaster] g2564657	BLAST-GenBank MOTIFS
30	91	T44 S84		Signal peptide: M1-A19 Transmembrane domain: P58-S82 Glycophorin A proteins signature: T22-S32; I63-G91 Glycophorin domain: M1-R86	Preglycophorin B [Homo sapiens] g4803699	BLAST-GenBank MOTIFS SPSCAN HMMER BLINPS-BLOCKS PROFILES SCAN BLAST-PRODOM BLAST-DOMO

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
31	295	S96 T113 S129 T155 T125 T157 T187 S222 T231 T263 Y212	N111 N169 N223	Signal peptide: M1-G48 Transmembrane domain: L241-L259 Immunoglobulin domain: K159-V216 Carcinoembryonic antigen domain: I38-P147 Glycoprotein antigen domain: M1-V140; Y141-Y234 G239-S295	Biliary glycoprotein [Mus musculus] g312590	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLAST-PRODOM BLAST-DOMO
32	724	T39 S47 T171 S205 T224 S225 T241 S285 S301 T323 S352 T353 S439 S509 S517 S537 T659 T707 S8 S18 S49 S72 T85 T159 S173 S271 S367 S560 S588 Y499	N279 N348	Transmembrane domain: I611-F630 Membrane protein domain: T4-L209		MOTIFS HMMER BLAST-DOMO

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
33	331	S117 S147 S149 T320 S138 S174 T274 T319 S328 Y198	N222	Signal peptide: M1-S16 Transmembrane domains: A67-N87; I118-C134 W240-V269; L294-Y310 Transmembrane protein domain: A6-T311	Putative Golgi UDP-GlcNAc transporter [S. pombe] g3738167	BLAST-GenBank MOTIFS SPSCAN HMMER BLAST-PRODOM
34	398	T42 T158 S271 S28 S285 T334 S375		Transmembrane domain: I59-L79 Band 7 family domain: F64-A231, A78-V90; R116-L154 Stomatin signature: T84-L106; L131-P152 T166-L183; I186-G209 L54-Q227	Stomatin-like protein UNC24 [Homo sapiens] g5326747	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO
35	220	S199 T120 S192	N107	Signal peptide: M1-G19 Leucine rich repeats: A62-F85; Q86-S109 G110-G133; A134-R157 A158-S181; H184-P207	Similar to Leucine-rich transmembrane proteins [Homo sapiens] g2781386	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLIMPS-PRINTS

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
36	706	T564 T74 T113 S291 S452 S632 S14 T42 S66 T115 T142 S286 T551 T575 S701	N101	Transmembrane domains: F158-M178; L344-V368 L425-L442; M478-F498 A581-I604; L641-V665 Glycosaminoglycan attachment site: S223-G226	LAK-4p [Homo sapiens] g7209574	BLAST-GenBank MOTIFS HMMER
37	466	T326 S10 T46 T105 S187 S98 T164 T310 S321 Y388	N368	Signal peptide: M1-G23 Transmembrane domain: A236-I255 SPRY domain: A338-S464; E123-S136 E322-W343; V407-F420 Butyrophilin domain: W19-C114	Butyrophilin like receptor [Homo sapiens] g4587209	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLIMPS-PFAM BLAST-PRODOM BLAST-DOMO

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
38	844-888	Nervous (0.377) Reproductive (0.180) Cardiovascular (0.115) Gastrointestinal (0.115)	Cancer (0.410) Inflammation/Trauma (0.296) Cell Proliferation (0.131)	PBLUESCRIPT
39	579-623	Developmental (0.400) Musculoskeletal (0.200) Nervous (0.200) Urologic (0.200)	Cancer (0.400) Cell Proliferation (0.400)	PSPORT1
40	336-380	Cardiovascular (0.267) Hematopoietic/Immune (0.200) Endocrine (0.133) Reproductive (0.133)	Cancer (0.400) Inflammation/Trauma (0.400) Cell Proliferation (0.133)	pINCY
41	596-640	Nervous (0.588) Gastrointestinal (0.118) Reproductive (0.118)	Inflammation/Trauma (0.470) Cancer (0.235) Cell Proliferation (0.176)	pINCY
42	1281-1325	Reproductive (0.237) Hematopoietic/Immune (0.145) Nervous (0.145)	Cancer (0.441) Inflammation/Trauma (0.323) Cell Proliferation (0.178)	pINCY
43	227-271	Reproductive (0.444) Dermatologic (0.222) Endocrine (0.111) Gastrointestinal (0.111) Nervous (0.111)	Cancer (0.333) Cell Proliferation (0.222) Inflammation/Trauma (0.222)	pINCY

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
44	1368-1412	Nervous (0.339) Reproductive (0.278) Gastrointestinal (0.104)	Cancer (0.478) Inflammation/Trauma (0.278) Cell Proliferation (0.165)	pINCY
45	543-587	Hematopoietic/Immune (0.500) Gastrointestinal (0.188)	Inflammation/Trauma (0.500) Cancer (0.250) Cell Proliferation (0.188)	pINCY
46	280-324	Reproductive (0.267) Nervous (0.233) Gastrointestinal (0.112)	Cancer (0.483) Inflammation/Trauma (0.345) Cell Proliferation (0.155)	pINCY
47	380-424 875-919	Reproductive (0.412) Gastrointestinal (0.176) Cardiovascular (0.118)	Cancer (0.647) Inflammation/Trauma (0.178)	PSPORT1
48	272-316 1514-1558	Nervous (0.645) Developmental (0.129)	Cancer (0.355) Cell Proliferation (0.258) Neurological (0.194)	PSPORT1
49	282-326 768-812	Hematopoietic/Immune (0.238) Gastrointestinal (0.155) Reproductive (0.143)	Cancer (0.381) Inflammation/Trauma (0.381) Cell Proliferation (0.202)	PSPORT1
50	597-641 1074-1118	Reproductive (0.214) Nervous (0.196) Hematopoietic/Immune (0.143)	Cancer (0.464) Inflammation/Trauma (0.304) Cell Proliferation (0.196)	PSPORT1
51	973-1017	Reproductive (0.266) Nervous (0.234) Hematopoietic/Immune (0.125)	Cancer (0.516) Inflammation/Trauma (0.359) Cell Proliferation (0.109)	PSPORT1

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
52	299-343	Gastrointestinal (1.000)	Cancer (0.500) Inflammation/Trauma (0.500)	pINCY
53	380-424 1199-1243	Gastrointestinal (0.289) Reproductive (0.244) Cardiovascular (0.111) Hematopoietic/Immune (0.111)	Cancer (0.578) Inflammation/Trauma (0.311) Cell Proliferation (0.178)	pINCY
54	1135-1179	Nervous (0.195) Reproductive (0.186) Gastrointestinal (0.144)	Cancer (0.449) Inflammation/Trauma (0.305) Cell Proliferation (0.144)	pINCY
55	325-369 820-864	Hematopoietic/Immune (0.750)	Inflammation/Trauma (0.625) Cancer (0.125)	pINCY
56	487-531 1090-1134	Nervous (0.583)	Cancer (0.458) Inflammation/Trauma (0.250)	pINCY
57	569-613 1360-1405	Reproductive (0.429) Hematopoietic/Immune (0.286) Musculoskeletal (0.143) Urologic (0.143)	Cancer (0.571) Inflammation/Trauma (0.286) Cell Proliferation (0.143)	pINCY
58	272-472 551-595 812-1012 1523-1567	Reproductive (0.350) Nervous (0.150) Cardiovascular (0.100) Gastrointestinal (0.100) Hematopoietic/Immune (0.100) Urologic (0.100)	Cancer (0.500) Inflammation/Trauma (0.500)	pINCY

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
59	217-261	Nervous (0.286) Developmental (0.143) Gastrointestinal (0.143) Hematopoietic/Immune (0.143) Reproductive (0.143)	Inflammation/Trauma (0.428) Cancer (0.357) Cell Proliferation (0.143)	PBLUESCRIPT
60	444-488	Nervous (0.207) Reproductive (0.207) Gastrointestinal (0.130) Hematopoietic/Immune (0.130)	Cancer (0.467) Inflammation/Trauma (0.359) Cell Proliferation (0.163)	PSPORT1
61	643-687	Reproductive (0.464) Endocrine (0.143) Cardiovascular (0.107) Gastrointestinal (0.107)	Cancer (0.500) Inflammation/Trauma (0.321)	PSPORT1
62	146-344 390-434 506-704 786-830	Gastrointestinal (0.500) Hematopoietic/Immune (0.250) Reproductive (0.250)	Cancer (0.750) Inflammation/Trauma (0.250)	pINCY
63	163-207	Reproductive (0.315) Gastrointestinal (0.161) Cardiovascular (0.147)	Cancer (0.594) Cell Proliferation (0.231) Inflammation/Trauma (0.210)	pINCY
64	201-506 525-569 606-912 975-1280 1362-1406	Gastrointestinal (0.455) Cardiovascular (0.273) Reproductive (0.189)	Cancer (0.455) Inflammation/Trauma (0.367) Cell Proliferation (0.189)	PSPORT1

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
65	703-747	Gastrointestinal (0.667) Cardiovascular (0.167) Reproductive (0.167)	Cancer (1.000)	pINCY
66	271-315 319-363	Nervous (0.314) Reproductive (0.314) Developmental (0.114) Urologic (0.114)	Cancer (0.429) Cell Proliferation (0.171) Inflammation/Trauma (0.143)	pINCY
67	319-363	Developmental (0.364) Hematopoietic/Immune (0.364) Gastrointestinal (0.182)	Cell Proliferation (0.727) Cancer (0.273) Inflammation/Trauma (0.182)	pINCY
68	812-856	Reproductive (0.444) Nervous (0.222) Endocrine (0.111) Hematopoietic/Immune (0.111) Musculoskeletal (0.111)	Cancer (0.556) Inflammation/Trauma (0.333)	pINCY
69	596-640 1577-1621	Reproductive (0.255) Nervous (0.184) Developmental (0.122) Gastrointestinal (0.122)	Cancer (0.429) Inflammation/Trauma (0.337) Cell Proliferation (0.255)	pINCY
70	379-675 703-747 766-1062 1081-1347	Nervous (0.467) Hematopoietic/Immune (0.200) Reproductive (0.133) Urologic (0.133)	Cancer (0.467) Cell Proliferation (0.267) Inflammation/Trauma (0.267)	pINCY

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
71	18-62	Nervous (0.265) Reproductive (0.206) Musculoskeletal (0.147)	Cancer (0.500) Inflammation/Trauma (0.264) Cell Proliferation (0.147)	pINCY
72	290-488 507-704 759-803	Gastrointestinal (0.333) Hematopoietic/Immune (0.333) Nervous (0.333)	Inflammation/Trauma (0.667) Cancer (0.333)	pINCY
73	649-693 1711-1755	Reproductive (0.392) Gastrointestinal (0.294) Cardiovascular (0.118)	Cancer (0.686) Inflammation/Trauma (0.294)	pINCY
74	704-748	Gastrointestinal (0.923)	Cancer (0.462) Inflammation/Trauma (0.385)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
38	PITUNOT01	This library was constructed using RNA obtained from Clontech (CLON 6584-2, lot 35278). The RNA was isolated from pituitary glands removed from a pool of 18 male and female Caucasian donors, 16 to 70 years old, who died from trauma.
39	KIDNTUT01	This library was constructed using RNA isolated from kidney tumor tissue removed from an 8-month-old female during nephroureterectomy. Pathology indicated Wilms' tumor (nephroblastoma), which involved 90 percent of the renal parenchyma. Prior to surgery, the patient was receiving heparin anticoagulant therapy.
40	BLADTUT04	This library was constructed using RNA isolated from bladder tumor tissue removed from a 60-year-old Caucasian male during a radical cystectomy, prostatectomy, and vasectomy. Pathology indicated grade 3 transitional cell carcinoma in the left bladder wall. Carcinoma in-situ was identified in the dome and trigone. Patient history included tobacco use. Family history included type I diabetes, malignant neoplasm of the stomach, atherosclerotic coronary artery disease, and acute myocardial infarction.
41	PROSNOT19	This library was constructed using RNA isolated from diseased prostate tissue removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+3). The patient presented with elevated prostate-specific antigen (PSA). Patient history included colon diverticuli, asbestosis, and thrombophlebitis. Family history included benign hypertension, multiple myeloma, hyperlipidemia and rheumatoid arthritis.
42	ISLTNOT01	This library was constructed using RNA isolated from a pooled collection of pancreatic islet cells.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
43	ENDCNOT03	This library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male.
44	SMCANOT01	This library was constructed using RNA isolated from an aortic smooth muscle cell line derived from the explanted heart of a male obtained during a heart transplant.
45	THYMNOT04	This library was constructed using RNA isolated from thymus tissue removed from a 3-year-old Caucasian male, who died from anoxia.
46	FIBAUNT02	This library was constructed using RNA isolated from untreated aortic adventitial fibroblasts removed from a 65-year-old Caucasian female.
47	BRSTTUT01	This library was constructed using RNA isolated from breast tumor tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 4 mammary adenocarcinoma. Patient history included atrial tachycardia and a benign breast neoplasm. Family history included cardiovascular and cerebrovascular disease and depressive disorder.
48	BRAITUT01	This library was constructed using RNA isolated from brain tumor tissue removed from a 50-year-old Caucasian female during a frontal lobectomy. Pathology indicated recurrent grade 3 oligoastrocytoma with focal necrosis and extensive calcification. Patient history included a speech disturbance and epilepsy. The patient's brain had also been irradiated with a total dose of 5,082 cys (Fraction 8). Family history included a brain tumor.
49	RATRNOT02	This library was constructed using RNA isolated from the right atrium tissue of a 39-year-old Caucasian male, who died from a gunshot wound.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
50	BRSTNOT02	This library was constructed using RNA isolated from diseased breast tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated proliferative fibrocystic changes characterized by apocrine metaplasia, sclerosing adenosis, cyst formation, and ductal hyperplasia without atypia. Pathology for the associated tumor tissue indicated an invasive grade 4 mammary adenocarcinoma. Patient history included atrial tachycardia and a benign neoplasm. Family history included cardiovascular and cerebrovascular disease.
51	BRSTNOT03	This library was constructed using RNA isolated from diseased breast tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy. Pathology for the associated tumor tissue indicated residual invasive grade 3 mammary ductal adenocarcinoma. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia and a malignant neoplasm of the colon.
52	COLNNOT13	This library was constructed using RNA isolated from ascending colon tissue of a 28-year-old Caucasian male with moderate chronic ulcerative colitis.
53	COLNNOT13	This library was constructed using RNA isolated from ascending colon tissue of a 28-year-old Caucasian male with moderate chronic ulcerative colitis.
54	PENITUT01	This library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old Caucasian male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
55	SPLNNOT04	This library was constructed using RNA isolated from the spleen tissue of a 2-year-old Hispanic male, who died from cerebral anoxia.
56	BRAITUT12	This library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 40-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated grade 4 gemistocytic astrocytoma.
57	URETTUT01	This library was constructed using RNA isolated from right ureter tumor tissue of a 69-year-old Caucasian male during ureterectomy and lymph node excision. Pathology indicated invasive grade 3 transitional cell carcinoma. Patient history included benign colon neoplasm, tobacco use, asthma, emphysema, acute duodenal ulcer, and hyperplasia of the prostate. Family history included atherosclerotic coronary artery disease, congestive heart failure, and malignant lung neoplasm.
58	PROSNOT18	This library was constructed using RNA isolated from diseased prostate tissue removed from a 58-year-old Caucasian male during a radical cystectomy, radical prostatectomy, and gastrectomy. Pathology indicated adenofibromatous hyperplasia; this tissue was associated with a grade 3 transitional cell carcinoma. Patient history included angina and emphysema. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
59	PITUNOT01	This library was constructed using RNA obtained from Clontech (CLON 6584-2, lot 35278). The RNA was isolated from the pituitary glands removed from a pool of 18 male and female Caucasian donors, 16 to 70 years old, who died from trauma.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
60	BRSTTUT03	This library was constructed using RNA isolated from breast tumor tissue removed from a 58-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated multicentric invasive grade 4 lobular carcinoma. The mass was identified in the upper outer quadrant, and three separate nodules were found in the lower outer quadrant of the left breast. Patient history included skin cancer, rheumatic heart disease, osteoarthritis, and tuberculosis. Family history included cerebrovascular disease, coronary artery aneurysm, breast cancer, prostate cancer, atherosclerotic coronary artery disease, and type I diabetes.
61	OVARNOT03	This library was constructed using RNA isolated from ovarian tissue removed from a 43-year-old Caucasian female during removal of the fallopian tubes and ovaries. Pathology for the associated tumor tissue indicated grade 2 mucinous cystadenocarcinoma. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, pancreatic cancer, stress reaction, cerebrovascular disease, breast cancer, and uterine cancer.
62	TESTTUT02	This library was constructed using RNA isolated from testicular tumor tissue removed from a 31-year-old Caucasian male during unilateral orchiectomy. Pathology indicated embryonal carcinoma.
63	NPOLNOT01	This library was constructed using RNA isolated from nasal polyp tissue removed from a 78-year-old Caucasian male during a nasal polypectomy. Pathology indicated a nasal polyp and striking eosinophilia. Patient history included asthma and nasal polyps.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
64	OVARTUN01	This normalized library was constructed from 5.36 million independent clones obtained from an ovarian tumor library. RNA was isolated from tumor tissue removed from the left ovary of a 58-year-old Caucasian female during a total abdominal hysterectomy, removal of a single ovary, and inguinal hernia repair. Pathology indicated a metastatic grade 3 adenocarcinoma of colonic origin, forming a partially cystic and necrotic tumor mass in the left ovary, and a nodule in the left mesovarium. A single intramural leiomyoma was identified in the myometrium. The cervix showed mild chronic cystic cervicitis. Patient history included benign hypertension, follicular ovarian cyst, colon cancer, benign colon neoplasm, and osteoarthritis. Family history included emphysema, myocardial infarction, atherosclerotic coronary artery disease, benign hypertension, hyperlipidemia, and primary tuberculous complex. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9228) and Bonaldo et al. (Genome Research (1996) 6:791).
65	LUNGNOT28	This library was constructed using RNA isolated from lung tissue removed from a 53-year-old male. Pathology for the associated tumor tissue indicated grade 4 adenocarcinoma.
66	EPIGNOT01	This library was constructed using RNA isolated from epiglottic tissue removed from a 71-year-old male during laryngectomy with right parathyroid biopsy. Pathology for the associated tumor tissue indicated recurrent grade 1 papillary thyroid carcinoma.
67	LIVRNOT03	This library was constructed using RNA isolated from liver tissue removed from a Caucasian male fetus, who died from Patau's syndrome (trisomy 13) at 20 weeks' gestation.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
68	CONFNOT03	This library was constructed using RNA isolated from mesenteric fat tissue removed from a 71-year-old Caucasian male during a partial colectomy and permanent colostomy. Pathology indicated mesenteric fat tissue associated with diverticulosis and diverticulitis with abscess formation. Approximately 50 diverticula were noted, one of which was perforated and associated with abscess formation in adjacent mesenteric fat. The patient presented with atrial fibrillation. Patient history included viral hepatitis, a hemangioma, and diverticulitis of colon. Family history included extrinsic asthma, atherosclerotic coronary artery disease, and myocardial infarction.
69	COLSTUT01	This library was constructed using RNA isolated from colon tumor tissue removed from the sigmoid colon of a 62-year-old Caucasian male during a sigmoidectomy and permanent colostomy. Pathology indicated invasive grade 2 adenocarcinoma, with invasion through the muscularis. Patient history included hyperlipidemia, cataract disorder and dermatitis. Family history included benign hypertension, atherosclerotic coronary artery disease, hyperlipidemia, breast cancer, and prostate cancer.
70	BRSTNOT33	This library was constructed using RNA isolated from right breast tissue removed from a 46-year-old Caucasian female during a unilateral extended simple mastectomy with breast reconstruction. Pathology for the associated tumor tissue indicated invasive grade 3 adenocarcinoma, ductal type, with apocrine features, nuclear grade 3 forming a mass in the outer quadrant. There was greater than 50% intraductal component. Patient history included breast cancer.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
71	BRAINOT22	This library was constructed using RNA isolated from right temporal lobe tissue removed from a 45-year-old Black male during a brain lobectomy. Pathology for the associated tumor tissue indicated dysembryoplastic neuroepithelial tumor of the right temporal lobe. The right temporal region dura was consistent with calcifying pseudotumor of the neuraxis. Patient history included obesity, meningitis, backache, unspecified sleep apnea, acute stress reaction, acquired knee deformity, and chronic sinusitis. Family history included obesity, benign hypertension, cirrhosis of the liver, obesity, hyperlipidemia, cerebrovascular disease, and type II diabetes.
72	LIVRDIR01	This library was constructed using RNA isolated from diseased liver tissue removed from a 63-year-old Caucasian female during a liver transplant. Patient history included primary biliary cirrhosis. Serology was positive for anti-mitochondrial antibody.
73	COLADIT05	This library was constructed using RNA isolated from diseased ascending colon tissue removed from a 32-year-old Caucasian male during a total intra-abdominal colectomy, abdominal-perineal rectal resection, and temporary ileostomy. Pathology indicated chronic ulcerative colitis extending in a continuous fashion from the mid-portion of the ascending colon distally to the rectum. This was characterized microscopically by crypt abscess formation and inflammation confined to the mucosa and submucosa. The terminal ileum exhibited ileitis and the rectal mucosa showed crypt abscess formation. Patient history included tobacco use. Family history included ulcerative colitis, malignant neoplasm of the breast and acute myocardial infarction.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
74	SINITMT04	<p>Library was constructed using RNA isolated from ileum tissue removed from a 70-year-old Caucasian female during right hemicolectomy, open liver biopsy, flexible sigmoidoscopy, colonoscopy, and permanent colostomy. Pathology indicated a non-tumorous margin of ileum. Pathology for the associated tumor indicated invasive grade 2 adenocarcinoma forming an ulcerated mass, situated 2 cm distal to the ileocecal valve. The tumor invaded through the muscularis propria just into the serosal adipose tissue. One (of 16) regional lymph node was positive for a microfocus of metastatic adenocarcinoma. Patient history included a malignant breast neoplasm, type II diabetes, hyperlipidemia, viral hepatitis, an unspecified thyroid disorder, osteoarthritis, and a malignant skin neoplasm. Family history included breast cancer, atherosclerotic coronary artery disease, benign hypertension, cerebrovascular disease, ovarian cancer, and hyperlipidemia.</p>

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	PE Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	PE Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	PE Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.0E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score > GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29,
 - 10 SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37,
- b) a naturally occurring amino acid sequence having at least 70% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10,
 - 15 SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37,
- 20 c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID
 - 25 NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, and
- d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID
 - 30 NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37.

35 2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1,

SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37.

3. An isolated polynucleotide encoding a polypeptide of claim 1.

10 4. An isolated polynucleotide encoding a polypeptide of claim 2.

5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74.

20 6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3..

7. A cell transformed with a recombinant polynucleotide of claim 6.

25 8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

9. A method for producing a polypeptide of claim 1, the method comprising:

a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

35

11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74,
- b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74,
- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37.

18. A method for treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide

of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

5 23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional MEMAP, comprising administering to a patient in need of such treatment a composition of claim 23.

10 25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- 15 b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- 20 a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound
- 25 with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- 30 a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

35 28. A method for assessing toxicity of a test compound, said method comprising:

a) treating a biological sample containing nucleic acids with the test compound;

b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;

c) quantifying the amount of hybridization complex; and

d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

SEQUENCE LISTING

<110> INCYTE GENOMICS, INC.

LAL, Preeti

YUE, Henry

TANG, Y. Tom

BANDMAN, Olga

BURFORD, Neil

AZIMZAI, Yalda

BAUGHN, Mariah R.

LU, Dyung Aina M.

PATTERSON, Chandra

<120> MEMBRANE ASSOCIATED PROTEINS

<130> PF-0731 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/149,641; 60/164,203

<151> 1999-08-17; 1999-11-09

<160> 74

<170> PERL Program

<210> 1

<211> 351

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 112301CD1

<400> 1

Met	Thr	Leu	Arg	Leu	Leu	Glu	Asp	Trp	Cys	Arg	Gly	Met	Asp	Met	
1				5					10					15	
Asn	Pro	Arg	Lys	Ala	Leu	Leu	Ile	Ala	Gly	Ile	Ser	Gln	Ser	Cys	
				20					25					30	
Ser	Val	Ala	Glu	Ile	Glu	Glu	Ala	Leu	Gln	Ala	Gly	Leu	Ala	Pro	
				35					40					45	
Leu	Gly	Glu	Tyr	Arg	Leu	Leu	Gly	Arg	Met	Phe	Arg	Arg	Asp	Glu	
				50					55					60	
Asn	Arg	Lys	Val	Ala	Leu	Val	Gly	Leu	Thr	Ala	Glu	Thr	Ser	His	
				65					70					75	
Ala	Leu	Val	Pro	Lys	Glu	Ile	Pro	Gly	Lys	Gly	Gly	Ile	Trp	Arg	
				80					85					90	
Val	Ile	Phe	Lys	Pro	Pro	Asp	Pro	Asp	Asn	Thr	Phe	Leu	Ser	Arg	
				95					100					105	
Leu	Asn	Glu	Phe	Leu	Ala	Gly	Glu	Gly	Met	Thr	Val	Gly	Glu	Leu	
				110					115					120	
Ser	Arg	Ala	Leu	Gly	His	Glu	Asn	Gly	Ser	Leu	Asp	Pro	Glu	Gln	
				125					130					135	
Gly	Met	Ile	Pro	Glu	Met	Trp	Ala	Pro	Met	Leu	Ala	Gln	Ala	Leu	
				140					145					150	
Glu	Ala	Leu	Gln	Pro	Ala	Leu	Gln	Cys	Leu	Lys	Tyr	Lys	Lys	Leu	
				155					160					165	
Arg	Val	Phe	Ser	Gly	Arg	Glu	Ser	Pro	Glu	Pro	Gly	Glu	Glu	Glu	
				170					175					180	
Phe	Gly	Arg	Trp	Met	Phe	His	Thr	Thr	Gln	Met	Ile	Lys	Ala	Trp	

Gln Val Pro Asp	185	Val Glu Lys Arg Arg	190	Arg Leu Leu Glu Ser	195
	200		205		210
Arg Gly Pro Ala	215	Leu Asp Val Ile Arg	220	Val Leu Lys Ile Asn	225
	230		235		240
Pro Leu Ile Thr	245	Val Asp Glu Cys Leu	250	Gln Ala Leu Glu Glu	255
	260		265		270
Phe Gly Val Thr	275	Asp Asn Pro Arg Glu	280	Leu Gln Val Lys Tyr	285
	290		295		300
Thr Thr Tyr Gln	305	Lys Asp Glu Glu Lys	310	Leu Ser Ala Tyr Val	315
	320		325		330
Arg Leu Glu Pro	335	Leu Leu Gln Lys Leu	340	Val Gln Arg Gly Ala	345
	350				
Glu Arg Asp Ala		Val Asn Gln Ala Arg		Leu Asp Gln Val Ile	
Gly Ala Val His		Lys Thr Ile Arg Arg		Glu Leu Asn Leu Pro	
Asp Gly Pro Ala		Pro Gly Phe Leu Gln		Leu Leu Val Leu Ile	
Asp Tyr Glu Ala		Ala Glu Glu Glu Glu		Ala Leu Leu Gln Ala	
Leu Glu Gly Asn		Phe Thr			

<210> 2

<211> 458

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 997947CD1

<400> 2

Met Gln Ala Thr	1	Ser Asn Leu Leu Asn	10	Leu Leu Leu Ser	15
	5		10		15
Phe Ala Gly Leu	20	Asp Pro Ser Lys Thr	25	Gln Ile Ser Pro Lys	30
	35		40		45
Gly Trp Gln Val	50	Tyr Ser Ser Ala Gln	55	Asp Pro Asp Gly Arg	60
	65		70		75
Ile Cys Thr Val	80	Val Ala Pro Glu Gln	85	Asn Leu Cys Ser Arg	90
	95		100		105
Ala Lys Ser Arg	110	Gln Leu Arg Gln Leu	115	Leu Glu Lys Val Gln	120
	125		130		135
Met Ser Gln Ser	140	Ile Glu Val Leu Asn	145	Leu Arg Thr Gln Arg	150
	155		160		165
Phe Gln Tyr Val	170	Leu Lys Met Glu Thr	175	Gln Met Lys Gly Leu	180
	185		190		195
Ala Lys Phe Arg	200	Gln Ile Glu Asp Asp	205	Arg Lys Thr Leu Met	210
	215		220		225
Lys His Phe Gln		Glu Leu Lys Glu Lys		Met Asp Glu Leu Leu	
Leu Ile Pro Val		Leu Glu Gln Tyr Lys		Thr Asp Ala Lys Leu	
Thr Gln Phe Lys		Glu Glu Ile Arg Asn		Leu Ser Ala Val Leu	
Gly Ile Gln Glu		Glu Ile Gly Ala Tyr		Asp Tyr Glu Glu Leu	
Gln Arg Val Leu		Ser Leu Glu Thr Arg		Leu Arg Asp Cys Met	
Lys Leu Thr Cys		Gly Lys Leu Met Lys		Ile Thr Gly Pro Val	
Val Lys Thr Ser		Gly Thr Arg Phe Gly		Ala Trp Met Thr Asp	

Leu	Ala	Ser	Glu	Lys	Asn	Asn	Arg	Val	Trp	Tyr	Met	Asp	Ser	Tyr	
				230					235					240	
Thr	Asn	Asn	Lys	Ile	Val	Arg	Glu	Tyr	Lys	Ser	Ile	Ala	Asp	Phe	
				245					250					255	
Val	Ser	Gly	Ala	Glu	Ser	Arg	Thr	Tyr	Asn	Leu	Pro	Phe	Lys	Trp	
				260					265					270	
Ala	Gly	Thr	Asn	His	Val	Val	Tyr	Asn	Gly	Ser	Leu	Tyr	Phe	Asn	
				275					280					285	
Lys	Tyr	Gln	Ser	Asn	Ile	Ile	Ile	Lys	Tyr	Ser	Phe	Asp	Met	Gly	
				290					295					300	
Arg	Val	Leu	Ala	Gln	Arg	Ser	Leu	Glu	Tyr	Ala	Gly	Phe	His	Asn	
				305					310					315	
Val	Tyr	Pro	Tyr	Thr	Trp	Gly	Gly	Phe	Ser	Asp	Ile	Asp	Leu	Met	
				320					325					330	
Ala	Asp	Glu	Ile	Gly	Leu	Trp	Ala	Val	Tyr	Ala	Thr	Asn	Gln	Asn	
				335					340					345	
Ala	Gly	Asn	Ile	Val	Ile	Ser	Gln	Leu	Asn	Gln	Asp	Thr	Leu	Glu	
				350					355					360	
Val	Met	Lys	Ser	Trp	Ser	Thr	Gly	Tyr	Pro	Lys	Arg	Ser	Ala	Gly	
				365					370					375	
Glu	Ser	Phe	Met	Ile	Cys	Gly	Thr	Leu	Tyr	Val	Thr	Asn	Ser	His	
				380					385					390	
Leu	Thr	Gly	Ala	Lys	Val	Tyr	Tyr	Ser	Tyr	Ser	Thr	Lys	Thr	Ser	
				395					400					405	
Thr	Tyr	Glu	Tyr	Thr	Asp	Ile	Pro	Phe	His	Asn	Gln	Tyr	Phe	His	
				410					415					420	
Ile	Ser	Met	Leu	Asp	Tyr	Asn	Ala	Arg	Asp	Arg	Ala	Leu	Tyr	Ala	
				425					430					435	
Trp	Asn	Asn	Gly	His	Gln	Val	Leu	Phe	Asn	Val	Thr	Leu	Phe	His	
				440					445					450	
Ile	Ile	Lys	Thr	Glu	Asp	Asp	Thr								
				455											

<210> 3

<211> 219

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1521513CD1

<400> 3

Met	Asn	Ser	Ser	Lys	Ser	Ser	Glu	Thr	Gln	Cys	Thr	Glu	Arg	Gly	
1				5					10					15	
Cys	Phe	Ser	Ser	Gln	Met	Phe	Leu	Trp	Thr	Val	Ala	Gly	Ile	Pro	
				20					25					30	
Ile	Leu	Phe	Leu	Ser	Ala	Cys	Phe	Ile	Thr	Arg	Cys	Val	Val	Thr	
				35					40					45	
Phe	Arg	Ile	Phe	Gln	Thr	Cys	Asp	Glu	Lys	Lys	Phe	Gln	Leu	Pro	
				50					55					60	
Glu	Asn	Phe	Thr	Glu	Leu	Ser	Cys	Tyr	Asn	Tyr	Gly	Ser	Gly	Ser	
				65					70					75	
Val	Lys	Asn	Cys	Cys	Pro	Leu	Asn	Trp	Glu	Tyr	Phe	Gln	Ser	Ser	
				80					85					90	
Cys	Tyr	Phe	Phe	Ser	Thr	Asp	Thr	Ile	Ser	Trp	Ala	Leu	Ser	Leu	
				95					100					105	
Lys	Asn	Cys	Ser	Ala	Met	Gly	Ala	His	Leu	Val	Val	Ile	Asn	Ser	
				110					115					120	
Gln	Glu	Glu	Gln	Glu	Phe	Leu	Ser	Tyr	Lys	Lys	Pro	Lys	Met	Arg	
				125					130					135	
Glu	Phe	Phe	Ile	Gly	Leu	Ser	Asp	Gln	Val	Val	Glu	Gly	Gln	Trp	
				140					145					150	
Gln	Trp	Val	Asp	Gly	Thr	Pro	Leu	Thr	Lys	Ser	Leu	Ser	Phe	Trp	

	155		160		165
Asp Val Gly Glu Pro	Asn Asn Ile Ala	Thr Leu Glu Asp Cys	Ala		
	170		175		180
Thr Met Arg Asp Ser	Ser Asn Pro Arg	Gln Asn Trp Asn Asp	Val		
	185		190		195
Thr Cys Phe Leu Asn	Tyr Phe Arg Ile	Cys Glu Met Val Gly	Ile		
	200		205		210
Asn Pro Leu Asn Lys	Gly Lys Ser Leu				
	215				

<210> 4

<211> 276

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1863994CD1

<400> 4	
Met Glu Ser Arg Met	Trp Pro Ala Leu Leu Ser His Leu Leu
1	5 10 15
Pro Leu Trp Pro Leu	Leu Leu Leu Pro Leu Pro Pro Pro Ala Gln
	20 25 30
Gly Ser Ser Ser Ser	Pro Arg Thr Pro Pro Ala Pro Ala Arg Pro
	35 40 45
Pro Cys Ala Arg Gly	Gly Pro Ser Ala Pro Arg His Val Cys Val
	50 55 60
Trp Glu Arg Ala Pro	Pro Pro Ser Arg Ser Pro Arg Val Pro Arg
	65 70 75
Ser Arg Arg Gln Val	Leu Pro Gly Thr Ala Pro Pro Ala Thr Pro
	80 85 90
Ser Gly Phe Glu Glu	Gly Pro Pro Ser Ser Gln Tyr Pro Trp Ala
	95 100 105
Ile Val Trp Gly Pro	Thr Val Ser Arg Glu Asp Gly Gly Asp Pro
	110 115 120
Asn Ser Ala Asn Pro	Gly Phe Leu Asp Tyr Gly Phe Ala Ala Pro
	125 130 135
His Gly Leu Ala Thr	Pro His Pro Asn Ser Asp Ser Met Arg Gly
	140 145 150
Asp Gly Asp Gly Leu	Ile Leu Gly Glu Ala Pro Ala Thr Leu Arg
	155 160 165
Pro Phe Leu Phe Gly	Gly Arg Gly Glu Gly Val Asp Pro Gln Leu
	170 175 180
Tyr Val Thr Ile Thr	Ile Ser Ile Ile Ile Val Leu Val Ala Thr
	185 190 195
Gly Ile Ile Phe Lys	Phe Cys Trp Asp Arg Ser Gln Lys Arg Arg
	200 205 210
Arg Pro Ser Gly Gln	Gln Gly Ala Leu Arg Gln Glu Glu Ser Gln
	215 220 225
Gln Pro Leu Thr Asp	Leu Ser Pro Ala Gly Val Thr Val Leu Gly
	230 235 240
Ala Phe Gly Asp Ser	Pro Thr Pro Thr Pro Asp His Glu Glu Pro
	245 250 255
Arg Gly Gly Pro Arg	Pro Gly Met Pro His Pro Lys Gly Ala Pro
	260 265 270
Ala Phe Gln Leu Asn	Arg
	275

<210> 5

<211> 375

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2071941CD1

<400> 5

Met	Ser	Ser	His	Lys	Gly	Ser	Val	Val	Ala	Gln	Gly	Asn	Gly	Ala	
1				5					10					15	
Pro	Ala	Ser	Asn	Arg	Glu	Ala	Asp	Thr	Val	Glu	Leu	Ala	Glu	Leu	
				20					25					30	
Gly	Pro	Leu	Leu	Glu	Glu	Lys	Gly	Lys	Arg	Val	Ile	Ala	Asn	Pro	
				35					40					45	
Pro	Lys	Ala	Glu	Glu	Glu	Gln	Thr	Cys	Pro	Val	Pro	Gln	Glu	Glu	
				50					55					60	
Glu	Glu	Glu	Val	Arg	Val	Leu	Thr	Leu	Pro	Leu	Gln	Ala	His	His	
				65					70					75	
Ala	Met	Glu	Lys	Met	Glu	Glu	Phe	Val	Tyr	Lys	Val	Trp	Glu	Gly	
				80					85					90	
Arg	Trp	Arg	Val	Ile	Pro	Tyr	Asp	Val	Leu	Pro	Asp	Trp	Leu	Lys	
				95					100					105	
Asp	Asn	Asp	Tyr	Leu	Leu	His	Gly	His	Arg	Pro	Pro	Met	Pro	Ser	
				110					115					120	
Phe	Arg	Ala	Cys	Phe	Lys	Ser	Ile	Phe	Arg	Ile	His	Thr	Glu	Thr	
				125					130					135	
Gly	Asn	Ile	Trp	Thr	His	Leu	Leu	Gly	Phe	Val	Leu	Phe	Leu	Phe	
				140					145					150	
Leu	Gly	Ile	Leu	Thr	Met	Leu	Arg	Pro	Asn	Met	Tyr	Phe	Met	Ala	
				155					160					165	
Pro	Leu	Gln	Glu	Lys	Val	Val	Phe	Gly	Met	Phe	Phe	Leu	Gly	Ala	
				170					175					180	
Val	Leu	Cys	Leu	Ser	Phe	Ser	Trp	Leu	Phe	His	Thr	Val	Tyr	Cys	
				185					190					195	
His	Ser	Glu	Lys	Val	Ser	Arg	Thr	Phe	Ser	Lys	Leu	Asp	Tyr	Ser	
				200					205					210	
Gly	Ile	Ala	Leu	Leu	Ile	Met	Gly	Ser	Phe	Val	Pro	Trp	Leu	Tyr	
				215					220					225	
Tyr	Ser	Phe	Tyr	Cys	Ser	Pro	Gln	Pro	Arg	Leu	Ile	Tyr	Leu	Ser	
				230					235					240	
Ile	Val	Cys	Val	Leu	Gly	Ile	Ser	Ala	Ile	Ile	Val	Ala	Gln	Trp	
				245					250					255	
Asp	Arg	Phe	Ala	Thr	Pro	Lys	His	Arg	Gln	Thr	Arg	Ala	Gly	Val	
				260					265					270	
Phe	Leu	Gly	Leu	Gly	Leu	Ser	Gly	Val	Val	Pro	Thr	Met	His	Phe	
				275					280					285	
Thr	Ile	Ala	Glu	Gly	Phe	Val	Lys	Ala	Thr	Thr	Val	Gly	Gln	Met	
				290					295					300	
Gly	Trp	Phe	Phe	Leu	Met	Ala	Val	Met	Tyr	Ile	Thr	Gly	Ala	Gly	
				305					310					315	
Leu	Tyr	Ala	Ala	Arg	Ile	Pro	Glu	Arg	Phe	Phe	Pro	Gly	Lys	Phe	
				320					325					330	
Asp	Ile	Trp	Phe	Gln	Ser	His	Gln	Ile	Phe	His	Val	Leu	Val	Val	
				335					340					345	
Ala	Ala	Ala	Phe	Val	His	Phe	Tyr	Gly	Val	Ser	Asn	Leu	Gln	Glu	
				350					355					360	
Phe	Arg	Tyr	Gly	Leu	Glu	Gly	Gly	Cys	Thr	Asp	Asp	Thr	Leu	Leu	
				365					370					375	

<210> 6

<211> 249

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2172512CD1

<400> 6

Met	Ser	Gly	Val	Val	Pro	Thr	Ala	Pro	Glu	Gln	Pro	Ala	Gly	Glu
1				5					10					15
Met	Glu	Asn	Gln	Thr	Lys	Pro	Pro	Asp	Pro	Arg	Pro	Asp	Ala	Pro
				20					25					30
Pro	Glu	Tyr	Ser	Ser	His	Phe	Leu	Pro	Gly	Pro	Pro	Gly	Thr	Ala
				35					40					45
Val	Pro	Pro	Pro	Thr	Gly	Tyr	Pro	Gly	Gly	Leu	Pro	Met	Gly	Tyr
				50					55					60
Tyr	Ser	Pro	Gln	Gln	Pro	Ser	Thr	Phe	Pro	Leu	Tyr	Gln	Pro	Val
				65					70					75
Gly	Gly	Ile	His	Pro	Val	Arg	Tyr	Gln	Pro	Gly	Lys	Tyr	Pro	Met
				80					85					90
Pro	Asn	Gln	Ser	Val	Pro	Ile	Thr	Trp	Met	Pro	Gly	Pro	Thr	Pro
				95					100					105
Met	Ala	Asn	Cys	Pro	Pro	Gly	Leu	Glu	Tyr	Leu	Val	Gln	Leu	Asp
				110					115					120
Asn	Ile	His	Val	Leu	Gln	His	Phe	Glu	Pro	Leu	Glu	Met	Met	Thr
				125					130					135
Cys	Phe	Glu	Thr	Asn	Asn	Arg	Tyr	Asp	Ile	Lys	Asn	Asn	Ser	Asp
				140					145					150
Gln	Met	Val	Tyr	Ile	Val	Thr	Glu	Asp	Thr	Asp	Asp	Phe	Thr	Arg
				155					160					165
Asn	Ala	Tyr	Arg	Thr	Leu	Arg	Pro	Phe	Val	Leu	Arg	Val	Thr	Asp
				170					175					180
Cys	Met	Gly	Arg	Glu	Ile	Met	Thr	Met	Gln	Arg	Pro	Phe	Arg	Cys
				185					190					195
Thr	Cys	Cys	Cys	Phe	Cys	Cys	Pro	Ser	Ala	Arg	Gln	Glu	Leu	Glu
				200					205					210
Val	Gln	Cys	Pro	Pro	Gly	Val	Thr	Ile	Gly	Phe	Val	Ala	Glu	His
				215					220					225
Trp	Asn	Leu	Cys	Arg	Ala	Val	Tyr	Ser	Ile	Gln	Lys	Lys	Lys	Lys
				230					235					240
Lys	Ile	Ala	Ala	Gln	Ala	Tyr	Ser	Leu						
				245										

<210> 7

<211> 353

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2483172CD1

<400> 7

Met	Ala	Met	Thr	Leu	Leu	Glu	Asp	Trp	Cys	Arg	Gly	Met	Asp	Val
1				5					10					15
Asn	Ser	Gln	Arg	Ala	Leu	Leu	Val	Trp	Gly	Ile	Pro	Val	Asn	Cys
				20					25					30
Asp	Glu	Ala	Glu	Ile	Glu	Glu	Thr	Leu	Gln	Ala	Ala	Met	Pro	Gln
				35					40					45
Val	Ser	Tyr	Arg	Met	Leu	Gly	Arg	Met	Phe	Trp	Arg	Glu	Glu	Asn
				50					55					60
Ala	Lys	Ala	Ala	Leu	Leu	Glu	Leu	Thr	Gly	Ala	Val	Asp	Tyr	Ala
				65					70					75
Ala	Ile	Pro	Arg	Glu	Met	Pro	Gly	Lys	Gly	Gly	Val	Trp	Lys	Val
				80					85					90
Leu	Phe	Lys	Pro	Pro	Thr	Ser	Asp	Ala	Glu	Phe	Leu	Glu	Arg	Leu
				95					100					105
His	Leu	Phe	Leu	Ala	Arg	Glu	Gly	Trp	Thr	Val	Gln	Asp	Val	Ala
				110					115					120
Arg	Val	Leu	Gly	Phe	Gln	Asn	Pro	Thr	Pro	Thr	Pro	Gly	Pro	Glu
				125					130					135

Met	Pro	Ala	Glu	Met	Leu	Asn	Tyr	Ile	Leu	Asp	Asn	Val	Ile	Gln	
				140					145					150	
Pro	Leu	Val	Glu	Ser	Ile	Trp	Tyr	Lys	Arg	Leu	Thr	Leu	Phe	Ser	
				155					160					165	
Gly	Arg	Asp	Ile	Pro	Gly	Pro	Gly	Glu	Glu	Thr	Phe	Asp	Pro	Trp	
				170					175					180	
Leu	Glu	His	Thr	Asn	Glu	Val	Leu	Glu	Glu	Trp	Gln	Val	Ser	Asp	
				185					190					195	
Val	Glu	Lys	Arg	Arg	Arg	Leu	Met	Glu	Ser	Leu	Arg	Gly	Pro	Ala	
				200					205					210	
Ala	Asp	Val	Ile	Arg	Ile	Leu	Lys	Ser	Asn	Asn	Pro	Ala	Ile	Thr	
				215					220					225	
Thr	Ala	Glu	Cys	Leu	Lys	Ala	Leu	Glu	Gln	Val	Phe	Gly	Ser	Val	
				230					235					240	
Glu	Ser	Ser	Arg	Asp	Ala	Gln	Ile	Lys	Phe	Leu	Asn	Thr	Tyr	Gln	
				245					250					255	
Asn	Pro	Gly	Glu	Lys	Leu	Ser	Ala	Tyr	Val	Ile	Arg	Leu	Glu	Pro	
				260					265					270	
Leu	Leu	Gln	Lys	Val	Val	Glu	Lys	Gly	Ala	Ile	Asp	Lys	Asp	Asn	
				275					280					285	
Val	Asn	Gln	Ala	Arg	Leu	Glu	Gln	Val	Ile	Ala	Gly	Ala	Asn	His	
				290					295					300	
Ser	Gly	Ala	Ile	Arg	Arg	Gln	Leu	Trp	Leu	Thr	Gly	Ala	Gly	Glu	
				305					310					315	
Gly	Pro	Ala	Pro	Asn	Leu	Phe	Gln	Leu	Leu	Val	Gln	Ile	Arg	Glu	
				320					325					330	
Glu	Glu	Ala	Lys	Glu	Glu	Glu	Glu	Glu	Ala	Glu	Ala	Thr	Leu	Leu	
				335					340					345	
Gln	Leu	Gly	Leu	Glu	Gly	His	Phe								
				350											

<210> 8

<211> 194

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2656128CD1

<400> 8

Met	His	Asp	Ser	Asn	Asn	Val	Glu	Lys	Asp	Ile	Thr	Pro	Ser	Glu	
1				5					10					15	
Leu	Pro	Ala	Asn	Pro	Gly	Cys	Leu	His	Ser	Lys	Glu	His	Ser	Ile	
				20					25					30	
Lys	Ala	Thr	Leu	Ile	Trp	Arg	Leu	Phe	Phe	Leu	Ile	Met	Phe	Leu	
				35					40					45	
Thr	Ile	Ile	Val	Cys	Gly	Met	Val	Ala	Ala	Leu	Ser	Ala	Ile	Arg	
				50					55					60	
Ala	Asn	Cys	His	Gln	Glu	Pro	Ser	Val	Cys	Leu	Gln	Ala	Ala	Cys	
				65					70					75	
Pro	Glu	Ser	Trp	Ile	Gly	Phe	Gln	Arg	Lys	Cys	Phe	Tyr	Phe	Ser	
				80					85					90	
Asp	Asp	Thr	Lys	Asn	Trp	Thr	Ser	Ser	Gln	Arg	Phe	Cys	Asp	Ser	
				95					100					105	
Gln	Asp	Ala	Asp	Leu	Ala	Gln	Val	Glu	Ser	Phe	Gln	Glu	Leu	Asn	
				110					115					120	
Phe	Leu	Leu	Arg	Tyr	Lys	Gly	Pro	Ser	Asp	His	Trp	Ile	Gly	Leu	
				125					130					135	
Ser	Arg	Glu	Gln	Gly	Gln	Pro	Trp	Lys	Trp	Ile	Asn	Gly	Thr	Glu	
				140					145					150	
Trp	Thr	Arg	Gln	Leu	Val	Met	Lys	Glu	Asp	Gly	Ala	Asn	Leu	Tyr	
				155					160					165	
Val	Ala	Lys	Val	Ser	Gln	Val	Pro	Arg	Met	Asn	Pro	Arg	Pro	Val	

	170		175		180
Met Val Ser Tyr	Pro Gly Ser Arg Arg	Val Cys Leu Phe Glu			
	185	190			

<210> 9

<211> 322

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5855841CD1

<400> 9

Met	Ser	Ser	Leu	Gly	Gly	Gly	Ser	Gln	Asp	Ala	Gly	Gly	Ser	Ser
1				5					10					15
Ser	Ser	Ser	Thr	Asn	Gly	Ser	Gly	Gly	Ser	Gly	Ser	Ser	Gly	Pro
				20					25					30
Lys	Ala	Gly	Ala	Ala	Asp	Lys	Ser	Ala	Val	Val	Ala	Ala	Ala	Ala
				35					40					45
Pro	Ala	Ser	Val	Ala	Asp	Asp	Thr	Pro	Pro	Pro	Glu	Arg	Arg	Asn
				50					55					60
Lys	Ser	Gly	Ile	Ile	Ser	Glu	Pro	Leu	Asn	Lys	Ser	Leu	Arg	Arg
				65					70					75
Ser	Arg	Pro	Leu	Ser	His	Tyr	Ser	Ser	Phe	Gly	Ser	Ser	Gly	Gly
				80					85					90
Ser	Gly	Gly	Gly	Ser	Met	Met	Gly	Gly	Glu	Ser	Ala	Asp	Lys	Ala
				95					100					105
Thr	Ala	Ala	Ala	Ala	Ala	Ala	Ser	Leu	Leu	Ala	Asn	Gly	His	Asp
				110					115					120
Leu	Ala	Ala	Ala	Met	Ala	Val	Asp	Lys	Ser	Asn	Pro	Thr	Ser	Lys
				125					130					135
His	Lys	Ser	Gly	Ala	Val	Ala	Ser	Leu	Leu	Ser	Lys	Ala	Glu	Arg
				140					145					150
Ala	Thr	Glu	Leu	Ala	Ala	Glu	Gly	Gln	Leu	Thr	Leu	Gln	Gln	Phe
				155					160					165
Ala	Gln	Ser	Thr	Glu	Met	Leu	Lys	Arg	Val	Val	Gln	Glu	His	Leu
				170					175					180
Pro	Leu	Met	Ser	Glu	Ala	Gly	Ala	Gly	Leu	Pro	Asp	Met	Glu	Ala
				185					190					195
Val	Ala	Gly	Ala	Glu	Ala	Leu	Asn	Gly	Gln	Ser	Asp	Phe	Pro	Tyr
				200					205					210
Leu	Gly	Ala	Phe	Pro	Ile	Asn	Pro	Gly	Leu	Phe	Ile	Met	Thr	Pro
				215					220					225
Ala	Gly	Val	Phe	Leu	Ala	Glu	Ser	Ala	Leu	His	Met	Ala	Gly	Leu
				230					235					240
Ala	Glu	Tyr	Pro	Met	Gln	Gly	Glu	Leu	Ala	Ser	Ala	Ile	Ser	Ser
				245					250					255
Gly	Lys	Lys	Lys	Arg	Lys	Arg	Cys	Gly	Met	Cys	Ala	Pro	Cys	Arg
				260					265					270
Arg	Arg	Ile	Asn	Cys	Glu	Gln	Cys	Ser	Ser	Cys	Arg	Asn	Arg	Lys
				275					280					285
Thr	Gly	His	Gln	Ile	Cys	Lys	Phe	Arg	Lys	Cys	Glu	Glu	Leu	Lys
				290					295					300
Lys	Lys	Pro	Ser	Ala	Ala	Leu	Glu	Lys	Val	Met	Leu	Pro	Thr	Gly
				305					310					315
Ala	Ala	Phe	Arg	Trp	Phe	Gln								
				320										

<210> 10

<211> 335

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 603462CD1

<400> 10

```

Met Leu Gln Gly His Ser Ser Val Phe Gln Ala Leu Leu Gly Thr
 1          5          10          15
Phe Phe Thr Trp Gly Met Thr Ala Ala Gly Ala Ala Leu Val Phe
          20          25          30
Val Phe Ser Ser Gly Gln Arg Arg Ile Leu Asp Gly Ser Leu Gly
          35          40          45
Phe Ala Ala Gly Val Met Leu Ala Ala Ser Tyr Trp Ser Leu Leu
          50          55          60
Ala Pro Ala Val Glu Met Ala Thr Ser Ser Gly Gly Phe Gly Ala
          65          70          75
Phe Ala Phe Phe Pro Val Ala Val Gly Phe Thr Leu Gly Ala Ala
          80          85          90
Phe Val Tyr Leu Ala Asp Leu Leu Met Pro His Leu Gly Ala Ala
          95          100          105
Glu Asp Pro Gln Thr Ala Leu Ala Leu Asn Phe Gly Ser Thr Leu
          110          115          120
Met Lys Lys Lys Ser Asp Pro Glu Gly Pro Ala Leu Leu Phe Pro
          125          130          135
Glu Ser Glu Leu Ser Ile Arg Ile Asp Lys Ser Glu Asn Gly Glu
          140          145          150
Ala Tyr Gln Arg Lys Lys Ala Ala Ala Thr Gly Leu Pro Glu Gly
          155          160          165
Pro Ala Val Pro Val Pro Ser Arg Gly Asn Leu Ala Gln Pro Gly
          170          175          180
Gly Ser Ser Trp Arg Arg Ile Ala Leu Leu Ile Leu Ala Ile Thr
          185          190          195
Ile His Asn Val Pro Glu Gly Leu Ala Val Gly Val Gly Phe Gly
          200          205          210
Ala Ile Glu Lys Thr Ala Ser Ala Thr Phe Glu Ser Ala Arg Asn
          215          220          225
Leu Ala Ile Gly Ile Gly Ile Gln Asn Phe Pro Glu Gly Leu Ala
          230          235          240
Val Ser Leu Pro Leu Arg Gly Ala Gly Phe Ser Thr Trp Arg Ala
          245          250          255
Phe Trp Tyr Gly Gln Leu Ser Gly Met Val Glu Pro Leu Ala Gly
          260          265          270
Val Phe Gly Ala Phe Ala Val Val Leu Ala Glu Pro Ile Leu Pro
          275          280          285
Tyr Ala Leu Ala Phe Ala Ala Gly Ala Met Val Tyr Val Val Met
          290          295          300
Asp Asp Ile Ile Pro Glu Ala Gln Ile Ser Gly Asn Gly Lys Leu
          305          310          315
Ala Ser Trp Ala Ser Ile Leu Gly Phe Val Val Met Met Ser Leu
          320          325          330
Asp Val Gly Leu Gly
          335

```

<210> 11

<211> 620

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 747681CD1

<400> 11

```

Met Gln Val Ser Lys Arg Met Leu Ala Gly Gly Val Arg Ser Met
 1          5          10          15
Pro Ser Pro Leu Leu Ala Cys Trp Gln Pro Ile Leu Leu Leu Val

```

	20		25		30
Leu Gly Ser Val	Leu Ser Gly Ser Ala Thr	Gly Cys Pro Pro	Arg		
	35		40		45
Cys Glu Cys Ser	Ala Gln Asp Arg Ala Val	Leu Cys His Arg	Lys		
	50		55		60
Arg Phe Val Ala	Val Pro Glu Gly Ile Pro	Thr Glu Thr Arg	Leu		
	65		70		75
Leu Asp Leu Gly	Lys Asn Arg Ile Lys Thr	Leu Asn Gln Asp	Glu		
	80		85		90
Phe Ala Ser Phe	Pro His Leu Glu Glu Leu	Glu Leu Asn Glu	Asn		
	95		100		105
Ile Val Ser Ala	Val Glu Pro Gly Ala Phe	Asn Asn Leu Phe	Asn		
	110		115		120
Leu Arg Thr Leu	Gly Leu Arg Ser Asn Arg	Leu Lys Leu Ile	Pro		
	125		130		135
Leu Gly Val Phe	Thr Gly Leu Ser Asn Leu	Thr Lys Leu Asp	Ile		
	140		145		150
Ser Glu Asn Lys	Ile Val Ile Leu Leu Asp	Tyr Met Phe Gln	Asp		
	155		160		165
Leu Tyr Asn Leu	Lys Ser Leu Glu Val Gly	Asp Asn Asp Leu	Val		
	170		175		180
Tyr Ile Ser His	Arg Ala Phe Ser Gly Leu	Asn Ser Leu Glu	Gln		
	185		190		195
Leu Thr Leu Glu	Lys Cys Asn Leu Thr Ser	Ile Pro Thr Glu	Ala		
	200		205		210
Leu Ser His Leu	His Gly Leu Ile Val Leu	Arg Leu Arg His	Leu		
	215		220		225
Asn Ile Asn Ala	Ile Arg Asp Tyr Ser Phe	Lys Arg Leu Tyr	Arg		
	230		235		240
Leu Lys Val Leu	Glu Ile Ser His Trp Pro	Tyr Leu Asp Thr	Met		
	245		250		255
Thr Pro Asn Cys	Leu Tyr Gly Leu Asn Leu	Thr Ser Leu Ser	Ile		
	260		265		270
Thr His Cys Asn	Leu Thr Ala Val Pro Tyr	Leu Ala Val Arg	His		
	275		280		285
Leu Val Tyr Leu	Arg Phe Leu Asn Leu Ser	Tyr Asn Pro Ile	Ser		
	290		295		300
Thr Ile Glu Gly	Ser Met Leu His Glu Leu	Leu Arg Leu Gln	Glu		
	305		310		315
Ile Gln Leu Val	Gly Gly Gln Leu Ala Val	Val Glu Pro Tyr	Ala		
	320		325		330
Phe Arg Gly Leu	Asn Tyr Leu Arg Val Leu	Asn Val Ser Gly	Asn		
	335		340		345
Gln Leu Thr Thr	Leu Glu Glu Ser Val Phe	His Ser Val Gly	Asn		
	350		355		360
Leu Glu Thr Leu	Ile Leu Asp Ser Asn Pro	Leu Ala Cys Asp	Cys		
	365		370		375
Arg Leu Leu Trp	Val Phe Arg Arg Arg Trp	Arg Leu Asn Phe	Asn		
	380		385		390
Arg Gln Gln Pro	Thr Cys Ala Thr Pro Glu	Phe Val Gln Gly	Lys		
	395		400		405
Glu Phe Lys Asp	Phe Pro Asp Val Leu Leu	Pro Asn Tyr Phe	Thr		
	410		415		420
Cys Arg Arg Ala	Arg Ile Arg Asp Arg Lys	Ala Gln Gln Val	Phe		
	425		430		435
Val Asp Glu Gly	His Thr Val Gln Phe Val	Cys Arg Ala Asp	Gly		
	440		445		450
Asp Pro Pro Pro	Ala Ile Leu Trp Leu Ser	Pro Arg Lys His	Leu		
	455		460		465
Val Ser Ala Lys	Ser Asn Gly Arg Leu Thr	Val Phe Pro Asp	Gly		
	470		475		480
Thr Leu Glu Val	Arg Tyr Ala Gln Val Gln	Asp Asn Gly Thr	Tyr		
	485		490		495

Leu Cys Ile Ala	Ala Asn Ala Gly Gly	Asn Asp Ser Met Pro	Ala
500	505	510	
His Leu His Val	Arg Ser Tyr Ser Pro	Asp Trp Pro His Gln	Pro
515	520	525	
Asn Lys Thr Phe	Ala Phe Ile Ser Asn	Gln Pro Gly Glu Gly	Glu
530	535	540	
Ala Asn Ser Thr	Arg Ala Thr Val Pro	Phe Pro Phe Asp Ile	Lys
545	550	555	
Thr Leu Ile Ile	Ala Thr Thr Met Gly	Phe Ile Ser Phe Leu	Gly
560	565	570	
Val Val Leu Phe	Cys Leu Val Leu Leu	Phe Leu Trp Ser Arg	Gly
575	580	585	
Lys Gly Asn Thr	Lys His Asn Ile Glu	Ile Glu Tyr Val Pro	Arg
590	595	600	
Lys Ser Asp Ala	Gly Ile Ser Ser Ala	Asp Ala Pro Arg Lys	Phe
605	610	615	
Asn Met Lys Met	Ile		
620			

<210> 12

<211> 491

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 919469CD1

<400> 12

Met Ala Gly Gln Gly	Leu Pro Leu His Val	Ala Thr Leu Leu Thr
1	5	10
Gly Leu Leu Glu Cys	Leu Gly Phe Ala Gly	Val Leu Phe Gly Trp
20	25	30
Pro Ser Leu Val Phe	Val Phe Lys Asn Glu	Asp Tyr Phe Lys Asp
35	40	45
Leu Cys Gly Pro Asp	Ala Gly Pro Ile Gly	Asn Ala Thr Gly Gln
50	55	60
Ala Asp Cys Lys Ala	Gln Asp Glu Arg Phe	Ser Leu Ile Phe Thr
65	70	75
Leu Gly Ser Phe Met	Asn Asn Phe Met Thr	Phe Pro Thr Gly Tyr
80	85	90
Ile Phe Asp Arg Phe	Lys Thr Thr Val Ala	Arg Leu Ile Ala Ile
95	100	105
Phe Phe Tyr Thr Thr	Ala Thr Leu Ile Ile	Ala Phe Thr Ser Ala
110	115	120
Gly Ser Ala Val Leu	Leu Phe Leu Ala Met	Pro Met Leu Thr Ile
125	130	135
Gly Gly Ile Leu Phe	Leu Ile Thr Asn Leu	Gln Ile Gly Asn Leu
140	145	150
Phe Gly Gln His Arg	Ser Thr Ile Ile Thr	Leu Tyr Asn Gly Ala
155	160	165
Phe Asp Ser Ser Ser	Ala Val Phe Leu Ile	Ile Lys Leu Leu Tyr
170	175	180
Glu Lys Gly Ile Ser	Leu Arg Ala Ser Phe	Ile Phe Ile Ser Val
185	190	195
Cys Ser Thr Trp His	Val Ala Arg Thr Phe	Leu Leu Met Pro Arg
200	205	210
Gly His Ile Pro Tyr	Pro Leu Pro Pro Asn	Tyr Ser Tyr Gly Leu
215	220	225
Cys Pro Gly Asn Gly	Thr Thr Lys Glu Glu	Lys Glu Thr Ala Glu
230	235	240
His Glu Asn Arg Glu	Leu Gln Ser Lys Glu	Phe Leu Ser Ala Lys
245	250	255
Glu Glu Thr Pro Gly	Ala Gly Gln Lys Gln	Glu Leu Arg Ser Phe

260	265	270
Trp Ser Tyr Ala Phe Ser Arg Arg Phe	Ala Trp His Leu Val Trp	
275	280	285
Leu Ser Val Ile Gln Leu Trp His Tyr	Leu Phe Ile Gly Thr Leu	
290	295	300
Asn Ser Leu Leu Thr Asn Met Ala Gly	Gly Asp Met Ala Arg Val	
305	310	315
Ser Thr Tyr Thr Asn Ala Phe Ala Phe	Thr Gln Phe Gly Val Leu	
320	325	330
Cys Ala Pro Trp Asn Gly Leu Leu Met	Asp Arg Leu Lys Gln Lys	
335	340	345
Tyr Gln Lys Glu Ala Arg Lys Thr Gly	Ser Ser Thr Leu Ala Val	
350	355	360
Ala Leu Cys Ser Thr Val Pro Ser Leu	Ala Leu Thr Ser Leu Leu	
365	370	375
Cys Leu Gly Phe Ala Leu Cys Ala Ser	Val Pro Ile Leu Pro Leu	
380	385	390
Gln Tyr Leu Thr Phe Ile Leu Gln Val	Ile Ser Arg Ser Phe Leu	
395	400	405
Tyr Gly Ser Asn Ala Ala Phe Leu Thr	Leu Ala Phe Pro Ser Glu	
410	415	420
His Phe Gly Lys Leu Phe Gly Leu Val	Met Ala Leu Ser Ala Val	
425	430	435
Val Ser Leu Leu Gln Phe Pro Ile Phe	Thr Leu Ile Lys Gly Ser	
440	445	450
Leu Gln Asn Asp Pro Phe Tyr Val Asn	Val Met Phe Met Leu Ala	
455	460	465
Ile Leu Leu Thr Phe Phe His Pro Phe	Leu Val Tyr Arg Glu Cys	
470	475	480
Arg Thr Trp Lys Glu Ser Pro Ser Ala	Ile Ala	
485	490	

<210> 13

<211> 580

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 977658CD1

<400> 13

Met Thr Ala Pro Ala Gly Pro Arg Gly Ser Glu Thr Glu Arg Leu	
1 5 10 15	
Leu Thr Pro Asn Pro Gly Tyr Gly Thr Gln Ala Gly Pro Ser Pro	
20 25 30	
Ala Pro Pro Thr Pro Pro Glu Glu Glu Asp Leu Arg Arg Arg Leu	
35 40 45	
Lys Tyr Phe Phe Met Ser Pro Cys Asp Lys Phe Arg Ala Lys Gly	
50 55 60	
Arg Lys Pro Cys Lys Leu Met Leu Gln Val Val Lys Ile Leu Val	
65 70 75	
Val Thr Val Gln Leu Ile Leu Phe Gly Leu Ser Asn Gln Leu Ala	
80 85 90	
Val Thr Phe Arg Glu Glu Asn Thr Ile Ala Phe Arg His Leu Phe	
95 100 105	
Leu Leu Gly Tyr Ser Asp Gly Ala Asp Thr Phe Ala Ala Tyr	
110 115 120	
Thr Arg Glu Gln Leu Tyr Gln Ala Ile Phe His Ala Val Asp Gln	
125 130 135	
Tyr Leu Ala Leu Pro Asp Val Ser Leu Gly Arg Tyr Ala Tyr Val	
140 145 150	
Arg Gly Gly Gly Asp Pro Trp Thr Asn Gly Ser Gly Leu Ala Leu	
155 160 165	

Cys	Gln	Arg	Tyr	Tyr	His	Arg	Gly	His	Val	Asp	Pro	Ala	Asn	Asp	
				170					175					180	
Thr	Phe	Asp	Ile	Asp	Pro	Met	Val	Val	Thr	Asp	Cys	Ile	Gln	Val	
				185					190					195	
Asp	Pro	Pro	Glu	Arg	Pro	Pro	Pro	Pro	Pro	Ser	Asp	Asp	Leu	Thr	
				200					205					210	
Leu	Leu	Glu	Ser	Ser	Ser	Ser	Tyr	Lys	Asn	Leu	Thr	Leu	Lys	Phe	
				215					220					225	
His	Lys	Leu	Val	Asn	Val	Thr	Ile	His	Phe	Arg	Leu	Lys	Thr	Ile	
				230					235					240	
Asn	Leu	Gln	Ser	Leu	Ile	Asn	Asn	Glu	Ile	Pro	Asp	Cys	Tyr	Thr	
				245					250					255	
Phe	Ser	Val	Leu	Ile	Thr	Phe	Asp	Asn	Lys	Ala	His	Ser	Gly	Arg	
				260					265					270	
Ile	Pro	Ile	Ser	Leu	Glu	Thr	Gln	Ala	His	Ile	Gln	Glu	Cys	Lys	
				275					280					285	
His	Pro	Ser	Val	Phe	Gln	His	Gly	Asp	Asn	Ser	Phe	Arg	Leu	Leu	
				290					295					300	
Phe	Asp	Val	Val	Ile	Leu	Thr	Cys	Ser	Leu	Ser	Phe	Leu	Leu		
				305					310					315	
Cys	Ala	Arg	Ser	Leu	Leu	Arg	Gly	Phe	Leu	Leu	Gln	Asn	Glu	Phe	
				320					325					330	
Val	Gly	Phe	Met	Trp	Arg	Gln	Arg	Gly	Arg	Val	Ile	Ser	Leu	Trp	
				335					340					345	
Glu	Arg	Leu	Glu	Phe	Val	Asn	Gly	Trp	Tyr	Ile	Leu	Leu	Val	Thr	
				350					355					360	
Ser	Asp	Val	Leu	Thr	Ile	Ser	Gly	Thr	Ile	Met	Lys	Ile	Gly	Ile	
				365					370					375	
Glu	Ala	Lys	Asn	Leu	Ala	Ser	Tyr	Asp	Val	Cys	Ser	Ile	Leu	Leu	
				380					385					390	
Gly	Thr	Ser	Thr	Leu	Leu	Val	Trp	Val	Gly	Val	Ile	Arg	Tyr	Leu	
				395					400					405	
Thr	Phe	Phe	His	Asn	Tyr	Asn	Ile	Leu	Ile	Ala	Thr	Leu	Arg	Val	
				410					415					420	
Ala	Leu	Pro	Ser	Val	Met	Arg	Phe	Cys	Cys	Cys	Val	Ala	Val	Ile	
				425					430					435	
Tyr	Leu	Gly	Tyr	Cys	Phe	Cys	Gly	Trp	Ile	Val	Leu	Gly	Pro	Tyr	
				440					445					450	
His	Val	Lys	Phe	Arg	Ser	Leu	Ser	Met	Val	Ser	Glu	Cys	Leu	Phe	
				455					460					465	
Ser	Leu	Ile	Asn	Gly	Asp	Asp	Met	Phe	Val	Thr	Phe	Ala	Ala	Met	
				470					475					480	
Gln	Ala	Gln	Gln	Gly	Arg	Ser	Ser	Leu	Val	Trp	Leu	Phe	Ser	Gln	
				485					490					495	
Leu	Tyr	Leu	Tyr	Ser	Phe	Ile	Ser	Leu	Phe	Ile	Tyr	Met	Val	Leu	
				500					505					510	
Ser	Leu	Phe	Ile	Ala	Leu	Ile	Thr	Gly	Ala	Tyr	Asp	Thr	Ile	Lys	
				515					520					525	
His	Pro	Gly	Gly	Ala	Gly	Ala	Glu	Glu	Ser	Glu	Leu	Gln	Ala	Tyr	
				530					535					540	
Ile	Ala	Gln	Cys	Gln	Asp	Ser	Pro	Thr	Ser	Gly	Lys	Phe	Arg	Arg	
				545					550					555	
Gly	Ser	Gly	Ser	Ala	Cys	Ser	Leu	Leu	Cys	Cys	Cys	Gly	Arg	Asp	
				560					565					570	
Pro	Ser	Glu	Glu	His	Ser	Leu	Leu	Val	Asn						
				575					580						

<210> 14

<211> 455

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1004703CD1

<400> 14

Met	Ser	Phe	Leu	Ile	Asp	Ser	Ser	Ile	Met	Ile	Thr	Ser	Gln	Ile
1				5					10					15
Leu	Phe	Phe	Gly	Phe	Gly	Trp	Leu	Phe	Phe	Met	Arg	Gln	Leu	Phe
			20						25					30
Lys	Asp	Tyr	Glu	Ile	Arg	Gln	Tyr	Val	Val	Gln	Val	Ile	Phe	Ser
			35						40					45
Val	Thr	Phe	Ala	Phe	Ser	Cys	Thr	Met	Phe	Glu	Leu	Ile	Ile	Phe
			50						55					60
Glu	Ile	Leu	Gly	Val	Leu	Asn	Ser	Ser	Ser	Arg	Tyr	Phe	His	Trp
			65						70					75
Lys	Met	Asn	Leu	Cys	Val	Ile	Leu	Leu	Ile	Leu	Val	Phe	Met	Val
			80						85					90
Pro	Phe	Tyr	Ile	Gly	Tyr	Phe	Ile	Val	Ser	Asn	Ile	Arg	Leu	Leu
			95						100					105
His	Lys	Gln	Arg	Leu	Leu	Phe	Ser	Cys	Leu	Leu	Trp	Leu	Thr	Phe
			110						115					120
Met	Tyr	Phe	Phe	Trp	Lys	Leu	Gly	Asp	Leu	Phe	Pro	Ile	Leu	Ser
			125						130					135
Pro	Lys	His	Gly	Ile	Leu	Ser	Ile	Glu	Gln	Leu	Ile	Ser	Arg	Val
			140						145					150
Gly	Val	Ile	Gly	Val	Thr	Leu	Met	Ala	Leu	Leu	Ser	Gly	Phe	Gly
			155						160					165
Ala	Val	Asn	Cys	Pro	Tyr	Thr	Tyr	Met	Ser	Tyr	Phe	Leu	Arg	Asn
			170						175					180
Val	Thr	Asp	Thr	Asp	Ile	Leu	Ala	Leu	Glu	Arg	Arg	Leu	Leu	Gln
			185						190					195
Thr	Met	Asp	Met	Ile	Ile	Ser	Lys	Lys	Lys	Arg	Met	Ala	Met	Ala
			200						205					210
Arg	Arg	Thr	Met	Phe	Gln	Lys	Gly	Glu	Val	His	Asn	Lys	Pro	Ser
			215						220					225
Gly	Phe	Trp	Gly	Met	Ile	Lys	Ser	Val	Thr	Thr	Ser	Ala	Ser	Gly
			230						235					240
Ser	Glu	Asn	Leu	Thr	Leu	Ile	Gln	Gln	Glu	Val	Asp	Ala	Leu	Glu
			245						250					255
Glu	Leu	Ser	Arg	Gln	Leu	Phe	Leu	Glu	Thr	Ala	Asp	Leu	Tyr	Ala
			260						265					270
Thr	Lys	Glu	Arg	Ile	Glu	Tyr	Ser	Lys	Thr	Phe	Lys	Gly	Lys	Tyr
			275						280					285
Phe	Asn	Phe	Leu	Gly	Tyr	Phe	Phe	Ser	Ile	Tyr	Cys	Val	Trp	Lys
			290						295					300
Ile	Phe	Met	Ala	Thr	Ile	Asn	Ile	Val	Phe	Asp	Arg	Val	Gly	Lys
			305						310					315
Thr	Asp	Pro	Val	Thr	Arg	Gly	Ile	Glu	Ile	Thr	Val	Asn	Tyr	Leu
			320						325					330
Gly	Ile	Gln	Phe	Asp	Val	Lys	Phe	Trp	Ser	Gln	His	Ile	Ser	Phe
			335						340					345
Ile	Leu	Val	Gly	Ile	Ile	Ile	Val	Thr	Ser	Ile	Arg	Gly	Leu	Leu
			350						355					360
Ile	Thr	Leu	Thr	Lys	Phe	Phe	Tyr	Ala	Ile	Ser	Ser	Ser	Lys	Ser
			365						370					375
Ser	Asn	Val	Ile	Val	Leu	Leu	Leu	Ala	Gln	Ile	Met	Gly	Met	Tyr
			380						385					390
Phe	Val	Ser	Ser	Val	Leu	Leu	Ile	Arg	Met	Ser	Met	Pro	Leu	Glu
			395						400					405
Tyr	Arg	Thr	Ile	Ile	Thr	Glu	Val	Leu	Gly	Glu	Leu	Gln	Phe	Asn
			410						415					420
Phe	Tyr	His	Arg	Trp	Phe	Asp	Val	Ile	Phe	Leu	Val	Ser	Ala	Leu
			425						430					435
Ser	Ser	Ile	Leu	Phe	Leu	Tyr	Leu	Ala	His	Lys	Gln	Ala	Pro	Glu
			440						445					450

<223> Incyte ID No: 1334051CD1

<213> Homo sapiens

<223> Incyte ID No: 1336728CD1

Leu Val Val Thr Ala Thr Ala Ser Pro Pro Ala Gly Leu Leu Ser

	20		25		30
Leu Leu Thr Ser Gly	Gln Gly Ala Leu Asp	Gln Glu Ala Leu Gly			
35	40	45			
Gly Leu Leu Asn Thr	Leu Ala Asp Arg Val	His Cys Thr Asn Gly			
50	55	60			
Pro Cys Gly Lys Cys	Leu Ser Val Glu Asp	Ala Leu Gly Leu Gly			
65	70	75			
Glu Pro Glu Gly Ser	Gly Leu Pro Pro Gly	Pro Val Leu Glu Ala			
80	85	90			
Arg Tyr Val Ala Arg	Leu Ser Ala Ala Ala	Val Leu Tyr Leu Ser			
95	100	105			
Asn Pro Glu Gly Thr	Cys Glu Asp Thr Arg	Ala Gly Leu Trp Ala			
110	115	120			
Ser His Ala Asp His	Leu Leu Ala Leu Leu	Glu Ser Pro Lys Ala			
125	130	135			
Leu Thr Pro Gly Leu	Ser Trp Leu Leu Gln	Arg Met Gln Ala Arg			
140	145	150			
Ala Ala Gly Gln Thr	Pro Lys Thr Ala Cys	Val Asp Ile Pro Gln			
155	160	165			
Leu Leu Glu Glu Ala	Val Gly Ala Gly Ala	Pro Gly Ser Ala Gly			
170	175	180			
Gly Val Leu Ala Ala	Leu Leu Asp His Val	Arg Ser Gly Ser Cys			
185	190	195			
Phe His Ala Leu Pro	Ser Pro Gln Tyr Phe	Val Asp Phe Val Phe			
200	205	210			
Gln Gln His Ser Ser	Glu Val Pro Met Thr	Leu Ala Glu Leu Ser			
215	220	225			
Ala Leu Met Gln Arg	Leu Gly Val Gly Arg	Glu Ala His Ser Asp			
230	235	240			
His Ser His Arg His	Arg Gly Ala Ser Ser	Arg Asp Pro Val Pro			
245	250	255			
Leu Ile Ser Ser Ser	Asn Ser Ser Ser Ser	Val Trp Asp Thr Val Cys			
260	265	270			
Leu Ser Ala Arg Asp	Val Met Ala Ala Tyr	Gly Leu Ser Glu Gln			
275	280	285			
Ala Gly Val Thr Pro	Glu Ala Trp Ala Gln	Leu Ser Pro Ala Leu			
290	295	300			
Leu Gln Gln Gln Leu	Ser Gly Ala Cys Thr	Ser Gln Ser Arg Pro			
305	310	315			
Pro Val Gln Asp Gln	Leu Ser Gln Ser Glu	Arg Tyr Leu Tyr Gly			
320	325	330			
Ser Leu Ala Thr Leu	Leu Ile Cys Leu Cys	Ala Val Phe Gly Leu			
335	340	345			
Leu Leu Leu Thr Cys	Thr Gly Cys Arg Gly	Val Thr His Tyr Ile			
350	355	360			
Leu Gln Thr Phe Leu	Ser Leu Ala Val Gly	Ala Leu Thr Gly Asp			
365	370	375			
Ala Val Leu His Leu	Thr Pro Lys Val Leu	Gly Leu His Thr His			
380	385	390			
Ser Glu Glu Gly Leu	Ser Pro Gln Pro Thr	Trp Arg Leu Leu Ala			
395	400	405			
Met Leu Ala Gly Leu	Tyr Ala Phe Phe Leu	Phe Glu Asn Leu Phe			
410	415	420			
Asn Leu Leu Leu Pro	Arg Asp Pro Glu Asp	Leu Glu Asp Gly Pro			
425	430	435			
Cys Gly His Ser Ser	His Ser His Gly Gly	His Ser His Gly Val			
440	445	450			
Ser Leu Gln Leu Ala	Pro Ser Glu Leu Arg	Gln Pro Lys Pro Pro			
455	460	465			
His Glu Gly Ser Arg	Ala Asp Leu Val Ala	Glu Glu Ser Pro Glu			
470	475	480			
Leu Leu Asn Pro Glu	Pro Arg Arg Leu Ser	Pro Glu Leu Arg Leu			
485	490	495			

Leu	Pro	Tyr	Met	Ile	Thr	Leu	Gly	Asp	Ala	Val	His	Asn	Phe	Ala
				500					505					510
Asp	Gly	Leu	Ala	Val	Gly	Ala	Ala	Phe	Ala	Ser	Ser	Trp	Lys	Thr
				515					520					525
Gly	Leu	Ala	Thr	Ser	Leu	Ala	Val	Phe	Cys	His	Glu	Leu	Pro	His
				530					535					540
Glu	Leu	Gly	Asp	Phe	Ala	Ala	Leu	Leu	His	Ala	Gly	Leu	Ser	Val
				545					550					555
Arg	Gln	Ala	Leu	Leu	Leu	Asn	Leu	Ala	Ser	Ala	Leu	Thr	Ala	Phe
				560					565					570
Ala	Gly	Leu	Tyr	Val	Ala	Leu	Ala	Val	Gly	Val	Ser	Glu	Glu	Ser
				575					580					585
Glu	Ala	Trp	Ile	Leu	Ala	Val	Ala	Thr	Gly	Leu	Phe	Leu	Tyr	Val
				590					595					600
Ala	Leu	Cys	Asp	Met	Leu	Pro	Ala	Met	Leu	Lys	Val	Arg	Asp	Pro
				605					610					615
Arg	Pro	Trp	Leu	Leu	Phe	Leu	Leu	His	Asn	Val	Gly	Leu	Leu	Gly
				620					625					630
Gly	Trp	Thr	Val	Leu	Leu	Leu	Leu	Ser	Leu	Tyr	Glu	Asp	Asp	Ile
				635					640					645

Thr Phe

<210> 17

<211> 406

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1452856CD1

<400> 17

Met	Ala	Glu	Asn	Gly	Lys	Asn	Cys	Asp	Gln	Arg	Arg	Val	Ala	Met
1				5					10					15
Asn	Lys	Glu	His	His	Asn	Gly	Asn	Phe	Thr	Asp	Pro	Ser	Ser	Val
				20					25					30
Asn	Glu	Lys	Lys	Arg	Arg	Glu	Arg	Glu	Gln	Arg	Gln	Asn	Ile	Val
				35					40					45
Leu	Trp	Arg	Gln	Pro	Leu	Ile	Thr	Leu	Gln	Tyr	Phe	Ser	Leu	Glu
				50					55					60
Ile	Leu	Val	Ile	Leu	Lys	Glu	Trp	Thr	Ser	Lys	Leu	Trp	His	Arg
				65					70					75
Gln	Ser	Ile	Val	Val	Ser	Phe	Leu	Leu	Leu	Leu	Ala	Val	Leu	Ile
				80					85					90
Ala	Thr	Tyr	Tyr	Val	Glu	Gly	Val	His	Gln	Gln	Tyr	Val	Gln	Arg
				95					100					105
Ile	Glu	Lys	Gln	Phe	Leu	Leu	Tyr	Ala	Tyr	Trp	Ile	Gly	Leu	Gly
				110					115					120
Ile	Leu	Ser	Ser	Val	Gly	Leu	Gly	Thr	Gly	Leu	His	Thr	Phe	Leu
				125					130					135
Leu	Tyr	Leu	Gly	Pro	His	Ile	Ala	Ser	Val	Thr	Leu	Ala	Ala	Tyr
				140					145					150
Glu	Cys	Asn	Ser	Val	Asn	Phe	Pro	Glu	Pro	Pro	Tyr	Pro	Asp	Gln
				155					160					165
Ile	Ile	Cys	Pro	Asp	Glu	Glu	Gly	Thr	Glu	Gly	Thr	Ile	Ser	Leu
				170					175					180
Trp	Ser	Ile	Ile	Ser	Lys	Val	Arg	Ile	Glu	Ala	Cys	Met	Trp	Gly
				185					190					195
Ile	Gly	Thr	Ala	Ile	Gly	Glu	Leu	Pro	Pro	Tyr	Phe	Met	Ala	Arg
				200					205					210
Ala	Ala	Arg	Leu	Ser	Gly	Ala	Glu	Pro	Asp	Asp	Glu	Glu	Tyr	Gln
				215					220					225
Glu	Phe	Glu	Glu	Met	Leu	Glu	His	Ala	Glu	Ser	Ala	Gln	Asp	Phe

Ala Ser Arg Ala	230	235	240
Lys Leu Ala Val Gln	Lys Leu Val Gln Lys Val		
245	250	255	
Gly Phe Phe Gly	Ile Leu Ala Cys Ala Ser Ile Pro Asn Pro Leu		
260	265	270	
Phe Asp Leu Ala	Gly Ile Thr Cys Gly His Phe Leu Val Pro Phe		
275	280	285	
Trp Thr Phe Phe	Gly Ala Thr Leu Ile Gly Lys Ala Ile Ile Lys		
290	295	300	
Met His Ile Gln	Lys Ile Phe Val Ile Ile Thr Phe Ser Lys His		
305	310	315	
Ile Val Glu Gln	Met Val Ala Phe Ile Gly Ala Val Pro Gly Ile		
320	325	330	
Gly Pro Ser Leu	Gln Lys Pro Phe Gln Gly Tyr Leu Glu Ala Gln		
335	340	345	
Arg Gln Lys Leu	His His Lys Ser Glu Met Gly Thr Pro Gln Gly		
350	355	360	
Glu Asn Trp Leu	Ser Trp Met Phe Glu Lys Leu Val Val Val Met		
365	370	375	
Val Cys Tyr Phe	Ile Leu Ser Ile Ile Asn Ser Met Ala Gln Ser		
380	385	390	
Tyr Ala Lys Arg	Ile Gln Gln Arg Leu Asn Ser Glu Glu Lys Thr		
395	400	405	

Lys

<210> 18

<211> 290

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1562471CD1

<400> 18

Met Pro Leu Leu Thr	Leu Tyr Leu Leu Leu Phe Trp Leu Ser Gly	
1	5	10
Tyr Ser Ile Ala Thr	Gln Ile Thr Gly Pro Thr Thr Val Asn Gly	15
20	25	30
Leu Glu Arg Gly Ser	Leu Thr Val Gln Cys Val Tyr Arg Ser Gly	35
35	40	45
Trp Glu Thr Tyr Leu	Lys Trp Trp Cys Arg Gly Ala Ile Trp Arg	50
50	55	60
Asp Cys Lys Ile Leu	Val Lys Thr Ser Gly Ser Glu Gln Glu Val	65
65	70	75
Lys Arg Asp Arg Val	Ser Ile Lys Asp Asn Gln Lys Asn Arg Thr	80
80	85	90
Phe Thr Val Thr Met	Glu Asp Leu Met Lys Thr Asp Ala Asp Thr	95
95	100	105
Tyr Trp Cys Gly Ile	Glu Lys Thr Gly Asn Asp Leu Gly Val Thr	110
110	115	120
Val Gln Val Thr Ile	Asp Pro Ala Pro Val Thr Gln Glu Glu Thr	125
125	130	135
Ser Ser Ser Pro Thr	Leu Thr Gly His His Leu Asp Asn Arg His	140
140	145	150
Lys Leu Leu Lys Leu	Ser Val Leu Leu Pro Leu Ile Phe Thr Ile	155
155	160	165
Leu Leu Leu Leu Leu	Val Ala Ala Ser Leu Leu Ala Trp Arg Met	170
170	175	180
Met Lys Tyr Gln Gln	Lys Ala Ala Gly Met Ser Pro Glu Gln Val	185
185	190	195
Leu Gln Pro Leu Glu	Gly Asp Leu Cys Tyr Ala Asp Leu Thr Leu	200
200	205	210

Gln	Leu	Ala	Gly	Thr	Ser	Pro	Arg	Lys	Ala	Thr	Thr	Lys	Leu	Ser
				215					220					225
Ser	Ala	Gln	Val	Asp	Gln	Val	Glu	Val	Glu	Tyr	Val	Thr	Met	Ala
				230					235					240
Ser	Leu	Pro	Lys	Glu	Asp	Ile	Ser	Tyr	Ala	Ser	Leu	Thr	Leu	Gly
				245					250					255
Ala	Glu	Asp	Gln	Glu	Pro	Thr	Tyr	Cys	Asn	Met	Gly	His	Leu	Ser
				260					265					270
Ser	His	Leu	Pro	Gly	Arg	Gly	Pro	Glu	Glu	Pro	Thr	Glu	Tyr	Ser
				275					280					285
Thr	Ile	Ser	Arg	Pro										
				290										

<210> 19

<211> 390

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1618158CD1

<400> 19

Met	Phe	Ser	Thr	Asn	Tyr	Ser	His	Met	Glu	Asn	Tyr	Arg	Lys	Arg
1				5					10					15
Glu	Asp	Leu	Val	Tyr	Gln	Ser	Thr	Val	Arg	Leu	Pro	Glu	Val	Arg
				20					25					30
Ile	Ser	Asp	Asn	Gly	Pro	Tyr	Glu	Cys	His	Val	Gly	Ile	Tyr	Asp
				35					40					45
Arg	Ala	Thr	Arg	Glu	Lys	Val	Val	Leu	Ala	Ser	Gly	Asn	Ile	Phe
				50					55					60
Leu	Asn	Val	Met	Ala	Pro	Pro	Thr	Ser	Ile	Glu	Val	Val	Ala	Ala
				65					70					75
Asp	Thr	Pro	Ala	Pro	Phe	Ser	Arg	Tyr	Gln	Ala	Gln	Asn	Phe	Thr
				80					85					90
Leu	Val	Cys	Ile	Val	Ser	Gly	Gly	Lys	Pro	Ala	Pro	Met	Val	Tyr
				95					100					105
Phe	Lys	Arg	Asp	Gly	Glu	Pro	Ile	Asp	Ala	Val	Pro	Leu	Ser	Glu
				110					115					120
Pro	Pro	Ala	Ala	Ser	Ser	Gly	Pro	Leu	Gln	Asp	Ser	Arg	Pro	Phe
				125					130					135
Arg	Ser	Leu	Leu	His	Arg	Asp	Leu	Asp	Asp	Thr	Lys	Met	Gln	Lys
				140					145					150
Ser	Leu	Ser	Leu	Leu	Asp	Ala	Glu	Asn	Arg	Gly	Gly	Arg	Pro	Tyr
				155					160					165
Thr	Glu	Arg	Pro	Ser	Arg	Gly	Leu	Thr	Pro	Asp	Pro	Asn	Ile	Leu
				170					175					180
Leu	Gln	Pro	Thr	Thr	Glu	Asn	Ile	Pro	Glu	Thr	Val	Val	Ser	Arg
				185					190					195
Glu	Phe	Pro	Arg	Trp	Val	His	Ser	Ala	Glu	Pro	Thr	Tyr	Phe	Leu
				200					205					210
Arg	His	Ser	Arg	Thr	Pro	Ser	Ser	Asp	Gly	Thr	Val	Glu	Val	Arg
				215					220					225
Ala	Leu	Leu	Thr	Trp	Thr	Leu	Asn	Pro	Gln	Ile	Asp	Asn	Glu	Ala
				230					235					240
Leu	Phe	Ser	Cys	Glu	Val	Lys	His	Pro	Ala	Leu	Ser	Met	Pro	Met
				245					250					255
Gln	Ala	Glu	Val	Thr	Leu	Val	Ala	Pro	Lys	Gly	Pro	Lys	Ile	Val
				260					265					270
Met	Thr	Pro	Ser	Arg	Ala	Arg	Val	Gly	Asp	Thr	Val	Arg	Ile	Leu
				275					280					285
Val	His	Gly	Phe	Gln	Asn	Glu	Val	Phe	Pro	Glu	Pro	Met	Phe	Thr
				290					295					300
Trp	Thr	Arg	Val	Gly	Ser	Arg	Leu	Leu	Asp	Gly	Ser	Ala	Glu	Phe

Asp Gly Lys Glu	305	Val Leu Glu Arg	310	Val Pro Ala Glu Leu	315
	320		325		330
Gly Ser Met Tyr	335	Arg Cys Thr Ala Gln	340	Asn Pro Leu Gly Ser	345
	350		355		360
Asp Thr His Thr	365	Arg Leu Ile Val Phe	370	Glu Asn Pro Asn Ile	375
	380		385		390
Arg Gly Thr Glu		Asp Ser Asn Gly Ser		Ile Gly Pro Thr Gly	
Arg Leu Thr Leu		Val Leu Ala Leu Thr		Val Ile Leu Glu Leu	

<210> 20

<211> 427

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1656935CD1

<400> 20

Met Asn Val Asn Ser	Met Asp Met Thr Gly	Gly Leu Ser Val Lys
1	10	15
Asp Pro Ser Gln Ser	Gln Ser Arg Leu Pro	Gln Trp Thr His Pro
20	25	30
Asn Ser Met Asp Asn	Leu Pro Ser Ala Ala	Ser Pro Leu Glu Gln
35	40	45
Asn Pro Ser Lys His	Gly Ala Ile Pro Gly	Gly Leu Ser Ile Gly
50	55	60
Pro Pro Gly Lys Ser	Ser Ile Asp Asp Ser	Tyr Gly Arg Tyr Asp
65	70	75
Leu Ile Gln Asn Ser	Glu Ser Pro Ala Ser	Pro Pro Val Ala Val
80	85	90
Pro His Ser Trp Ser	Arg Ala Lys Ser Asp	Ser Asp Lys Ile Ser
95	100	105
Asn Gly Ser Ser Ile	Asn Trp Pro Pro Glu	Phe His Pro Gly Val
110	115	120
Pro Trp Lys Gly Leu	Gln Asn Ile Asp Pro	Glu Asn Asp Pro Asp
125	130	135
Val Thr Pro Gly Ser	Val Pro Thr Gly Pro	Thr Ile Asn Thr Thr
140	145	150
Ile Gln Asp Val Asn	Arg Tyr Leu Leu Lys	Ser Gly Gly Ser Ser
155	160	165
Pro Pro Ser Ser Gln	Asn Ala Thr Leu Pro	Ser Ser Ser Ala Trp
170	175	180
Pro Leu Ser Ala Ser	Gly Tyr Ser Ser Ser	Phe Ser Ser Ile Ala
185	190	195
Ser Ala Pro Ser Val	Ala Gly Lys Leu Ser	Asp Ile Lys Ser Thr
200	205	210
Trp Ser Ser Gly Pro	Thr Ser His Thr Gln	Ala Ser Leu Ser His
215	220	225
Glu Leu Trp Lys Val	Pro Arg Asn Ser Thr	Ala Pro Thr Arg Pro
230	235	240
Pro Pro Gly Leu Thr	Asn Pro Lys Pro Ser	Ser Thr Trp Gly Ala
245	250	255
Ser Pro Leu Gly Trp	Thr Ser Ser Tyr Ser	Ser Gly Ser Ala Trp
260	265	270
Ser Thr Asp Thr Ser	Gly Arg Thr Ser Ser	Trp Leu Val Leu Arg
275	280	285
Asn Leu Thr Pro Gln	Ile Asp Gly Ser Lys	Leu Arg Thr Leu Cys
290	295	300
Leu Gln His Gly Pro	Leu Ile Thr Phe His	Leu Asn Leu Thr Gln

	305		310		315
Gly Asn Ala Val	Val Arg Tyr Ser Ser	Lys Glu Glu Gly Leu	Pro		
	320		325		330
Lys Ala Gln Glu	Val Leu Cys Thr Ile	Val Arg Pro Trp Glu	Thr		
	335		340		345
Leu Ser His Ser	Leu Gly Pro Ser Phe	Arg Leu Val Gly Thr	Lys		
	350		355		360
Glu Val Gly Ile	Arg Val Ser Phe Lys	Pro Pro Glu Gly Pro	Gly		
	365		370		375
Arg Ile Gly Gln	Ser Thr Ile Phe Gln	Gly Leu Ala Gln Phe	His		
	380		385		390
Asp Gln Arg Gly	Val Ser Lys Leu Thr	Gly Arg Gly Gly Ile	His		
	395		400		405
Arg Pro Arg Gly	Arg Gly Lys Ala Ser	His Gln Leu Ala His	Met		
	410		415		420
Arg His Cys Glu	Leu Thr Phe				
	425				

<210> 21

<211> 459

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1859305CD1

<400> 21

Met Glu Lys Thr Cys	Ile Asp Ala Leu Pro	Leu Thr Met Asn Ser	
1	5	10	15
Ser Glu Lys Gln Glu	Thr Val Cys Ile Phe	Gly Thr Gly Asp Phe	
	20	25	30
Gly Arg Ser Leu Gly	Leu Lys Met Leu Gln	Cys Gly Tyr Ser Val	
	35	40	45
Val Phe Gly Ser Arg	Asn Pro Gln Lys Thr	Thr Leu Leu Pro Ser	
	50	55	60
Gly Ala Glu Val Leu	Ser Tyr Ser Glu Ala	Ala Lys Lys Ser Asp	
	65	70	75
Ile Ile Ile Ile Ala	Ile His Arg Glu His	Tyr Asp Phe Leu Thr	
	80	85	90
Glu Leu Thr Glu Val	Leu Asn Gly Lys Ile	Leu Val Asp Ile Ser	
	95	100	105
Asn Asn Leu Lys Ile	Asn Gln Tyr Pro Glu	Ser Asn Ala Glu Tyr	
	110	115	120
Leu Ala His Leu Val	Pro Gly Ala His Val	Val Lys Ala Phe Asn	
	125	130	135
Thr Ile Ser Ala Trp	Ala Leu Gln Ser Gly	Ala Leu Asp Ala Ser	
	140	145	150
Arg Gln Val Phe Val	Cys Gly Asn Asp Ser	Lys Ala Lys Gln Arg	
	155	160	165
Val Met Asp Ile Val	Arg Asn Leu Gly Leu	Thr Pro Met Asp Gln	
	170	175	180
Gly Ser Leu Met Ala	Ala Lys Glu Ile Glu	Lys Tyr Pro Leu Gln	
	185	190	195
Leu Phe Pro Met Trp	Arg Phe Pro Phe Tyr	Leu Ser Ala Val Leu	
	200	205	210
Cys Val Phe Leu Phe	Phe Tyr Cys Val Ile	Arg Asp Val Ile Tyr	
	215	220	225
Pro Tyr Val Tyr Glu	Lys Lys Asp Asn Thr	Phe Arg Met Ala Ile	
	230	235	240
Ser Ile Pro Asn Arg	Ile Phe Pro Ile Thr	Ala Leu Thr Leu Leu	
	245	250	255
Ala Leu Val Tyr Leu	Pro Gly Val Ile Ala	Ala Ile Leu Gln Leu	
	260	265	270

Tyr	Arg	Gly	Thr	Lys	Tyr	Arg	Arg	Phe	Pro	Asp	Trp	Leu	Asp	His
				275					280					285
Trp	Met	Leu	Cys	Arg	Lys	Gln	Leu	Gly	Leu	Val	Ala	Leu	Gly	Phe
				290					295					300
Ala	Phe	Leu	His	Val	Leu	Tyr	Thr	Leu	Val	Ile	Pro	Ile	Arg	Tyr
				305					310					315
Tyr	Val	Arg	Trp	Arg	Leu	Gly	Asn	Leu	Thr	Val	Thr	Gln	Ala	Ile
				320					325					330
Leu	Lys	Lys	Glu	Asn	Pro	Phe	Ser	Thr	Ser	Ser	Ala	Trp	Leu	Ser
				335					340					345
Asp	Ser	Tyr	Val	Ala	Leu	Gly	Ile	Leu	Gly	Phe	Phe	Leu	Phe	Val
				350					355					360
Leu	Leu	Gly	Ile	Thr	Ser	Leu	Pro	Ser	Val	Ser	Asn	Ala	Val	Asn
				365					370					375
Trp	Arg	Glu	Phe	Arg	Phe	Val	Gln	Ser	Lys	Leu	Gly	Tyr	Leu	Thr
				380					385					390
Leu	Ile	Leu	Cys	Thr	Ala	His	Thr	Leu	Val	Tyr	Gly	Gly	Lys	Arg
				395					400					405
Phe	Leu	Ser	Pro	Ser	Asn	Leu	Arg	Trp	Tyr	Leu	Pro	Ala	Ala	Tyr
				410					415					420
Val	Leu	Gly	Leu	Ile	Ile	Pro	Cys	Thr	Val	Leu	Val	Ile	Lys	Phe
				425					430					435
Val	Leu	Ile	Met	Pro	Cys	Val	Asp	Asn	Thr	Leu	Thr	Arg	Ile	Arg
				440					445					450
Gln	Gly	Trp	Glu	Arg	Asn	Ser	Lys	His						
				455										

<210> 22

<211> 229

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1949083CD1

<400> 22

Met	Leu	Pro	Val	Ser	Arg	Thr	Cys	Leu	Leu	Glu	Ser	Ser	Thr	Arg
1				5					10					15
Leu	Lys	Pro	His	Glu	Ala	Gln	Asn	Tyr	Arg	Lys	Lys	Ala	Leu	Trp
				20					25					30
Val	Ser	Trp	Phe	Ser	Ile	Ile	Val	Thr	Leu	Ala	Leu	Ala	Val	Ala
				35					40					45
Ala	Phe	Thr	Val	Ser	Val	Met	Arg	Tyr	Ser	Ala	Ser	Ala	Phe	Gly
				50					55					60
Phe	Ala	Phe	Asp	Ala	Ile	Leu	Asp	Val	Leu	Ser	Ser	Ala	Ile	Val
				65					70					75
Leu	Trp	Arg	Tyr	Ser	Asn	Ala	Ala	Ala	Val	His	Ser	Ala	His	Arg
				80					85					90
Glu	Tyr	Ile	Ala	Cys	Val	Ile	Leu	Gly	Val	Ile	Phe	Leu	Leu	Ser
				95					100					105
Ser	Ile	Cys	Ile	Val	Val	Lys	Ala	Ile	His	Asp	Leu	Ser	Thr	Arg
				110					115					120
Leu	Leu	Pro	Glu	Val	Asp	Asp	Phe	Leu	Phe	Ser	Val	Ser	Ile	Leu
				125					130					135
Ser	Gly	Ile	Leu	Cys	Ser	Ile	Leu	Ala	Val	Leu	Lys	Phe	Met	Leu
				140					145					150
Gly	Lys	Val	Leu	Thr	Ser	Arg	Ala	Leu	Ile	Thr	Asp	Gly	Phe	Asn
				155					160					165
Ser	Leu	Val	Gly	Gly	Val	Met	Gly	Phe	Ser	Ile	Leu	Leu	Ser	Ala
				170					175					180
Glu	Val	Phe	Lys	His	Asp	Ser	Ala	Val	Trp	Tyr	Leu	Asp	Gly	Ser
				185					190					195
Ile	Gly	Val	Leu	Ile	Gly	Leu	Thr	Ile	Phe	Ala	Tyr	Gly	Val	Lys

	200		205		210
Leu Leu Ile Asp Met Val Pro Lys Val Arg Gln Thr Arg His Tyr					
	215		220		225
Glu Met Phe Glu					

<210> 23
 <211> 311
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1996357CD1

<400> 23
 Met Ala Val Asp Ile Gln Pro Ala Cys Leu Gly Leu Tyr Cys Gly
 1 5 10 15
 Lys Thr Leu Leu Phe Lys Asn Gly Ser Thr Glu Ile Tyr Gly Glu
 20 25 30
 Cys Gly Val Cys Pro Arg Gly Gln Arg Thr Asn Ala Gln Lys Tyr
 35 40 45
 Cys Gln Pro Cys Thr Glu Ser Pro Glu Leu Tyr Asp Trp Leu Tyr
 50 55 60
 Leu Gly Phe Met Ala Met Leu Pro Leu Val Leu His Trp Phe Phe
 65 70 75
 Ile Glu Trp Tyr Ser Gly Lys Lys Ser Ser Ala Leu Phe Gln
 80 85 90
 His Ile Thr Ala Leu Phe Glu Cys Ser Met Ala Ala Ile Ile Thr
 95 100 105
 Leu Leu Val Ser Asp Pro Val Gly Val Leu Tyr Ile Arg Ser Cys
 110 115 120
 Arg Val Leu Met Leu Ser Asp Trp Tyr Thr Met Leu Tyr Asn Pro
 125 130 135
 Ser Pro Asp Tyr Val Thr Thr Val His Cys Thr His Glu Ala Val
 140 145 150
 Tyr Pro Leu Tyr Thr Ile Val Phe Ile Tyr Tyr Ala Phe Cys Leu
 155 160 165
 Val Leu Met Met Leu Leu Arg Pro Leu Leu Val Lys Lys Ile Ala
 170 175 180
 Cys Gly Leu Gly Lys Ser Asp Arg Phe Lys Ser Ile Tyr Ala Ala
 185 190 195
 Leu Tyr Phe Phe Pro Ile Leu Thr Val Leu Gln Ala Val Gly Gly
 200 205 210
 Gly Leu Leu Tyr Tyr Ala Phe Pro Tyr Ile Ile Leu Val Leu Ser
 215 220 225
 Leu Val Thr Leu Ala Val Tyr Met Ser Ala Ser Glu Ile Glu Asn
 230 235 240
 Cys Tyr Asp Leu Leu Val Arg Lys Lys Arg Leu Ile Val Leu Phe
 245 250 255
 Ser His Trp Leu Leu His Ala Tyr Gly Ile Ile Ser Ile Ser Arg
 260 265 270
 Val Asp Lys Leu Glu Gln Asp Leu Pro Leu Leu Ala Leu Val Pro
 275 280 285
 Thr Pro Ala Leu Phe Tyr Leu Phe Thr Ala Lys Phe Thr Glu Pro
 290 295 300
 Ser Arg Ile Leu Ser Glu Gly Ala Asn Gly His
 305 310

<210> 24
 <211> 92
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature
 <223> Incyte ID No: 2061330CD1

<400> 24
 Met Arg Phe Ile Phe Leu Lys Phe Trp Thr Tyr Thr Val Arg Ala
 1 5 10 15
 Ser Thr Asp Leu Thr Gln Thr Gly Asp Cys Ser Gln Cys Thr His
 20 25 30
 Gln Val Thr Glu Val Gly Gln Gln Ile Lys Thr Ile Phe Leu Phe
 35 40 45
 Tyr Ser Tyr Tyr Glu Cys Met Glu Thr Ile Lys Glu Thr Cys Leu
 50 55 60
 Tyr Asn Ala Thr Gln Tyr Lys Val Cys Ser Pro Arg Asn Asp Arg
 65 70 75
 Pro Asp Val Cys Tyr Asn Pro Ser Glu Pro Pro Ala Pro Pro Phe
 80 85 90
 Leu Lys

<210> 25
 <211> 258
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2346947CD1

<400> 25
 Met Ala Glu Ser Pro Gly Cys Cys Ser Val Trp Ala Arg Cys Leu
 1 5 10 15
 His Cys Leu Tyr Ser Cys His Trp Arg Lys Cys Pro Arg Glu Arg
 20 25 30
 Met Gln Thr Ser Lys Cys Asp Cys Ile Trp Phe Gly Leu Leu Phe
 35 40 45
 Leu Thr Phe Leu Leu Ser Leu Ser Trp Leu Tyr Ile Gly Leu Val
 50 55 60
 Leu Leu Asn Asp Leu His Asn Phe Asn Glu Phe Leu Phe Arg Arg
 65 70 75
 Trp Gly His Trp Met Asp Trp Ser Leu Ala Phe Leu Leu Val Ile
 80 85 90
 Ser Leu Leu Val Thr Tyr Ala Ser Leu Leu Leu Val Leu Ala Leu
 95 100 105
 Leu Leu Arg Leu Cys Arg Gln Pro Leu His Leu His Ser Leu His
 110 115 120
 Lys Val Leu Leu Leu Leu Ile Met Leu Leu Val Ala Ala Gly Leu
 125 130 135
 Val Gly Leu Asp Ile Gln Trp Gln Gln Glu Trp His Ser Leu Arg
 140 145 150
 Val Ser Leu Gln Ala Thr Ala Pro Phe Leu His Ile Gly Ala Ala
 155 160 165
 Ala Gly Ile Ala Leu Leu Ala Trp Pro Val Ala Asp Thr Phe Tyr
 170 175 180
 Arg Ile His Arg Arg Gly Pro Lys Ile Leu Leu Leu Leu Leu Phe
 185 190 195
 Phe Gly Val Val Leu Val Ile Tyr Leu Ala Pro Leu Cys Ile Ser
 200 205 210
 Ser Pro Cys Ile Met Glu Pro Arg Asp Leu Pro Pro Lys Pro Gly
 215 220 225
 Leu Val Gly His Arg Gly Ala Pro Met Leu Ala Pro Glu Asn Thr
 230 235 240
 Leu Met Ser Leu Arg Lys Thr Ala Glu Cys Gly Leu Leu Cys Leu
 245 250 255
 Arg Leu Met

<210> 26
 <211> 226
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2795577CD1

<400> 26
 Met Lys Met Val Ala Pro Trp Thr Arg Phe Tyr Ser Asn Ser Cys
 1 5 10 15
 Cys Leu Cys Cys His Val Arg Thr Gly Thr Ile Leu Leu Gly Val
 20 25 30
 Trp Tyr Leu Ile Ile Asn Ala Val Val Leu Leu Ile Leu Leu Ser
 35 40 45
 Ala Leu Ala Asp Pro Asp Gln Tyr Asn Phe Ser Ser Ser Glu Leu
 50 55 60
 Gly Gly Asp Phe Glu Phe Met Asp Asp Ala Asn Met Cys Ile Ala
 65 70 75
 Ile Ala Ile Ser Leu Leu Met Ile Leu Ile Cys Ala Met Ala Thr
 80 85 90
 Tyr Gly Ala Tyr Lys Gln Arg Ala Ala Trp Ile Ile Pro Phe Phe
 95 100 105
 Cys Tyr Gln Ile Phe Asp Phe Ala Leu Asn Met Leu Val Ala Ile
 110 115 120
 Thr Val Leu Ile Tyr Pro Asn Ser Ile Gln Glu Tyr Ile Arg Gln
 125 130 135
 Leu Pro Pro Asn Phe Pro Tyr Arg Asp Asp Val Met Ser Val Asn
 140 145 150
 Pro Thr Cys Leu Val Leu Ile Ile Leu Leu Phe Ile Ser Ile Ile
 155 160 165
 Leu Thr Phe Lys Gly Tyr Leu Ile Ser Cys Val Trp Asn Cys Tyr
 170 175 180
 Arg Tyr Ile Asn Gly Arg Asn Ser Ser Asp Val Leu Val Tyr Val
 185 190 195
 Thr Ser Asn Asp Thr Thr Val Leu Leu Pro Pro Tyr Asp Asp Ala
 200 205 210
 Thr Val Asn Gly Ala Ala Lys Glu Pro Pro Pro Pro Tyr Val Ser
 215 220 225
 Ala

<210> 27
 <211> 136
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3255825CD1

<400> 27
 Met Ile Ser Ile Thr Glu Trp Gln Lys Ile Gly Val Gly Ile Thr
 1 5 10 15
 Gly Phe Gly Ile Phe Phe Ile Leu Phe Gly Thr Leu Leu Tyr Phe
 20 25 30
 Asp Ser Val Leu Leu Ala Phe Gly Asn Leu Leu Phe Leu Thr Gly
 35 40 45
 Leu Ser Leu Ile Ile Gly Leu Arg Lys Thr Phe Trp Phe Phe Phe
 50 55 60
 Gln Arg His Lys Leu Lys Gly Thr Ser Phe Leu Leu Gly Gly Val
 65 70 75

Val	Ile	Val	Leu	Leu	Arg	Trp	Pro	Leu	Leu	Gly	Met	Phe	Leu	Glu
				80					85					90
Thr	Tyr	Gly	Phe	Phe	Ser	Leu	Phe	Lys	Gly	Phe	Phe	Pro	Val	Ala
				95					100					105
Phe	Gly	Ser	Trp	Ala	Met	Ser	Ala	Thr	Ser	Pro	Ser	Trp	Val	Arg
				110					115					120
Cys	Ser	Gly	Asp	Phe	Lys	Ala	Leu	Ala	Arg	Trp	Ser	Glu	Lys	Gln
				125					130					135

Arg

<210> 28

<211> 458

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3393430CD1

<400> 28

Met	Ala	Trp	Ala	Ser	Arg	Leu	Gly	Leu	Leu	Leu	Ala	Leu	Leu	Leu
1				5					10					15
Pro	Val	Val	Gly	Ala	Ser	Thr	Pro	Gly	Thr	Val	Val	Arg	Leu	Asn
				20					25					30
Lys	Ala	Ala	Leu	Ser	Tyr	Val	Ser	Glu	Ile	Gly	Lys	Ala	Pro	Leu
				35					40					45
Gln	Arg	Ala	Leu	Gln	Val	Thr	Val	Pro	His	Phe	Leu	Asp	Trp	Ser
				50					55					60
Gly	Glu	Ala	Leu	Gln	Pro	Thr	Arg	Ile	Arg	Ile	Leu	Asn	Val	His
				65					70					75
Val	Pro	Arg	Leu	His	Leu	Lys	Phe	Ile	Ala	Gly	Phe	Gly	Val	Arg
				80					85					90
Leu	Leu	Ala	Ala	Ala	Asn	Phe	Thr	Phe	Lys	Val	Phe	Arg	Ala	Pro
				95					100					105
Glu	Pro	Leu	Glu	Leu	Thr	Leu	Pro	Val	Glu	Leu	Leu	Ala	Asp	Thr
				110					115					120
Arg	Val	Thr	Gln	Ser	Ser	Ile	Arg	Thr	Pro	Val	Val	Ser	Ile	Ser
				125					130					135
Ala	Cys	Ser	Leu	Phe	Ser	Gly	His	Ala	Asn	Glu	Phe	Asp	Gly	Ser
				140					145					150
Asn	Ser	Thr	Ser	His	Ala	Leu	Leu	Val	Leu	Val	Gln	Lys	His	Ile
				155					160					165
Lys	Ala	Val	Leu	Ser	Asn	Lys	Leu	Cys	Leu	Ser	Ile	Ser	Asn	Leu
				170					175					180
Val	Gln	Gly	Val	Asn	Val	His	Leu	Gly	Thr	Leu	Ile	Gly	Leu	Asn
				185					190					195
Pro	Val	Gly	Pro	Glu	Ser	Gln	Ile	Arg	Tyr	Ser	Met	Val	Ser	Val
				200					205					210
Pro	Thr	Val	Thr	Ser	Asp	Tyr	Ile	Ser	Leu	Glu	Val	Asn	Ala	Val
				215					220					225
Leu	Phe	Leu	Leu	Gly	Lys	Pro	Ile	Ile	Leu	Pro	Thr	Asp	Ala	Thr
				230					235					240
Pro	Phe	Val	Leu	Pro	Arg	His	Val	Gly	Thr	Glu	Gly	Ser	Met	Ala
				245					250					255
Thr	Val	Gly	Leu	Ser	Gln	Gln	Leu	Phe	Asp	Ser	Ala	Leu	Leu	Leu
				260					265					270
Leu	Gln	Lys	Ala	Gly	Ala	Leu	Asn	Leu	Asp	Ile	Thr	Gly	Gln	Leu
				275					280					285
Arg	Ser	Asp	Asp	Asn	Leu	Leu	Asn	Thr	Ser	Ala	Leu	Gly	Arg	Leu
				290					295					300
Ile	Pro	Glu	Val	Ala	Arg	Gln	Phe	Pro	Glu	Pro	Met	Pro	Val	Val
				305					310					315
Leu	Lys	Val	Arg	Leu	Gly	Ala	Thr	Pro	Val	Ala	Met	Leu	His	Thr

320	325	330
Asn Asn Ala Thr Leu Arg Leu Gln Pro Phe Val Glu Val Leu Ala		
335	340	345
Thr Ala Ser Asn Ser Ala Phe Gln Ser Leu Phe Ser Leu Asp Val		
350	355	360
Val Val Asn Leu Arg Leu Gln Leu Ser Val Ser Lys Val Lys Leu		
365	370	375
Gln Gly Thr Thr Ser Val Leu Gly Asp Val Gln Leu Thr Val Ala		
380	385	390
Ser Ser Asn Val Gly Phe Ile Asp Thr Asp Gln Val Arg Thr Leu		
395	400	405
Met Gly Thr Val Phe Glu Lys Pro Leu Leu Asp His Leu Asn Ala		
410	415	420
Leu Leu Ala Met Gly Ile Ala Leu Pro Gly Val Val Asn Leu His		
425	430	435
Tyr Val Ala Pro Glu Ile Phe Val Tyr Glu Gly Tyr Val Val Ile		
440	445	450
Ser Ser Gly Leu Phe Tyr Gln Ser		
455		

<210> 29

<211> 368

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3490990CD1

<400> 29

Met Phe Gly Gln Asn Leu Glu Val Gln Leu Ser Ser Ala Arg Thr		
1 5 10 15		
Glu Asn Thr Thr Val Val Trp Lys Ser Phe His Asp Ser Ile Thr		
20 25 30		
Leu Ile Val Leu Ser Ser Glu Val Gly Ile Ser Glu Leu Arg Leu		
35 40 45		
Glu Arg Leu Leu Gln Met Val Phe Gly Ala Met Val Leu Leu Val		
50 55 60		
Gly Leu Glu Glu Leu Thr Asn Ile Arg Asn Val Glu Arg Leu Lys		
65 70 75		
Lys Asp Leu Arg Ala Ser Tyr Cys Leu Ile Asp Ser Phe Leu Gly		
80 85 90		
Asp Ser Glu Leu Ile Gly Asp Leu Thr Gln Cys Val Asp Cys Val		
95 100 105		
Ile Pro Pro Glu Gly Ser Leu Leu Gln Glu Ala Leu Ser Gly Phe		
110 115 120		
Ala Glu Ala Ala Gly Thr Thr Phe Val Ser Leu Val Val Ser Gly		
125 130 135		
Arg Val Val Ala Ala Thr Glu Gly Trp Trp Arg Leu Gly Thr Pro		
140 145 150		
Glu Ala Val Leu Leu Pro Trp Leu Val Gly Ser Leu Pro Pro Gln		
155 160 165		
Thr Ala Arg Asp Tyr Pro Val Tyr Leu Pro His Gly Ser Pro Thr		
170 175 180		
Val Pro His Arg Leu Leu Thr Leu Thr Leu Leu Pro Ser Leu Glu		
185 190 195		
Leu Cys Leu Leu Cys Gly Pro Ser Pro Pro Leu Ser Gln Leu Tyr		
200 205 210		
Pro Gln Leu Leu Glu Arg Trp Trp Gln Pro Leu Leu Asp Pro Leu		
215 220 225		
Arg Ala Cys Leu Pro Leu Gly Pro Arg Ala Leu Pro Ser Gly Phe		
230 235 240		
Pro Leu His Thr Asp Ile Leu Gly Leu Leu Leu His Leu Glu		
245 250 255		


```

Leu Lys Arg Cys Leu Phe Thr Val Glu Pro Leu Gly Asp Lys Glu
260 265 270
Pro Ser Pro Glu Gln Arg Arg Arg Leu Leu Arg Asn Phe Tyr Thr
275 280 285
Leu Val Thr Ser Thr His Phe Pro Pro Glu Pro Gly Pro Pro Glu
290 295 300
Lys Thr Glu Asp Glu Val Tyr Gln Ala Gln Leu Pro Arg Ala Cys
305 310 315
Tyr Leu Val Leu Gly Thr Glu Glu Pro Gly Thr Gly Val Arg Leu
320 325 330
Val Ala Leu Gln Leu Gly Leu Arg Arg Leu Leu Leu Leu Leu Ser
335 340 345
Pro Gln Ser Pro Thr His Gly Leu Arg Ser Leu Ala Thr His Thr
350 355 360
Leu His Ala Leu Thr Pro Leu Leu
365

```

<210> 30

<211> 91

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3635154CD1

<400> 30

```

Met Tyr Gly Lys Ile Ile Phe Val Leu Leu Leu Ser Glu Ile Val
1 5 10 15
Ser Ile Ser Ala Ser Ser Thr Thr Gly Val Ala Met His Thr Ser
20 25 30
Thr Ser Ser Ser Val Thr Lys Ser Tyr Ile Ser Ser Gln Thr Asn
35 40 45
Gly Glu Thr Gly Gln Leu Val His Arg Phe Thr Val Pro Ala Pro
50 55 60
Val Val Ile Ile Leu Ile Ile Leu Cys Val Met Ala Gly Ile Ile
65 70 75
Gly Thr Ile Leu Leu Phe Ser Tyr Ser Phe Arg Arg Leu Ile Lys
80 85 90
Gly

```

<210> 31

<211> 295

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4374347CD1

<400> 31

```

Met Gly Pro Pro Ser Ala Cys Pro His Arg Glu Cys Ile Pro Trp
1 5 10 15
Gln Gly Leu Leu Leu Thr Ala Ser Leu Leu Thr Phe Trp Asn Ala
20 25 30
Pro Thr Thr Ala Trp Leu Phe Ile Ala Ser Ala Pro Phe Glu Val
35 40 45
Ala Glu Gly Glu Asn Val His Leu Ser Val Val Tyr Leu Pro Glu
50 55 60
Asn Leu Tyr Ser Tyr Gly Trp Tyr Lys Gly Lys Thr Val Glu Pro
65 70 75
Asn Gln Leu Ile Ala Ala Tyr Val Ile Asp Thr His Val Arg Thr
80 85 90
Pro Gly Pro Ala Tyr Ser Gly Arg Glu Thr Ile Ser Pro Ser Gly

```

Asp	Leu	His	Phe	Gln	Asn	Val	Thr	Leu	Glu	Asp	Thr	Gly	Tyr	Tyr	105
				110					115						120
Asn	Leu	Gln	Val	Thr	Tyr	Arg	Asn	Ser	Gln	Ile	Glu	Gln	Ala	Ser	
				125					130						135
His	His	Leu	Arg	Val	Tyr	Glu	Ser	Val	Ala	Gln	Pro	Ser	Ile	Gln	
				140					145						150
Ala	Ser	Ser	Thr	Thr	Val	Thr	Glu	Lys	Gly	Ser	Val	Val	Leu	Thr	
				155					160						165
Cys	His	Thr	Asn	Asn	Thr	Gly	Thr	Ser	Phe	Gln	Trp	Ile	Phe	Asn	
				170					175						180
Asn	Gln	Arg	Leu	Gln	Val	Thr	Lys	Arg	Met	Lys	Leu	Ser	Trp	Phe	
				185					190						195
Asn	His	Val	Leu	Thr	Ile	Asp	Pro	Ile	Arg	Gln	Glu	Asp	Ala	Gly	
				200					205						210
Glu	Tyr	Gln	Cys	Glu	Val	Ser	Asn	Pro	Val	Ser	Ser	Asn	Arg	Ser	
				215					220						225
Asp	Pro	Leu	Lys	Leu	Thr	Val	Lys	Tyr	Asp	Asn	Thr	Leu	Gly	Ile	
				230					235						240
Leu	Ile	Gly	Val	Leu	Val	Gly	Ser	Leu	Leu	Val	Ala	Ala	Leu	Val	
				245					250						255
Cys	Phe	Leu	Leu	Leu	Arg	Lys	Thr	Gly	Arg	Ala	Ser	Asp	Gln	Ser	
				260					265						270
Asp	Phe	Arg	Glu	Gln	Gln	Pro	Pro	Ala	Ser	Thr	Pro	Gly	His	Gly	
				275					280						285
Pro	Ser	Asp	Ser	Ser	Asp	Ser	Ser	Ile	Ser						
				290					295						

<210> 32

<211> 724

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4596747CD1

<400> 32

Met	Phe	Asp	Thr	Thr	Pro	His	Ser	Gly	Arg	Ser	Thr	Pro	Ser	Ser	
1				5					10					15	
Ser	Pro	Ser	Leu	Arg	Lys	Arg	Leu	Gln	Leu	Leu	Pro	Pro	Ser	Arg	
				20					25					30	
Pro	Pro	Pro	Glu	Pro	Glu	Pro	Gly	Thr	Met	Val	Glu	Lys	Gly	Ser	
				35					40					45	
Asp	Ser	Ser	Ser	Glu	Lys	Gly	Gly	Val	Pro	Gly	Thr	Pro	Ser	Thr	
				50					55					60	
Gln	Ser	Leu	Gly	Ser	Arg	Asn	Phe	Ile	Arg	Asn	Ser	Lys	Lys	Met	
				65					70					75	
Gln	Ser	Trp	Tyr	Ser	Met	Leu	Ser	Pro	Thr	Tyr	Lys	Gln	Arg	Asn	
				80					85					90	
Glu	Asp	Phe	Arg	Lys	Leu	Phe	Ser	Lys	Leu	Pro	Glu	Ala	Glu	Arg	
				95					100					105	
Leu	Ile	Val	Asp	Tyr	Ser	Cys	Ala	Leu	Gln	Arg	Glu	Ile	Leu	Leu	
				110					115					120	
Gln	Gly	Arg	Leu	Tyr	Leu	Ser	Glu	Asn	Trp	Ile	Cys	Phe	Tyr	Ser	
				125					130					135	
Asn	Ile	Phe	Arg	Trp	Glu	Thr	Thr	Ile	Ser	Ile	Gln	Leu	Lys	Glu	
				140					145					150	
Val	Thr	Cys	Leu	Lys	Lys	Glu	Lys	Thr	Ala	Lys	Leu	Ile	Pro	Asn	
				155					160					165	
Ala	Ile	Gln	Ile	Cys	Thr	Glu	Ser	Glu	Lys	His	Phe	Phe	Thr	Ser	
				170					175					180	
Phe	Gly	Ala	Arg	Asp	Arg	Cys	Phe	Leu	Leu	Ile	Phe	Arg	Leu	Trp	
				185					190					195	

Gln	Asn	Ala	Leu	Leu	Glu	Lys	Thr	Leu	Ser	Pro	Arg	Glu	Leu	Trp
				200					205					210
His	Leu	Val	His	Gln	Cys	Tyr	Gly	Ser	Glu	Leu	Gly	Leu	Thr	Ser
				215					220					225
Glu	Asp	Glu	Asp	Tyr	Val	Ser	Pro	Leu	Gln	Leu	Asn	Gly	Leu	Gly
				230					235					240
Thr	Pro	Lys	Glu	Val	Gly	Asp	Val	Ile	Ala	Leu	Ser	Asp	Ile	Thr
				245					250					255
Ser	Ser	Gly	Ala	Ala	Asp	Arg	Ser	Gln	Glu	Pro	Ser	Pro	Val	Gly
				260					265					270
Ser	Arg	Arg	Gly	His	Val	Thr	Pro	Asn	Leu	Ser	Arg	Ala	Ser	Ser
				275					280					285
Asp	Ala	Asp	His	Gly	Ala	Glu	Glu	Asp	Lys	Glu	Glu	Gln	Val	Asp
				290					295					300
Ser	Gln	Pro	Asp	Ala	Ser	Ser	Ser	Gln	Thr	Val	Thr	Pro	Val	Ala
				305					310					315
Glu	Pro	Pro	Ser	Thr	Glu	Pro	Thr	Gln	Pro	Asp	Gly	Pro	Thr	Thr
				320					325					330
Leu	Gly	Pro	Leu	Asp	Leu	Leu	Pro	Ser	Glu	Glu	Leu	Leu	Thr	Asp
				335					340					345
Thr	Ser	Asn	Ser	Ser	Ser	Ser	Thr	Gly	Glu	Glu	Ala	Asp	Leu	Ala
				350					355					360
Ala	Leu	Leu	Pro	Asp	Leu	Ser	Gly	Arg	Leu	Leu	Ile	Asn	Ser	Val
				365					370					375
Phe	His	Val	Gly	Ala	Glu	Arg	Leu	Gln	Gln	Met	Leu	Phe	Ser	Asp
				380					385					390
Ser	Pro	Phe	Leu	Gln	Gly	Phe	Leu	Gln	Gln	Cys	Lys	Phe	Thr	Asp
				395					400					405
Val	Thr	Leu	Ser	Pro	Trp	Ser	Gly	Asp	Ser	Lys	Cys	His	Gln	Arg
				410					415					420
Arg	Val	Leu	Thr	Tyr	Thr	Ile	Pro	Ile	Ser	Asn	Pro	Leu	Gly	Pro
				425					430					435
Lys	Ser	Ala	Ser	Val	Val	Glu	Thr	Gln	Thr	Leu	Phe	Arg	Arg	Gly
				440					445					450
Pro	Gln	Ala	Gly	Gly	Cys	Val	Val	Asp	Ser	Glu	Val	Leu	Thr	Gln
				455					460					465
Gly	Ile	Pro	Tyr	Gln	Asp	Tyr	Phe	Tyr	Thr	Ala	His	Arg	Tyr	Cys
				470					475					480
Ile	Leu	Gly	Leu	Ala	Arg	Asn	Lys	Ala	Arg	Leu	Arg	Val	Ser	Ser
				485					490					495
Glu	Ile	Arg	Tyr	Arg	Lys	Gln	Pro	Trp	Ser	Leu	Val	Lys	Ser	Leu
				500					505					510
Ile	Glu	Lys	Asn	Ser	Trp	Ser	Gly	Ile	Glu	Asp	Tyr	Phe	His	His
				515					520					525
Leu	Glu	Arg	Glu	Leu	Ala	Lys	Ala	Glu	Lys	Leu	Ser	Leu	Glu	Glu
				530					535					540
Gly	Gly	Lys	Asp	Ala	Arg	Gly	Leu	Leu	Ser	Gly	Leu	Arg	Arg	Arg
				545					550					555
Lys	Arg	Pro	Leu	Ser	Trp	Arg	Ala	His	Gly	Asp	Gly	Pro	Gln	His
				560					565					570
Pro	Asp	Pro	Asp	Pro	Cys	Ala	Arg	Ala	Gly	Ile	His	Thr	Ser	Gly
				575					580					585
Ser	Leu	Ser	Ser	Arg	Phe	Ser	Glu	Pro	Ser	Val	Asp	Gln	Gly	Pro
				590					595					600
Gly	Ala	Gly	Ile	Pro	Ser	Ala	Leu	Val	Leu	Ile	Ser	Ile	Val	Ile
				605					610					615
Cys	Val	Ser	Leu	Ile	Ile	Leu	Ile	Ala	Leu	Asn	Val	Leu	Leu	Phe
				620					625					630
Tyr	Arg	Leu	Trp	Ser	Leu	Glu	Arg	Thr	Ala	His	Thr	Phe	Glu	Ser
				635					640					645
Trp	His	Ser	Leu	Ala	Leu	Ala	Lys	Gly	Lys	Phe	Pro	Gln	Thr	Ala
				650					655					660
Thr	Glu	Trp	Ala	Glu	Ile	Leu	Ala	Leu	Gln	Lys	Gln	Phe	His	Ser

Val	Glu	Val	His	665	Trp	Arg	Gln	Ile	670	Leu	Arg	Ala	Ser	Val	675	Glu
Leu	Leu	Asp	Glu	680	Met	Lys	Phe	Ser	685	Glu	Lys	Leu	His	Gln	690	Gly
Ile	Thr	Val	Ser	695	Asp	Pro	Pro	Phe	700	Thr	Gln	Pro	Arg	Pro	705	Asp
Asp	Ser	Phe	Ser	710					715						720	

<210> 33

<211> 331

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5052680CD1

<400> 33

Met	Arg	Pro	Ala	Leu	Ala	Val	Gly	Leu	Val	Phe	Ala	Gly	Cys	Cys		
1				5					10					15		
Ser	Asn	Val	Ile	Phe	Leu	Glu	Leu	Leu	Ala	Arg	Lys	His	Pro	Gly		
				20					25					30		
Cys	Gly	Asn	Ile	Val	Thr	Phe	Ala	Gln	Phe	Leu	Phe	Ile	Ala	Val		
				35					40					45		
Glu	Gly	Phe	Leu	Phe	Glu	Ala	Asp	Leu	Gly	Arg	Lys	Pro	Pro	Ala		
				50					55					60		
Ile	Pro	Ile	Arg	Tyr	Tyr	Ala	Ile	Met	Val	Thr	Met	Phe	Phe	Thr		
				65					70					75		
Val	Ser	Val	Val	Asn	Asn	Tyr	Ala	Leu	Asn	Leu	Asn	Ile	Ala	Met		
				80					85					90		
Pro	Leu	His	Met	Ile	Phe	Arg	Ser	Gly	Ser	Leu	Ile	Ala	Asn	Met		
				95					100					105		
Ile	Leu	Gly	Ile	Ile	Ile	Leu	Lys	Lys	Arg	Tyr	Ser	Ile	Phe	Lys		
				110					115					120		
Tyr	Thr	Ser	Ile	Ala	Leu	Val	Ser	Val	Gly	Ile	Phe	Ile	Cys	Thr		
				125					130					135		
Phe	Met	Ser	Ala	Lys	Gln	Val	Thr	Ser	Gln	Ser	Ser	Leu	Ser	Glu		
				140					145					150		
Asn	Asp	Gly	Phe	Gln	Ala	Phe	Val	Trp	Trp	Leu	Leu	Gly	Ile	Gly		
				155					160					165		
Ala	Leu	Thr	Phe	Ala	Leu	Leu	Met	Ser	Ala	Arg	Met	Gly	Ile	Phe		
				170					175					180		
Gln	Glu	Thr	Leu	Tyr	Lys	Arg	Phe	Gly	Lys	His	Ser	Lys	Glu	Ala		
				185					190					195		
Leu	Phe	Tyr	Asn	His	Ala	Leu	Pro	Leu	Pro	Gly	Phe	Val	Phe	Leu		
				200					205					210		
Ala	Ser	Asp	Ile	Tyr	Asp	His	Ala	Val	Leu	Phe	Asn	Lys	Ser	Glu		
				215					220					225		
Leu	Tyr	Glu	Ile	Pro	Val	Ile	Gly	Val	Thr	Leu	Pro	Ile	Met	Trp		
				230					235					240		
Phe	Tyr	Leu	Leu	Met	Asn	Ile	Ile	Thr	Gln	Tyr	Val	Cys	Ile	Arg		
				245					250					255		
Gly	Val	Phe	Ile	Leu	Thr	Thr	Glu	Cys	Ala	Ser	Leu	Thr	Val	Thr		
				260					265					270		
Leu	Val	Val	Thr	Leu	Arg	Lys	Phe	Val	Ser	Leu	Ile	Phe	Ser	Ile		
				275					280					285		
Leu	Tyr	Phe	Gln	Asn	Pro	Phe	Thr	Leu	Trp	His	Trp	Leu	Gly	Thr		
				290					295					300		
Leu	Phe	Val	Phe	Ile	Gly	Thr	Leu	Met	Tyr	Thr	Glu	Val	Trp	Asn		
				305					310					315		
Asn	Leu	Gly	Thr	Thr	Lys	Ser	Glu	Pro	Gln	Lys	Asp	Ser	Lys	Lys		
				320					325					330		

Asn

<210> 34

<211> 398

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5373575CD1

<400> 34

Met	Leu	Gly	Arg	Ser	Gly	Tyr	Arg	Ala	Leu	Pro	Leu	Gly	Asp	Phe
1				5					10					15
Asp	Arg	Phe	Gln	Gln	Ser	Ser	Phe	Gly	Phe	Leu	Gly	Ser	Gln	Lys
				20					25					30
Gly	Cys	Leu	Ser	Pro	Glu	Arg	Gly	Gly	Val	Gly	Thr	Gly	Ala	Asp
				35					40					45
Val	Pro	Gln	Ser	Trp	Pro	Ser	Cys	Leu	Cys	His	Gly	Leu	Ile	Ser
				50					55					60
Phe	Leu	Gly	Phe	Leu	Leu	Leu	Val	Thr	Phe	Pro	Ile	Ser	Gly	
				65					70					75
Trp	Phe	Ala	Leu	Lys	Ile	Val	Pro	Thr	Tyr	Glu	Arg	Met	Ile	Val
				80					85					90
Phe	Arg	Leu	Gly	Arg	Ile	Arg	Thr	Pro	Gln	Gly	Pro	Gly	Met	Val
				95					100					105
Leu	Leu	Leu	Pro	Phe	Ile	Asp	Ser	Phe	Gln	Arg	Val	Asp	Leu	Arg
				110					115					120
Thr	Arg	Ala	Phe	Asn	Val	Pro	Pro	Cys	Lys	Leu	Ala	Ser	Lys	Asp
				125					130					135
Gly	Ala	Val	Leu	Ser	Val	Gly	Ala	Asp	Val	Gln	Phe	Arg	Ile	Trp
				140					145					150
Asp	Pro	Val	Leu	Ser	Val	Met	Thr	Val	Lys	Asp	Leu	Asn	Thr	Ala
				155					160					165
Thr	Arg	Met	Thr	Ala	Gln	Asn	Ala	Met	Thr	Lys	Ala	Leu	Leu	Lys
				170					175					180
Arg	Pro	Leu	Arg	Glu	Ile	Gln	Met	Glu	Lys	Leu	Lys	Ile	Ser	Asp
				185					190					195
Gln	Leu	Leu	Leu	Glu	Ile	Asn	Asp	Val	Thr	Arg	Ala	Trp	Gly	Leu
				200					205					210
Glu	Val	Asp	Arg	Val	Glu	Leu	Ala	Val	Glu	Ala	Val	Leu	Gln	Pro
				215					220					225
Pro	Gln	Asp	Ser	Pro	Ala	Gly	Pro	Asn	Leu	Asp	Ser	Thr	Leu	Gln
				230					235					240
Gln	Leu	Ala	Leu	His	Phe	Leu	Gly	Gly	Ser	Met	Asn	Ser	Met	Ala
				245					250					255
Gly	Gly	Ala	Pro	Ser	Pro	Gly	Pro	Ala	Asp	Thr	Val	Glu	Met	Val
				260					265					270
Ser	Glu	Val	Glu	Pro	Pro	Ala	Pro	Gln	Val	Gly	Ala	Arg	Ser	Ser
				275					280					285
Pro	Lys	Gln	Pro	Leu	Ala	Glu	Gly	Leu	Leu	Thr	Ala	Leu	Gln	Pro
				290					295					300
Phe	Leu	Ser	Glu	Ala	Leu	Val	Ser	Gln	Val	Gly	Ala	Cys	Tyr	Gln
				305					310					315
Phe	Asn	Val	Val	Leu	Pro	Ser	Gly	Thr	Gln	Ser	Ala	Tyr	Phe	Leu
				320					325					330
Asp	Leu	Thr	Thr	Gly	Arg	Gly	Arg	Val	Gly	His	Gly	Val	Pro	Asp
				335					340					345
Gly	Ile	Pro	Asp	Val	Val	Val	Glu	Met	Ala	Glu	Ala	Asp	Leu	Arg
				350					355					360
Ala	Leu	Leu	Cys	Arg	Glu	Leu	Arg	Pro	Leu	Gly	Ala	Tyr	Met	Ser
				365					370					375
Gly	Arg	Leu	Lys	Val	Lys	Gly	Asp	Leu	Ala	Met	Ala	Met	Lys	Leu

380
Glu Ala Val Leu Arg Ala Leu Lys
395

385

390

<210> 35
<211> 220
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5524468CD1

<400> 35
Met Thr Trp Leu Val Leu Leu Gly Thr Leu Leu Cys Met Leu Arg
1 5 10 15
Val Gly Leu Gly Thr Pro Asp Ser Glu Gly Phe Pro Pro Arg Ala
20 25 30
Leu His Asn Cys Pro Tyr Lys Cys Ile Cys Ala Ala Asp Leu Leu
35 40 45
Ser Cys Thr Gly Leu Gly Leu Gln Asp Val Pro Ala Glu Leu Pro
50 55 60
Ala Ala Thr Ala Asp Leu Asp Leu Ser His Asn Ala Leu Gln Arg
65 70 75
Leu Arg Pro Gly Trp Leu Ala Pro Leu Phe Gln Leu Arg Ala Leu
80 85 90
His Leu Asp His Asn Glu Leu Asp Ala Leu Gly Arg Gly Val Phe
95 100 105
Val Asn Ala Ser Gly Leu Arg Leu Leu Asp Leu Ser Ser Asn Thr
110 115 120
Leu Arg Ala Leu Gly Arg His Asp Leu Asp Gly Leu Gly Ala Leu
125 130 135
Glu Lys Leu Leu Leu Phe Asn Asn Arg Leu Val His Leu Asp Glu
140 145 150
His Ala Phe His Gly Leu Arg Ala Leu Ser His Leu Tyr Leu Gly
155 160 165
Cys Asn Glu Leu Ala Ser Phe Ser Phe Asp His Leu His Gly Leu
170 175 180
Ser Ala Thr His Leu Leu Thr Leu Asp Leu Ser Ser Asn Arg Leu
185 190 195
Gly His Ile Ser Val Pro Glu Leu Ala Ala Leu Pro Ala Phe Leu
200 205 210
Lys Asn Gly Leu Tyr Leu His Asp Asn Thr
215 220

<210> 36
<211> 706
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5944279CD1

<400> 36
Met Glu Glu Asn Pro Thr Leu Glu Ser Glu Ala Trp Gly Ser Ser
1 5 10 15
Arg Gly Trp Leu Ala Pro Arg Glu Ala Arg Gly Gly Pro Ser Leu
20 25 30
Ser Ser Val Leu Asn Glu Leu Pro Ser Ala Ala Thr Leu Arg Tyr
35 40 45
Arg Asp Pro Gly Val Leu Pro Trp Gly Ala Leu Glu Glu Glu Glu
50 55 60
Glu Asp Gly Gly Arg Ser Arg Lys Ala Phe Thr Glu Val Thr Gln
65 70 75

Thr	Glu	Leu	Gln	Asp	Pro	His	Pro	Ser	Arg	Glu	Leu	Pro	Trp	Pro
				80					85					90
Met	Gln	Ala	Arg	Arg	Ala	His	Arg	Gln	Arg	Asn	Ala	Ser	Arg	Asp
				95					100					105
Gln	Val	Val	Tyr	Gly	Ser	Gly	Thr	Lys	Thr	Asp	Arg	Trp	Ala	Arg
				110					115					120
Leu	Leu	Arg	Arg	Ser	Lys	Glu	Lys	Thr	Lys	Glu	Gly	Leu	Arg	Ser
				125					130					135
Leu	Gln	Pro	Trp	Ala	Trp	Thr	Leu	Lys	Arg	Ile	Gly	Gly	Gln	Phe
				140					145					150
Gly	Ala	Gly	Thr	Glu	Ser	Tyr	Phe	Ser	Leu	Leu	Arg	Phe	Leu	Leu
				155					160					165
Leu	Leu	Asn	Val	Leu	Ala	Ser	Val	Leu	Met	Ala	Cys	Met	Thr	Leu
				170					175					180
Leu	Pro	Thr	Trp	Leu	Gly	Gly	Ala	Pro	Pro	Gly	Pro	Pro	Gly	Pro
				185					190					195
Asp	Ile	Ser	Ser	Pro	Cys	Gly	Ser	Tyr	Asn	Pro	His	Ser	Gln	Gly
				200					205					210
Leu	Val	Thr	Phe	Ala	Thr	Gln	Leu	Phe	Asn	Leu	Leu	Ser	Gly	Glu
				215					220					225
Gly	Tyr	Leu	Glu	Trp	Ser	Pro	Leu	Phe	Tyr	Gly	Phe	Tyr	Pro	Pro
				230					235					240
Arg	Pro	Arg	Leu	Ala	Val	Thr	Tyr	Leu	Cys	Trp	Ala	Phe	Ala	Val
				245					250					255
Gly	Leu	Ile	Cys	Leu	Leu	Leu	Ile	Leu	His	Arg	Ser	Val	Ser	Gly
				260					265					270
Leu	Lys	Gln	Thr	Leu	Leu	Ala	Glu	Ser	Glu	Ala	Leu	Thr	Ser	Tyr
				275					280					285
Ser	His	Arg	Val	Phe	Ser	Ala	Trp	Asp	Phe	Gly	Leu	Cys	Gly	Asp
				290					295					300
Val	His	Val	Arg	Leu	Arg	Gln	Arg	Ile	Ile	Leu	Tyr	Glu	Leu	Lys
				305					310					315
Val	Glu	Leu	Glu	Glu	Thr	Val	Val	Arg	Arg	Gln	Ala	Ala	Val	Arg
				320					325					330
Thr	Leu	Gly	Gln	Gln	Ala	Arg	Val	Trp	Leu	Val	Arg	Val	Leu	Leu
				335					340					345
Asn	Leu	Leu	Val	Val	Ala	Leu	Leu	Gly	Ala	Ala	Phe	Tyr	Gly	Val
				350					355					360
Tyr	Trp	Ala	Thr	Gly	Cys	Thr	Val	Glu	Leu	Gln	Glu	Met	Pro	Leu
				365					370					375
Val	Gln	Glu	Leu	Pro	Leu	Leu	Lys	Leu	Gly	Val	Asn	Tyr	Leu	Pro
				380					385					390
Ser	Ile	Phe	Ile	Ala	Gly	Val	Asn	Phe	Val	Leu	Pro	Pro	Val	Phe
				395					400					405
Lys	Leu	Ile	Ala	Pro	Leu	Glu	Gly	Tyr	Thr	Arg	Ser	Arg	Gln	Ile
				410					415					420
Val	Phe	Ile	Leu	Leu	Arg	Thr	Val	Phe	Leu	Arg	Leu	Ala	Ser	Leu
				425					430					435
Val	Val	Leu	Leu	Phe	Ser	Leu	Trp	Asn	Gln	Ile	Thr	Cys	Gly	Gly
				440					445					450
Asp	Ser	Glu	Ala	Glu	Asp	Cys	Lys	Thr	Cys	Gly	Tyr	Asn	Tyr	Lys
				455					460					465
Gln	Leu	Pro	Cys	Trp	Glu	Thr	Val	Leu	Gly	Gln	Glu	Met	Tyr	Lys
				470					475					480
Leu	Leu	Leu	Phe	Asp	Leu	Leu	Thr	Val	Leu	Ala	Val	Ala	Leu	Leu
				485					490					495
Ile	Gln	Phe	Pro	Arg	Lys	Leu	Leu	Cys	Gly	Leu	Cys	Pro	Gly	Ala
				500					505					510
Leu	Gly	Arg	Leu	Ala	Gly	Thr	Gln	Glu	Phe	Gln	Val	Pro	Asp	Glu
				515					520					525
Val	Leu	Gly	Leu	Ile	Tyr	Ala	Gln	Thr	Val	Val	Trp	Val	Gly	Ser
				530					535					540
Phe	Phe	Cys	Pro	Leu	Leu	Pro	Leu	Leu	Asn	Thr	Val	Lys	Phe	Leu

Leu Leu Phe Tyr	545	Leu Lys Lys Leu Thr	550	Leu Phe Ser Thr Cys Ser	555
Pro Ala Ala Arg	560	Thr Phe Arg Ala Ser	565	Ala Ala Asn Phe Phe Phe	570
Pro Leu Val Leu	575	Leu Leu Gly Leu Ala	580	Ile Ser Ser Val Pro Leu	585
Leu Tyr Ser Ile	590	Phe Leu Ile Pro Pro	595	Ser Lys Leu Cys Gly Pro	600
Phe Arg Gly Gln	605	Ser Ser Ile Trp Ala	610	Gln Ile Pro Glu Ser Ile	615
Ser Ser Leu Pro	620	Glu Thr Thr Gln Asn	625	Phe Leu Phe Phe Leu Gly	630
Thr Gln Ala Phe	635	Ala Val Pro Leu Leu	640	Leu Ile Ser Ser Ile Leu	645
Met Ala Tyr Thr	650	Val Ala Leu Ala Asn	655	Ser Tyr Gly Arg Leu Ile	660
Ser Glu Leu Lys	665	Arg Gln Arg Gln Thr	670	Glu Ala Gln Asn Lys Val	675
Phe Leu Ala Arg	680	Arg Ala Val Ala Leu	685	Thr Ser Thr Lys Pro Ala	690
Leu	695		700		705

<210> 37

<211> 466

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6114480CD1

<400> 37

Met Ala Phe Val Leu	1	Ile Leu Val Leu Ser	10	Phe Tyr Glu Leu Val	15
Ser Gly Gln Trp Gln	5	Val Thr Gly Pro Gly	20	Lys Phe Val Gln Ala	25
Leu Val Gly Glu Asp	10	Ala Val Phe Ser Cys	25	Ser Leu Phe Pro Glu	30
Thr Ser Ala Glu Ala	15	Met Glu Val Arg Phe	30	Phe Arg Asn Gln Phe	35
His Ala Val Val His	20	Leu Tyr Arg Asp Gly	35	Glu Asp Trp Glu Ser	40
Lys Gln Met Pro Gln	25	Tyr Arg Gly Arg Thr	40	Glu Phe Val Lys Asp	45
Ser Ile Ala Gly Gly	30	Arg Val Ser Leu Arg	45	Leu Lys Asn Ile Thr	50
Pro Ser Asp Ile Gly	35	Leu Tyr Gly Cys Trp	50	Phe Ser Ser Gln Ile	55
Tyr Asp Glu Glu Ala	40	Thr Trp Glu Leu Arg	55	Val Ala Ala Leu Gly	60
Ser Leu Pro Leu Ile	45	Ser Ile Val Gly Tyr	60	Val Asp Gly Gly Ile	65
Gln Leu Leu Cys Leu	50	Ser Ser Gly Trp Phe	65	Pro Gln Pro Thr Ala	70
Lys Trp Lys Gly Pro	55	Gln Gly Gln Asp Leu	70	Ser Ser Asp Ser Arg	75
Ala Asn Ala Asp Gly	60	Tyr Ser Leu Tyr Asp	75	Val Glu Ile Ser Ile	80
Ile Val Gln Glu Asn	65	Ala Gly Ser Ile Leu	80	Cys Ser Ile His Leu	85
Ala Glu Gln Ser His	70	Glu Val Glu Ser Lys	85	Val Leu Ile Gly Glu	90
	75		90		95
	80		95		100
	85		100		105
	90		105		110
	95		110		115
	100		115		120
	105		120		125
	110		125		130
	115		130		135
	120		135		140
	125		140		145
	130		145		150
	135		150		155
	140		155		160
	145		160		165
	150		165		170
	155		170		175
	160		175		180
	165		180		185
	170		185		190
	175		190		195
	180		195		200
	185		200		205
	190		205		210
	195		210		215
	200		215		220
	205		220		225

Thr	Phe	Phe	Gln	Pro	Ser	Pro	Trp	Arg	Leu	Ala	Ser	Ile	Leu	Leu	
				230					235						240
Gly	Leu	Leu	Cys	Gly	Ala	Leu	Cys	Gly	Val	Val	Met	Gly	Met	Ile	
				245					250						255
Ile	Val	Phe	Phe	Lys	Ser	Lys	Gly	Lys	Ile	Gln	Ala	Glu	Leu	Asp	
				260					265						270
Trp	Arg	Arg	Lys	His	Gly	Gln	Ala	Glu	Leu	Arg	Asp	Ala	Arg	Lys	
				275					280						285
His	Ala	Val	Glu	Val	Thr	Leu	Asp	Pro	Glu	Thr	Ala	His	Pro	Lys	
				290					295						300
Leu	Cys	Val	Ser	Asp	Leu	Lys	Thr	Val	Thr	His	Arg	Lys	Ala	Pro	
				305					310						315
Gln	Glu	Val	Pro	His	Ser	Glu	Lys	Arg	Phe	Thr	Arg	Lys	Ser	Val	
				320					325						330
Val	Ala	Ser	Gln	Gly	Phe	Gln	Ala	Gly	Arg	His	Tyr	Trp	Glu	Val	
				335					340						345
Asp	Val	Gly	Gln	Asn	Val	Gly	Trp	Tyr	Val	Gly	Val	Cys	Arg	Asp	
				350					355						360
Asp	Val	Asp	Arg	Gly	Lys	Asn	Asn	Val	Thr	Leu	Ser	Pro	Asn	Asn	
				365					370						375
Gly	Tyr	Trp	Val	Leu	Arg	Leu	Thr	Thr	Glu	His	Leu	Tyr	Phe	Thr	
				380					385						390
Phe	Asn	Pro	His	Phe	Ile	Ser	Leu	Pro	Pro	Ser	Thr	Pro	Pro	Thr	
				395					400						405
Arg	Val	Gly	Val	Phe	Leu	Asp	Tyr	Glu	Gly	Gly	Thr	Ile	Ser	Phe	
				410					415						420
Phe	Asn	Thr	Asn	Asp	Gln	Ser	Leu	Ile	Tyr	Thr	Leu	Leu	Thr	Cys	
				425					430						435
Gln	Phe	Glu	Gly	Leu	Leu	Arg	Pro	Tyr	Ile	Gln	His	Ala	Met	Tyr	
				440					445						450
Asp	Glu	Glu	Lys	Gly	Thr	Pro	Ile	Phe	Ile	Cys	Pro	Val	Ser	Trp	
				455					460						465

Gly

<210> 38

<211> 2801

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 112301CB1

<400> 38

```

cgccctcccc gagcgagacc aaaacaggtg gaatccgggc tggagccgga gctccggcgg 60
cgcggttggc ggcacgtccc tccagacagt accacaggca cctggagtac cggcatcggt 120
cgctgtggcc cccgagtgtc cgtcagagcc taggggagcc tgccctcccg cgcctcgteg 180
gggcccggcc aggcaccttg gccgccggcg cacggacgcg ggcacgagca ctatgacacg 240
gctgctggac ctccggcacgt tgacaagatt tctctggggt accgcgaggg attactttga 300
atttcggttg tcgctgtgtg tctggcatat ttagaactta agtctattat ttcgggcacc 360
atgactttga ggcttttaga agactggtgc aggggggatgg acatgaaccc tcggaaagcg 420
ctattgattg ccggcatctc ccagagctgc agtgtggcag aaatcgagga ggctctgcag 480
gctggtttag ctcccttggg ggagtacaga ctgcttgga ggatgttcag gagggatgag 540
aacaggaaag tagccttagt agggcttact gcggagacta gtcacgccct ggtccctaag 600
gagataccgg gaaaaggggg tatctggaga gtgatcttta agccccctga cccagataat 660
acatttttaa gcagattaaa tgaattttta gcgggagagg gcatgacagt ggggtgagttg 720
agcagagctc ttggacatga aaatggctcc ttagaccag agcagggcat gatccggaa 780
atgtgggccc ctatgttggc acaggcatta gaggtctctc agcctgccct gcaatgcttg 840
aagtataaaa agctgagagt gttctcgggc agggagtctc cagaaccagg agaagaagaa 900
tttggacgct ggatgtttca tactactcag atgataaagg cgtggcaggt gccagatgta 960
gagaagagaa ggcgattgct agagagcctt cgaggcccag cacttgatgt tattcggtgc 1020
ctcaagataa acaatccttt aattactgtc gatgaatgtc tgcaggctct tgaggaggtg 1080
tttgggggta cagataatcc tagggagttg caggtcaa atctaaccac ttaccagaag 1140

```

```

gatgaggaaa agttgtcggc ttatgtacta aggctggagc ctttgttaca gaagctggta 1200
cagagaggag caattgagag agatgctgtg aatcaggccc gcctagacca agtcattgct 1260
ggggcagtc acaaaacaat tcgcagagag cttaatctgc cagaggatgg cccagcccct 1320
ggtttcttgc agttattggt actaataaag gattatgagg cagctgagga ggaggaggcc 1380
cttctccagg caatattgga aggtaatttc acctgagctc cagggaacca cgaagggata 1440
tggcaattgag tagagcatga aggtagaaca gtctatatac tcttgtgaca catacaatcc 1500
ctacctgtgt ctgccaaagta actcattttt gtgcaattct cagtataagc cctttgtcgt 1560
ttctgtgcct atttaaagtc tcctaaaggt gtaattgact aggaaggatg tagttctaca 1620
ctgccattta cctattttaa ttcattcctt tgaatatctt tgttgttgtt gttgagacag 1680
agtctcgtct tgtcaccagg gctggagtgc agtgacgtga tcttggctca ctgcatectc 1740
cgcttccag gttaaagcta ttctcctgcc tcagttgccc gagtagctgg gactacagga 1800
atgactcacc acaccagct aagtttttgc ttatcagtag agacgggggt tctccatact 1860
ggtcaggctg gtctcgaact cctgacctca agtgatccac ctgccctggc ctcccaaagt 1920
gctgggatta caggcgtgag ccaccacacc cagcctcagt tcccttattt taataattgt 1980
cctatttctg ggggttgcta taaagattac atgagacttt aatgggaaag tatttaggac 2040
agtgtatggg acaaagttag taccagacat gcataaggca tttgaaaacc tgggtgcttt 2100
ctctctcact cttaccattt atagacaggg gtcatataat tttttttgtt gagacggagt 2160
ctcgtctctg ggcccaggct ggagagcagt ggtgtgatct cagctcatgg gttcaagcga 2220
ttctcgtgcc tcagctctcc gagtagctgg gactataggc atgcgccacc agggccagcc 2280
aattttttata ttttttagtag agatgggggt tctccatggt agccaggcta cccaggctgg 2340
tctcgatctc ctgacctcag gtgatccacc cacctggggc tcccaaagtg ctgggattac 2400
aggcgtgagc cactgcccc tgccttctta cactttttct ttgagactgg gtcgcacagt 2460
cgcacagtga gactccatct cctatgaaaa aaaagattgg cctagcgtct ggctcaggcc 2520
tgtaatccca gcactttggg agggcaaggc gggcggatca caaggtcagg agatcgaggc 2580
cagcatggcc aacatgggtg aaccctggtt ctactaaaaa taaaaaaaaa attagccggg 2640
catgggtggc catgcctata gtgtcagcta ctctggagaa tcgcttgaac ccgggaggcg 2700
gaggttgcag tgagccgaga ttgcgtcact gcactacagc ctgggcaaca gagcgagact 2760
cccgtctcag aaaaaaaaaa ttttggtgcc aaccggacga g 2801

```

<210> 39

<211> 2656

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 997947CB1

<220>

<221> unsure

<222> 2554, 2587, 2604, 2606, 2611, 2644, 2646-2647, 2651

<223> a, t, c, g, or other

<400> 39

```

ccactctcta gagctccccg actcctagct ctgagaatgc ctgcggaatg atcgcccccc 60
aggccggctg ccgccgctgc cgctgctgct gttattgcta ctgctgctgc cgccgcctct 120
gcttccactc ggctctgact ggcaggcaga aagtgcacct gacgagggaa aggtctctgc 180
agtgagtgga gagcctacat aaaagagagt aaagaggggc aaaaaccagg atcagaatgc 240
aggcgacgtc caaccttctc aacctcctgc tgcgtctctt gtttgccgga ttagatcctt 300
ccaagactca gattagtctt aaagaagggt ggcagggtga cagctcagct caggatcctg 360
atgggcgggt catttgca ca gttgttgctc cagaacaaaa cctgtgttcc cgggatgcc 420
aaagcaggca acttcgccaa ctactggaaa aggttcagaa catgtcccag tctattgaag 480
tcttaaactt gagaactcag agagatttcc aatatgtttt aaaaatggaa acccaaatga 540
aagggtgaa ggcaaaattt cggcagattg aagatgatcg aaagacactt atgaccaagc 600
attttcagga gttgaaagag aaaatggacg agctcctgcc tttgatcccc gtgctggaac 660
agtacaaaac agatgctaag ttaatcacc agttcaagga ggaaataagg aatctgtctg 720
ctgtcctcac tggatttcag gaggaaattg gtgcctatga ctacgaggaa ctacaccaa 780
gagtgtgtag cttggaaaca agacttcgtg actgcatgaa aaagctaaca tgtggcaaac 840
tgatgaaaat cacaggccca gttacagtca agacatctgg aacccgattt ggtgcttgg 900
tgacagaccc ttttagcatc gagaaaaaca acagagtctg gtacatggac agttatacta 960
acaataaaa tgttcgtgaa tacaatacaa ttgcagactt tgtcagtggt gctgaatcaa 1020
ggacatacaa ccttccttcc aagtgggcag gaactaacca tgttgtctac aatggctcac 1080
tctattttta caagtatcag agtaatatca tcatcaaata cagctttgat atggggagag 1140

```

tgcttgccca	acgaagcctg	gagtatgctg	gttttcataa	tgtttaccce	tacacatggg	1200
gtggattctc	tgacatcgac	ctaattggctg	atgaaatcgg	gctgtgggct	gtgtatgcaa	1260
ctaaccagaa	tgagggaat	attgtcatca	gccaacttaa	ccaagatacc	ttggagggtga	1320
tgaagagctg	gagcactggc	taccccaaga	gaagtgcagg	ggaatctttc	atgatctgtg	1380
ggacactgta	tgccaccaac	tcccacttaa	ctggagccaa	ggtgtattat	tcctattcca	1440
ccaaaacctc	cacatatgag	tacacagaca	tccccttcca	taaccaatac	tttcacatat	1500
ccatgcttga	ctacaatgca	agagatcgag	ctctctatgc	ctggaacaat	ggccaccagg	1560
tgctgttcaa	tgccaccctt	ttccatatca	tcaagacaga	ggatgacaca	taggcaaattg	1620
tgacatgttt	tcattgattt	aaacagtgtg	atttgtgata	aactctataa	gacccttcc	1680
gtttttttct	tcactattat	ttttcatcat	ttctccaaag	caaagcattt	ttattgtaaa	1740
gttgggtgtt	caaaaacata	gctgagcttg	tctaacttac	catgttggaa	acacatctta	1800
acttctaaat	ttacaaggcc	tatcatgtcc	ttgtcatgaa	aagcactaaa	aaaaaaaaaga	1860
gtttaagtgg	ctaaagtcac	agttttgcaa	gagattaatg	atctgcctta	tattagagtc	1920
agagactaat	ggtggcttaa	atgcacgaat	gtcttttttt	tttaaaactg	tcattttttta	1980
ctgtcttttg	ctccatatca	ggaaatatct	tggtaggaa	taggagaaca	aaaagcactt	2040
ttatccccatt	tatttcttta	aaaaatgtaa	ggatttcatt	tataattgaaa	aataatatta	2100
atcattttgc	tgtaaacaca	attctctgat	gcggtgctgt	acagtcattt	ttaaatctct	2160
tgctaacatt	ttattggcag	tatgtatttc	taccattgta	accaccattg	tgctattgta	2220
tctcttcact	tctgtgaaag	taatattttt	tataaaatac	actgaaattt	aacctcagta	2280
attgagtcca	ttttcaagtg	tggtcaagaa	taatcttctt	ggcttaccce	ttacataag	2340
cattataaac	taaaatgaaa	accaaaccag	acacctgaca	tagagtcttt	attttaccce	2400
aagttttttg	ggccactgac	attgaatgca	acaactgatt	tcatacaact	gagttactct	2460
gttcaactcca	ctgaatgcaa	cccatatagt	ttcttgacac	aggtgcatct	ggattctaaa	2520
tattggattt	gagttgactc	tactcattta	ttanacacta	aagaaatttt	gttcttcata	2580
gttaaangta	ctagcattta	attnanattt	nacatacaac	ttgcaataat	gaaatttcctt	2640
atgncnngac	nctgaa					2656

<210> 40

<211> 968

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1521513CB1

<400> 40

tctacatcta	tcgggagctga	acttctctaaa	agacaaagtg	tttatctttc	aagatttcatt	60
ctccctgaat	cttaccaaca	aaacactcct	gaggagaaag	aaagagaggg	agggagagaa	120
aaagagagag	agagaaacaa	aaaaccaaag	agagagaaaa	aatgaattca	tctaaatcat	180
ctgaaacaca	atgcacagag	agaggatgct	tctcttccca	aatgttctta	tggactgttg	240
ctgggatccc	catcctattt	ctcagtgcc	gtttcatcac	cagatgtgtt	gtgacatttc	300
gcatctttta	aacctgtgat	gagaaaaagt	ttcagctacc	tgagaatttc	acagagctct	360
cctgctacaa	ttatggatca	ggttcagtc	agaattgttg	tccattgaac	tgggaatatt	420
ttcaatccag	ctgctacttc	ttttctactg	acaccatttc	ctgggcgtta	agtttaaaga	480
actgctcagc	catgggggct	cacctgggtg	ttatcaactc	acaggaggag	caggaattcc	540
tttctacaa	gaaacctaaa	atgagagagt	tttttattgg	actgtcagac	caggttgctg	600
agggctcagt	gcaatgggtg	gacggcacac	ctttgacaaa	gtctctgagc	ttctgggatg	660
tagggggagcc	caacaacata	gctaccctgg	aggactgtgc	caccatgaga	gactcttcaa	720
acccaaggca	aaattggaat	gatgtaacct	gtttctctca	ttattttcgg	atttgtgaaa	780
tggttaggaat	aaatcctttg	aacaaaggaa	aatctcttta	agaacagaag	gcacaactca	840
aatgtgtaaa	gaaggaagag	caagaacatg	gccacacca	ccgccccaca	cgagaaattt	900
gtgcgctgaa	cttcaaagga	cttcataagt	atttgttact	ctgatataaa	taaaaataag	960
tagtttta						968

<210> 41

<211> 1837

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1863994CB1

<400> 41

```

ccccagcccc gctcagtecc gagcgcccgcg agtcgtcgcg ccgcccgcgcc aagcatgcag 60
taaaggctga aaatctgggt cacagctgag gaagacctca gacatggagt ccaggatgtg 120
gcctgcgctg ctgctgtccc acctcctccc tctctggcca ctgctgttgc tgccccctccc 180
accgcctgct cagggctctt catcctcccc tcgaacccca ccagccccag cccgccccccc 240
gtgtgccagg ggaggccctt cggccccacg tcatgtgtgc gtgtggggagc gagcacctcc 300
accaagccga tctcctcggg tcccaagatc acgtcggcaa gtcctgcctg gcactgcacc 360
ccagccacc ccatcaggct ttgaggagg gcccctcca tcccaatacc cctgggctat 420
cgtgtggggt cccaccgtgt ctcgagagga tggaggggac cccaactctg ccaatcccgg 480
atttctggac tatggttttg cagccccctca tgggctcgca accccacacc ccaactcaga 540
ctccatgcga ggtgatggag atgggcttat ccttgagagag gcacctgcca ccctgcggcc 600
attcctgttc gggggccgtg ggggaagggt ggacccccag ctctatgtca caattaccat 660
ctccatcatc attgttctcg tggccactgg catcatcttc aagtctctgt gggaccgcag 720
ccagaagcga cgcagacct cagggcagca aggtgccttg aggcaggagg agagccagca 780
ggcactgaca gacctgtccc cggtggagt cactgtgctg ggggccttcg gggactcacc 840
tacccccacc cctgaccatg agggaccccg agggggaccc cggcctggga tgccccaccc 900
caaggggggt ccagccttcc agttgaaccg gtgagggcag gggcaatggg atgggagggc 960
aaagagggaa ggcaacttag gtcttcagag ctgggggtgg ggtgccctct ggatgggtag 1020
tgaggaggga ggcgtggcct cccacagccc ctggccctcc caaggggggt ggaccagctc 1080
ctctctggga ggcacccttc ctctctccag tctctcagga tctgtgtcct attctctgct 1140
gcccataact ccaactctgc cctctttggt tttttctcat gccaccttgt ctaagacaac 1200
tctgccctct taaccttgat tccccctctt tgtcttgaa tcccccttct attctggcct 1260
accccttggt tctgactgt gccctttccc tcttctctc aggattcccc tggatgaatct 1320
gtgatgcccc caatgttggg gtgcagccaa gcaggaggcc aaggggcccg cacagccccc 1380
atcccactga ggggtggggca gctgtgggga gctggggcca caggggctcc tggctcctgc 1440
cccttgacac ccaccggaa cactccccag cccacggggc aatcctatct gctgcacctc 1500
ctgcaggtag gggcctcaca tatctgtgac tctgggtccc tgtccccacc cttgtgcact 1560
cacatgaaag ccttgacac tcacctccac ctacacaggc catttgacac cgctcctgca 1620
ccctctcccc gtccataccg ctccgctcag ctgactctca tgttctctcg tctcacattt 1680
gcactctctc cttcccatat tctgtgctca gctcactcag tggtcagcgt ttcctgcaca 1740
ctttacctct catgtgcgtt tcccggcctg atgttgtggt ggtgtgcggc gtgtcactc 1800
tctccctcat gaacaccac ccacctcgtt tcgcagc 1837

```

<210> 42

<211> 2124

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2071941CB1

<400> 42

```

gtgacgcggc tgcggagggt acgcgggagg tcgcgcgccc cttccggcgc ggggagggcg 60
ctgaagatcg gggccgctcg gccgcaggcg cctccagcgc cgcgggatgt agcgcggggg 120
accgcggccc ccagcagagc ccgcctgccc ggcttgtcta ccatcagagg gagatctctg 180
ccccctgggg ctgagagacc ccaaccttcc cccaagctga agctgcaggg tattgaggta 240
ccagccagat gtcttccac aaaggatctg tgggtggcaca ggggaatggg gctcctgcca 300
gtaacaggga agctgacacg gtggaactgg ctgaactggg acccctgcta gaagagaagg 360
gcaaacgggt aatcgccaac ccacccaaag ctgaagaaga gcaaaccatg ccagtgcacc 420
aggaagaaga ggaggagggt cgggtactga cacttcccc gcaagcccac cagcccatgg 480
agaagatgga agagtttgtg tacaaggctt gggagggacg ttggagggtc atcccatatg 540
atgtgtctcc tgaactggta aaggacaacg actatctgct acatgggtcat agacctcca 600
tgccctcctt tcgggcttgc ttcaagagca tcttccgcat tcatacagaa actggcaaca 660
tctggaccac tctgcttggg ttcgtgctgt ttctctttt gggaatcttg accatgctca 720
gaccaaatat gtacttcatg gcccctctac aggagaaggg ggtttttggg atgttctttt 780
tgggtgcagt agcttgcctc agcttctcct ggctcttcca caccgtctat tgtcattcag 840
agaaagtctc tcggactttt tccaaactgg actattcagg gattgctctt ctaattatgg 900
ggagctttgt cccctggctc tattatctct tctactgtc cccacagcca cggctcatct 960
acctctccat cgtctgtgtc ctgggcattt ctgccatcat tgtggcgag tgggaccggt 1020
ttgccactcc taagcaccgg cagacaagag caggcggtgt cctgggactt ggcttgagt 1080
gcgtcgtgcc caccatgcac tttactatcg ctgagggtt tgtcaaggcc accacagtgg 1140
gccagatggg ctggttcttc ctcatggctg tgatgtacat cactggagct ggcctttatg 1200

```

```

ctgctcgaat tcttgagcgc ttcttttctg gaaaatttga catatgggtc cagtctcatc 1260
agattttcca tgtcctggtg gtggcagcag cctttgtcca cttctatgga gtctccaacc 1320
ttcaggaatt ccgttacggc ctagaaggcg gctgtactga tgacaccctt ctctgagcct 1380
tcccacctgc ggggtggagg aggaacttcc caagtgtttt taaaaataac ttctttgctg 1440
aagtgagagg aagagtctga gttgtctgtt tctagaagaa acctcttaga gaattcagta 1500
ccaaccaagc ttcagcccac ttccacaccc actgggcaat aaactttcca ttccattctt 1560
cctagctggg gatggggcat ggtcaaaact agccatcccc tcctcagcaa ggcattctacc 1620
ggccccctac agagacagta ctttgaaact catgttgaga ttttaccctc tcctccaacc 1680
attttgggaa aattatggac tgggactctt cagaaattct gtcttttctt ctggaagaaa 1740
atgtccctcc cttaccccca tccttaactt tgtatcctgg cttataacag gccattcatt 1800
tttgtagcac acttttcaaa aacaattata taccctgggt ccattcttct agggcctgga 1860
tctgcttata gagcaggaag aataaagcca ccaactttta cctagcccgg ctaatcatgg 1920
aagtgtgtcc aggcttcaag taacttgagt ttttaatttt ttttttttct tggcagagta 1980
atgtaaaatt taaatgggga aagatattta atatttaata ctaagcttta aaaagaaacc 2040
tgctatcatt gctatgtatc ttgatgcaaa gactatgatg ttaataaaaag aaagtacaga 2100
agagacttgg cattcaaaaa aaaa 2124

```

<210> 43

<211> 993

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2172512CB1

<400> 43

```

ccgcgcgctt actttgttta taacttgaaa aatcctctcc gtctcccttc cctgcctcct 60
ttcctttccc tttcctctgc cagtacaact agaccggcg tctggcgctc ccggtgcccc 120
gcattctgcg gggcaggcgg attaattgga attcttcaaa atgtcagggt tggtagccac 180
agcccttgaa cagcctgcag gtgaaatgga aaatcaaaca aaaccaccag atccaaggcc 240
tgatgtctct cctgaatata gttctcattt tttaccagga ccccttgaa cagctgtccc 300
tccacctact ggctaccag gaggttgcc tatgggatac tacagtccac agcaaccag 360
taccttccct ttgtaccagc cagttgggtg tatccatcct gtccggtatc agcctggcaa 420
atatactatg ccaaatcagt ctgttccaat aacatggatg ccaggggcaa ctcctatggc 480
aaactgcctt cctggtctgg aatacttagt tcagttggac aacatacatg ttcttcagca 540
ttttgagcct ctggaaaatga tgacatgttt tgaaactaat aatagatatg atattaaaaa 600
caactcagac cagatggttt acattgtaac cgaagacaca gatgacttta ccaggaatgc 660
ctatcgggaca ctaaggccct tcgtcctccg ggtcactgat tgtatgggcc gagaaatcat 720
gacaatgcag agacccttca gatgcacctg ctgttgcttc tgttgccctt ctgccagaca 780
agagctggag gtgcagtgtc ctctgggtgt caccattggc tttgttgagg aacattggaa 840
cctgtgcagg gcggtgtaca gcatccaaaa aaaaaaaaaa aaaattgcgg cgcaagctta 900
ttcccttttag tgaggttaat ttttgggggc atgtgccgtg gcagtgcggg ggtgttggtg 960
gttgtgtggg ggtgggggtg ggggttggtg ggt

```

<210> 44

<211> 2214

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2483172CB1

<400> 44

```

ggctctcggc ccagcgcgcc tgccttcgcc gcccgccgtc gctcctcgcc cgctgcacga 60
cgacgcgagc cccctgtctg aggcggcgga cccgaccgga ccagaccga gacgcaagat 120
ggcgacggcc gcgtgactgc ctcagcgtcc ccgagctcgg ctccgagtgc acctacggac 180
tgactgtggg ggcagagaag ggcgagatca ggactctgtc tttgttaatc gtgactgcat 240
gaaggctgcc tccctcgggc ctacttggtg ggagtgtctg gtattgttct aaggccagga 300
gcacggtgag ccacagtctg ttggtagaat ttggcgtctt gatagttgag aaaatggcga 360
tgacactgtt ggaagactgg tgccggggga tggatgtgaa ctcccagaga gctctgttag 420
tctggggcat cccagtgaac tgtgatgagg ctgaaatcga agagaccctc caggctgcga 480

```

```

tgccccaggt ctcctaccga atgcttggga gaatgttctg gagggaagaa aatgcgaaag 540
cagccttatt agagctcact ggcgtgttag attacgccgc gatccccagg gagatgccgg 600
gcaaaggagg ggtctggaaa gtgttattta agcccccaac ttctgatgct gaatttttag 660
aaagattgca cctcttccta gctagagagg ggtggaccgt gcaagatgtt gcccgtgtcc 720
ttgggtttca gaaccctact ccgaccccg gcccagagat gccagcagag atgctaaact 780
atattttgga taatgttatt cagcctcttg ttgagtccat atggtacaag aggctgacac 840
ttttctcggg gagggacatc ccagggcctg gagaggaaac ctttgatccc tggctggagc 900
acactaatga ggtcctagag gagtggcagg tgtccgatgt agaaaagagg cggcggttga 960
tggagagtct tagaggcccc gccgctgatg ttattcgcac ccttaagtcc aacaaccccg 1020
cgataaccac tgccgaatgc ctgaaggcgc ttgagcaggt gtttgggagc gttgagagct 1080
ctaggggatgc ccagatcaaa tttctgaaca cttatcagaa cccgggagaa aaattgtctg 1140
cttatgtcat tcgtctggag cctctgctac agaaggtggt agagaagggg gccattgata 1200
aagataatgt gaaccaggcc cgcctagagg aggtcattgc cggggccaac cacagcgggg 1260
ccatccgaag gcagctgtgg cttaccgggg ctgggggaagg gccagcccca aacctcttcc 1320
agttgctggt gcagatccgt gaggaggaag ccaaggagga ggaggaggag gctgaggcca 1380
cccttctgca gttaggcctg gaagggcact tctgagtgcc aggaaaggca gcttttagtg 1440
agacctagat cacagctact tttcttgctc ctgtgggggtc ttacagatgt gtctctgagt 1500
agtaaaggct tagccttggt ctgttttgtt gttttttgga ggggaagggt agtcaggcct 1560
gagtattcat gtaacattct aaaattgtgc cagcgagcac cgtgaacgac tgcaatgcaa 1620
gcgggtcttg ctggctaagg taaagggttg gttggacaca gcgcttagtg cacgctgtca 1680
tcatggacat cataatcagt tgtgaaaaac acgcgaacct atgacacttc ttattccaca 1740
ctgaatgtga aattgcatgt tcagatgttt actacgaggc ctggctcaca ggaagtgttc 1800
agtaaaagta tgcactgtta gattactgat aacgcggata gatttttgtt taccataaat 1860
tggtccagat ttatattaat ggaaggaaat gtgcatttat tagctattac tcaactttac 1920
aatgcaaaca tcttatttct catctttaaa catgtcgacc agtttaattg aaaagtattc 1980
tgagactgca aaatgggggtg ttaaaaata ctgcagttac ggagctgtgt aaaccagttt 2040
ctcattgcat aagatacaga tgtaaattgc atggagaggt tgatatgcac ctgtacagta 2100
attcactccc ccatttcaca tctttgtcag agaatagttc ttgttcatac tgagtgttct 2160
aaatttgaag ttatatatac aaattaaaa atttttaaaaa ttcaaaaaaa aaaa 2214

```

<210> 45

<211> 897

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2656128CB1

<400> 45

```

gcaaaatgca tgacagtaac aatgtggaga aagacattac accatctgaa ttgcctgcaa 60
acccagggtg tctgcattca aaagagcatt ctattaaagc taccttaatt tggcgcttat 120
ttttcttaat catgtttctg acaatcatag tgtgtggaat gggtgctgct ttaagcgcaa 180
taagagctaa ctgccatcaa gagccatcag tatgtcttca agctgcatgc ccagaaagct 240
ggattgggtt tcaaagaaag tgtttctatt tttctgatga caccaagaac tggacatcaa 300
gtcagagggt ttgtgactca caagatgctg atcttgctca gggtgaaagc ttccaggaa 360
tgaatttcct gttgagatat aaaggcccat ctgatcactg gattgggctg agcagagaac 420
aaggccaacc atggaaatgg ataaatggta ctgaatggac aagacagtta gtcatgaaag 480
aagatggtgc caacttgtat gttgcaaagg tttcacaagt toctcgaatg aatccaagac 540
ctgtcatggt ttctatcctt gggagcagga gagtgtgcct atttgaatga caaagggtgc 600
agtagtgcca ggactacac agagaggaag tggatttggt ccaaatacaga tatacatgtc 660
tagatgttac agcaaagccc caactaatct ttagaagcat attggaactg ataactccat 720
tttaaaatga gcaaagaatt tattttctat accaacaggt atatgaaaat atgctcaata 780
tcactaataa ctgggaaaat acaaatacaa atcatagtaa aatattacct gttttcatgg 840
tgctaataat acctgttctc ccactgctaa tgacataccc gagactgagt aatttaa 897

```

<210> 46

<211> 2167

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5855841CB1

<400> 46

```
gccggggccg gagttcgctg caagtcggcg gaaagttttg ctgcgcgggg tccccgaag 60
ttcagagtga agacatttcc acctggacac ctgaccatgt gcctgccctg agcagcgagg 120
cccaccaggc atctctgttg tgggcagcag ggccagggtcc tggctctgtg accctcgga 180
gttggcaggc tccctctgca gtggggctct ggccctcgcc ccaccatgtc gagcctcggc 240
ggtggctccc aggatgccgg cggcagtagc agcagcagca ccaatggcag cgggtggcag 300
ggcagcagtg gcccaaaggc aggagcagca gacaagagtg cagtgggtgg tgcggccgca 360
ccagcctcag tggcagatga cacaccaccc cccgagcgtc ggaacaagag cggtatcatc 420
agttagcccc tcaacaagag cctgcgcgcg tcccgcgcgc tctcccacta ctcttctttt 480
ggcagcagtg gtggtagtgg cgggtggcagc atgatgggag gagagtctgc tgacaaggcc 540
actgcggctg cagccgctgc ctccctgttg gccaatgggc atgacctggc ggcgccatg 600
gcggtggaca aaagcaaccg agcggggcac ggagctggca gccgagggac agctgacgct gcagcagttt 720
agcaaggcag agcggggcac gaagcgcgtg gtgcaggagc atctcccgtc gatgagcgag 780
gcgcagtgca cagagatgct catggaggct gtggcagggt ccgaagccct caatggccag 840
gcgggtgctg gcctgcctga catggaggct gtggcagggt ccgaagccct caatggccag 900
tccgacttcc cctacctggg cgctttcccc atcaaccagg gcctcttcat tatgaccccg 960
gcaggtgtgt tcttgccgga gagcgcgctg cacatggcgg gcctggctga gtaccccatg 1020
cagggagagc tggcctctgc catcagctcc ggcaagaaga agcgggaaacg ctgcgccatg 1080
tgcgcgccct gccggcgccg catcaactgc gagcagtgca gcagttgtag gaatcgaaag 1140
actggccatc agattttgcaa attcagaaaa tgtgaggaac tcaaaaagaa gccttccgct 1200
gctctggaga aggtgatgct tccgacggga gccgccttcc ggtgggtttca gtgacggcgg 1260
cggaacccaa agctgccctc tccgtgcaat gtcactgctc gtgtgggtctc cagcaaggga 1320
tccggcgcaa gacaaacgga tgcacccgctc tttagaacca aaaatattct ctcacagatt 1380
tcattctgtg ttttatatat atattttttg ttgtcgtttt aacatctcca cgtccctagc 1440
ataaaaagaa aaagaaaaaa atttaaaactg ctttttcgga agaacaacaa caaaaaagag 1500
gtaaagacga atctataaag taccgagact tcctgggcaa agaattggaca atcagtttcc 1560
ttcctgtgtc gatgtcgatg ttgtctgtgc aggagatgca gtttttgtgt agagaatgta 1620
aattttctgt aaccttttga aatctagtta ctaataagca ctactgtaat ttagcacagt 1680
ttaactccac cctcatttaa acttcctttg agaatgttg atggcgccgt gtagaagccg 1740
ttgcctggag aatccactca cgttcataaa gagaatgttg cagcagggag ctcacagaag 1800
ctctgtatcc atccacgctg gcagagctgc cgagactggg atccccacc ccaacagtga 1860
accaggccag ctgagctgca cccacagtcg tttatccatt gcgatctggg gagccccatc 1920
ttttggaaaa aaaaatgaaa gttctgttcg tttatccatt agagaaaata agtccttttt 1980
tcgatatttc caatcctggc tactttttct agagaaaata agtccttttt ttctggcctt 2040
gctaattggc acagaagaaa gggcttcttt cgctgggtcc ctgctgggtg ggggtgggtcc 2100
ccaggggggc cctgcggcc tgggcccccc cagcttctct ctgatgaaca 2160
tgctgtttgt attgttttag gaaaccaggc tgttttgtga ataaaacgaa tgcagtgttt 2160
tgtcacg
```

<210> 47

<211> 1235

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 603462CB1

<400> 47

```
gtggagccgg cgagagagtg gcagcggggg ctgatggaag tgcagtgggg gctggagagg 60
gcaccctact gtatccagca tgctccaagg ccacagctct gtgttccagg ccttgctggg 120
gaccttcttc acctggggga tgacagcagc tggggcagct ctcgtgttcg tattctctag 180
tggacagagg cggatcttag atggaagtct tggctttgct gcaggggtca tgttggcagc 240
ttcctattgg tctcttctgg cccagcagtg tgagatggcc acgtcctctg ggggcttcgg 300
tgcccttggc ttcttccctg tggctgttgg cttcacccct ggagcggctt ttgtctactt 360
ggctgacctc ctgatgcctc acttgggtgc agcagaagac cccagacgg ccctggcact 420
gaacttcggc tctacgttga tgaagaagaa gtctgacctc gaggggtccc cgctgctctt 480
ccctgagagt gaactttcca tccggataga caagagttag aatgggtgagg catatcagag 540
aaagaaggcg gcagccactg gccttccaga gggctctgct gtcctctgtc cttctcgagg 600
gaatctggca cagcccgccg gcagcagctg gagggagatc gcaactgctc tcttggccat 660
cactatacac aacgttccag agggctctgc tggttggagt ggatttgggg ctatagaaaa 720
```

gacggcatct	gctacctttg	agagtgccag	gaatttggcc	attggaatcg	ggatccagaa	780
tttccccgag	ggcctggctg	tcagccttcc	cttgcgaggg	gcaggcttct	ccacctggag	840
agctttctgg	tatgggcagc	tgagcggcat	ggtggagccc	ctggccgggg	tctttgggtg	900
ctttgccgtg	gtgctggctg	agcccatact	gccctacgct	ctggcctttg	ctgccggtgc	960
catggtctac	gtgggtcatg	acgacatcat	ccccgaagcc	cagatcagtg	gtaatgggaa	1020
actggcatcc	tgggcctcca	tcctgggatt	tgtagtgatg	atgtcactgg	acgttggcct	1080
gggctagggc	tgagacgctt	cggaccctgg	gaaaggccat	acgaagaaac	agcagtgggt	1140
ggcttctatg	ggacaacaag	cttctttctt	cacattaaaa	cttttttctt	tcctctcttc	1200
ttcatctcat	tatcctgatt	gactctaatt	ataat			1235

<210> 48

<211> 2257

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 747681CB1

<400> 48

ccgagccgag	cggaccgaag	gcgcgcccga	gatgcaggtg	agcaagagga	tgctggcggg	60
gggctgagg	agcatgcca	gccccctcct	ggcctgctgg	cagcccatcc	tcctgctggt	120
gctgggctca	gtgctgtcag	gctcggccac	gggctgcccg	ccccgctgcg	agtgtctccg	180
ccaggaccgc	gctgtgctgt	gccaccgcaa	gcgctttgtg	gcagtccccg	agggcatccc	240
caccgagacg	cgctgtctgg	acctaggcaa	gaaccgcatc	aaaacgctca	accaggacga	300
gttcgcccag	ttcccgcacc	tggaggagct	ggagctcaac	gagaacatcg	tgagcgccgt	360
ggagcccggc	gccttcaaca	acctcttcaa	cctccggacg	ctgggtctcc	gcagcaaccg	420
cctgaagctc	atcccgttag	gcgtcttcac	tggcctcagc	aacctgacca	agctggacat	480
cagcgagaac	aagattgtta	tcctgctgga	ctacatgttt	caggacctgt	acaacctcaa	540
gtcactggag	gttggcgaca	atgacctcgt	ctacatctct	caccgcgcct	tcagcggcct	600
caacagcctg	gagcagctga	cgctggagaa	atgcaacctg	acctccatcc	ccaccgaggc	660
gctgtcccac	ctgcacggcc	tcctcgtcct	gaggctccgg	cacctcaaca	tcaatgccat	720
ccgggactac	tccttcaaga	ggctctaccg	actcaaggct	ttggagatct	ccactggcc	780
ctacttggac	accatgacac	ccaactgcct	ctacggcctc	aacctgacgt	ccctgtccat	840
cacacactgc	aatctgaccg	ctgtgcccct	cctggccgtc	cgccacctag	tctatctccg	900
cttcctcaac	ctctcctaca	acccccatcag	caccattgag	ggctccatgt	tgcattgagct	960
gctccggctg	caggagatcc	agctgggtgg	cgggcagctg	gccgtgggtg	agccctatgc	1020
cttcgcgggc	ctcaactacc	tgcgcgtgct	caatgtctct	ggcaaccagc	tgaccacact	1080
ggaggaatca	gtcttccact	cggtgggcaa	cctggagaca	ctcatcctgg	actccaaccc	1140
gctggcctgc	gactgtcggc	tcctgtgggt	gttcgcggcg	cgctggcggc	tcaacttcaa	1200
ccggcagcag	cccacgtgcg	ccacgcccga	gtttgtccag	ggcaaggagt	tcaaggactt	1260
ccctgatgtg	ctactgcccc	actacttcac	ctgccgccgc	gcccgcctcc	gggaccgcaa	1320
ggcccagcag	gtgtttgtgg	acgagggccca	cacggtgcag	tttgtgtgcc	gggcccgtgg	1380
cgaccgcggc	cccgccatcc	tctggctctc	accccgaag	cacctggtct	cagccaagag	1440
caatgggcgg	ctcacagtct	tcctgatgg	cacgctggag	gtgcgctacg	cccagggtaca	1500
ggacaacggc	acgtacctgt	gcctgcgggc	caacgcgggc	ggcaacgact	ccatgcccgc	1560
ccacctgcat	gtgcgcagct	actcgcccga	ctggccccat	cagcccaaca	agaccttcgc	1620
tttcatctcc	aaccagccgg	gcgagggaga	ggccaacagc	acccgcgcca	ctgtgccttt	1680
ccccttcgag	atcaagacc	tcctcatcgc	caccacctgt	ggcttcatct	ctttcctggg	1740
cgctgtcctc	ttctgcttgg	tgctgctgtt	tctctggagc	cggggcaagg	gcaacacaaa	1800
gcacaacatc	gagatcgagt	atgtgccccg	aaagtcggac	gcaggcatca	gctccgccga	1860
cgcgccccgc	aagttcaaca	tgaagatgat	atgaggccgg	ggcggggggc	agggaccccc	1920
gggcgggccg	gcaggggaag	gggcctggcc	gccacctgct	cactctccag	tccttcccac	1980
ctcctcccta	cccttctaca	cacgttctct	ttctccctcc	cgctccgctc	ccctgctgcc	2040
ccccgccagc	cctcaccacc	tgcctcctct	ctaccaggac	ctcagaagcc	cagacctggg	2100
gacccacact	acacaggggc	attgacagac	tggagttgaa	agccgacgaa	ccgacacgcg	2160
gcagagtcaa	taattcaata	aaaaagttac	gaactttctc	tgtaacttgg	gtttcaataa	2220
ttatggattt	ttatgaaaac	ttgaaataaa	aaaaaaa			2257

<210> 49

<211> 2359

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 919469CB1

<400> 49

```
gtgggatttg aagatccact ccacttctgc tcatggcggg ccagggcctg cccctgcacg 60
tggccacact gctgactggg ctgctggaat gcctgggctt tgctggcgct ctctttggct 120
ggccttcact agtgtttgtc ttcaagaatg aagattactt taaggatctg tgtggaccag 180
atgctggggc gattggcaat gccacagggc aggctgactg caaagcccag gatgagagg 240
tctcactcat cttcacccctg gggtccttca tgaacaactt catgacattc cccactggct 300
acatctttga cgggttcaag accaccgtgg cagcctcat agccatattt ttctacacca 360
ccgccacact catcatagcc ttcacctctg caggctcagc cgtgctgctc ttctggcca 420
tgccaatgct caccattggg ggaatcctgt ttctcatcac caacctgcag attgggaacc 480
tatttggcca acaccgttcg accatcatca ctctgtacaa tggagcattt gactcttctt 540
cggcagtctt ccttattatt aagcttcttt atgaaaaagg catcagcctc agggcctcct 600
tcatcttcat ctctgtctgc agtacctggc atgtagcacg cactttcttc ctgatgcccc 660
ggggggcacat cccataccca ctgcccccca actacageta tggcctgtgc cctgggaatg 720
gcaccacaaa ggaagagaag gaaacagctg agcatgaaaa cagggagcta cagtcaagg 780
agttcctttc agcgaaggaa gagacccag gggcagggca gaagcaggaa ctccgctcct 840
tctggagcta cgctttctct cggcgctttg cctggcacct ggtgtggctg tctgtgatac 900
agttgtggca ctacctcttc attggcactc tcaactcctt gctgaccaac atggccggtg 960
gggacatggc acgagtcagc acctacacaa atgcctttgc cttcactcag ttcggagtgc 1020
tgtgtgcccc ctggaatggc ctgctcatgg accggcttaa acagaagtac cagaaggaa 1080
caagaaagac aggttcctcc actttggcgg tggcctctg ctcgacggtg ccttcgctgg 1140
ccctgacatc cctgctgtgc ctgggcttcg ccctctgtgc ctgagtcctc atcctccctc 1200
tccagtacct cacccttcac ctgcaagtga tcagccgctc ctctctctat gggagcaacg 1260
cggccttctc cacccttgct ttcccttcag agcactttgg caagctcttt gggctgggta 1320
tggccttgte ggctgtgggtg tctctgtctc agttccccat cttcaccctc atcaaaggct 1380
cccttcagaa tgacccattt tacgtgaatg tgatgttcat gcttgccatt cttctgacat 1440
tcttccaccc ctttctggta tatcggaat gccgtacttg gaaagaaagt ccctctgcaa 1500
ttgcatagtt cagaagcctt cacttttcag ccccgaggat ggttttgttc atcttccacc 1560
acctttgagg acctcggtgc ccaaaagact ttgcctatcc cagcaaaaaca cacacacaca 1620
cacacacaca caaaaataaa agacacacaa ggacgtctgc gcagcaagaa aagaatctca 1680
gttgccaagc agattgatat cacacagact caaagcaaag gcatgtggaa cttctttatt 1740
tcaaaacaga agtgtctcct tgcacttagc cttggcagac ccttgactcc aggggagatg 1800
acctggggga ggaagtgtgt caactatttc tttaggcctg tttggctccg aagcctatat 1860
gtgcctggat cctctgccat ccctcaagag tgcagccttg gctagagaac tcacagctct 1980
cagctatgct tcctggctct ccctcaagag tgcagccttg gctagagaac tcacagctct 1980
gggaaaaaga ggagcagaca ggggttcctg gggcccttg cagcccagcc actgatgctg 2040
gatgaccttg gcctgacctt ggtctgggtc cagaatcact tttcccatct gtaaaattga 2100
gatgaatttt ggtgttgaaa gttcttctct gagcagatgt cctagaaggt tttaggaata 2160
gtgacagagt caggccaccc caagggccat gggagccagc tgacctgctt gaccgaagga 2220
tttctgacag actatctttg gggatgtttt caagaaggga tataagttat ttactttggg 2280
catttaaaag aaaatttctc tcgggaataa ttttatagaa aaataaagct tctgtgtcta 2340
aggcaaaaaa aaaaaaaaaa 2359
```

<210> 50

<211> 2052

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 977658CB1

<400> 50

```
gggtttgaag ccgcgccgcg agggagcgag gtcgcagtga cagcggcggg cgatcggacc 60
caggetgccc cgccgtaccc gcctgcgtcc cgcgctcccc cccagcatg acagccccgg 120
cgggtccgcg cggtcagag accgagcggc ttctgacccc caacccccgg tatgggaccc 180
aggcggggcc ttcaccggcc cctccgacac ccccagaaga ggaagacctt cgccgtcgct 240
tcaaatactt tttcatgagt ccttgcgaca agtttctgagc caagggccgc aagccctgca 300
agctgatgct gcaagtgggt aagatcctgg tggtcacggt gcagctcatc ctgtttgggc 360
tcagtaatca gctggctgtg acattccggg aagagaacac catcgccctc cgacacctct 420
```

```

tcctgctggg ctactcggac ggagcggatg acaccttcgc agcctacacg cgggagcagc 480
tgtaccaggc catcttccat gctgtggacc agtacctggc gttgcctgac gtgtcactgg 540
gccggtatgc gtatgtccgt ggtgggggtg acccttggac caatggctca gggcttgctc 600
tctgccagcg gtactaccac cgaggccacg tggaccggc caacgacaca tttgacattg 660
atccgatggt ggttactgac tgcattccagg tggatcccc cgagcggccc cctccgcccc 720
ccagcgacga tctcaccctc ttggaaagca gctccagtta caagaacctc acgctcaaat 780
tccacaagct ggtcaatgtc accatccact tccggctgaa gaccattaac ctccagagcc 840
tcatcaataa tgagatcccc gactgctata ccttcagcgt cctgatcacg tttgacaaca 900
aagcacacag tgggaggatc cccatcagcc tggagaccca ggcccacatc caggagtgtg 960
agcacccccag tgtcttccag cacggagaca acagcttccg gctcctgttt gacgtgggtg 1020
tcatcctcac ctgctccctg tcttctctcc tctgcgccc gctcactcct cgaggcttcc 1080
tgctgcagaa tgagtttgtg gggttcatgt ggcggcagcg gggacgggtc atcagcctgt 1140
gggagcggct ggaatttgtc aatggctggt acatcctgct cgtcaccagc gatgtgctca 1200
ccatctcggg caccatcagc aagatcggca tccaggccaa gaacttggcg agctacgacg 1260
tctgcagcat cctcctgggc acctcgacgc tgcgtggtgtg ggtgggctg atccgctacc 1320
tgaccttctt ccacaactac aatatactca tccccacact gcgggtggcc ctgccacg 1380
tcatgcgctt ctgctgctgc gtggctgtca tctacctggg ctactgcttc tgtggctgga 1440
tcgtgctggg gccctatcat gtgaagtccc gctcactctc catggtgtct gactgctgt 1500
tctcgctcat caatggggag gacatgtttg tgacgttcgc cgccatgcag gcgcagcagg 1560
gccgcagcag cctgggtgtg ctcttctccc agctctacct ttactccttc atcagcctct 1620
tcatctacat ggtgctcagc ctcttcatcg cgctcatcac cggcgccctac gacacctca 1680
agcatcccgg cggcgagcgc gcagaggaga gcgagctgca ggccctacat gcacagtgcc 1740
aggacagccc cactcgggc aagttccgcc gcgggagcgg ctcgccctgc agccttctct 1800
gctgctgcgg aagggacccc tccgaggagc attcgcgtgt ggtgaattga ttcgacctga 1860
ctgccgttgg accgtaggcc ctggactgca gagacccccg ccccgacccc cgcttattta 1920
ttgttagggt ttgcttttaa ggatcggctc cctgtcgcgc ccgaggaggg cctggacctt 1980
tcgtgtcgga cccttggggg cggggagact ggggtggggag ggtgttgaat aaaaggga 2040
ataaaaaaaaa aa 2052

```

<210> 51

<211> 1939

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1004703CB1

<400> 51

```

cttgatggcg tcgggctgga gagccgcagt cccggctgca gcacctggga gaaggcagac 60
cgtgtgaggg ggcctgtggc ccagcgtgct gtggcctcgg ggagtgggaa gtggaggcag 120
gagccttctt tacacttcgc catgagtttc ctcatcgact ccagcatcat gattacctcc 180
cagatactat tttttggatt tgggtggcct ttcttcagtc gccaatgtgt taaagactat 240
gagatacgtc agtatgttgt acaggtgata ttctccgtga cgtttgcatt ttcttgacc 300
atgtttgagc tcatcatctt tgaaatctta ggagtattga atagcagctc ccgttatttt 360
cactggaaaa tgaacctgtg tgtaattctg ctgactcctg ttttcatggg gcctttttac 420
attggctatt ttattgtgag caatatccga ctactgcata aacaacgact gcttttttcc 480
tgtctcttat ggctgacctt tatgtatttc ttctggaaac taggagatct ctttcccatt 540
ctcagcccaa aacatgggat cttatccata gaacagctca tcagccgggt tgggtgtgatt 600
ggagtgactc tcatggctct tctttctgga ttgtgtgctg tcaactgccc atacacttac 660
atgtcttact tctcaggaa tgtgactgac acggatattc tagccctgga acggcgactg 720
ctgcaaacca tggatatgat cataagcaaa aagaaaagga tggcaatggc acggagaaca 780
atgttccaga agggggaagt gcataacaaa ccatcagggt tctggggaat gataaaaagt 840
gttaccactt cagcatcagg aagtgaatat cttactctta ttcaacagga agtggatgct 900
ttggaagaat taagcagga gctttttctg gaaacagctg atctatatgc taccaaggag 960
agaatagaat actccaaaac cttcaagggt aaatatctta attttcttg ttactttttc 1020
tctatttact gtgtttggaa aattttcatg gctaccatca atattgttt tgatcgagtt 1080
gggaaaacgg atcctgtcac aagaggcatt gagatcactg tgaattatct gggaatccaa 1140
tttgatgtga agttttggc ccaacacatt tcttctattc ttgttggaat aatcatcgct 1200
acatccatca gaggattgct gatcactctt accaagttct tttatgccat ctctagcagt 1260
aagtctcca atgtcattgt cctgctatta gcacagataa tgggcatgta ctttgtctcc 1320
tctgtgctgc tgatccgaat gagtatgctt ttagaatacc gcaccataat cactgaagtc 1380
cttgaggaaac tgcagttcaa cttctatcac cgttgggttg atgtgatctt cctggctcag 1440

```

```

gctctctcta gcatactctt cctctatttg gctcacaaac aggcaccaga gaagcaaatg 1500
gcaccttgaa cttaagccta ctacagactg ttagaggcca gtggtttcaa aatttagata 1560
taagaggggg gaaaaatgga accagggcct gacattttat aaacaaacaa atgctatgg 1620
tagcattttt caccttcata gcatactcct tccccgtcag gtgatactat gaccatgagt 1680
agcatcagcc agaacatgag agggagaact aactcaagac aatactcagc agagagcatc 1740
ccgtgtggat atgaggctgg tgtagaggcg gagaggagcc aagaaactaa aggtgaaaaa 1800
tacactggaa ctctggggca agacatgtct atggtagctg agccaaacac gtaggatttc 1860
cgttttaagg ttcacatgga aaagggtata gctttgcctt gagattgact cattaataatc 1920
agagactgta aaaaaaaaaa

```

<210> 52

<211> 1138

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1334051CB1

<400> 52

```

caaactgcaa cttatatctg caatttattt tggatatagac aagaggtatg ccagtagcac 60
actgggtggc tcagaagaaa ttctcaacac cttagctcgcc agagagtcta tgtatgggat 120
tgaacaatct gtaaaactaa ggatcctaata catgaaaata agtatgataa attataagtc 180
actattggca ctgttggtta tattagcctc ctggatcatt ttacagttt tccagaactc 240
cacaaagggt tggctctgctc taaacttatc catctccctc cattactgga acaactccac 300
aaagtcctta ttccctaaaa caccactgat atcattaaag ccactaacag agactgaact 360
cagaataaag gaaatcatag agaaactaga tcagcagatc ccaccagac ctttcaccca 420
cgtgaacacc accaccagcg ccacacatag cacagccacc atcctcaacc ctcgagatac 480
gtactgcagg ggagaccagc tgcacatcct gctggagggtg agggaccact tgggacgcag 540
gaagcaatat ggcggggatt tcctgagggc caggatgtct tcccagcgc tgatggcagg 600
tgcttcagga aagggtgactg acttcaacaa cggcacctac ctggtcagct tcactctgtt 660
ctgggagggc caggtctctc tgtctctgct gctcatccac ccagtgaaag ggggtgtcagc 720
tctctggagt gcaaggaacc aaggctatga cagggtgatc ttcactggcc agtttgtcaa 780
tggcacttcc caagtccact ctgaatgtgg cctgatccca aacacaaatg ctgaattgtg 840
ccagtacctg gacaacagag accaagaagg cttctactgt gtgaggcctc aacacatgcc 900
ctgtgctgca ctactcaca tgtattctaa gaacaagaaa gtttcttata ttagcaaaca 960
agaaaagagc ctctttgaaa ggtaaaaata attacttctt gagactacct gtgcaaatat 1020
tgtgatttgg cctatatact gatccaaaga aaagtcttgt gagtgattta attttgggtg 1080
tctttagtaa gagcctttgg ggaaaggatc tgtgaattca tttagagaca gtgccccat 1138

```

<210> 53

<211> 2117

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1336728CB1

<400> 53

```

tggccacagc aacaccact gagcacgtg ggagctgagt atggcgctccc tggctcgtc 60
ggagctgggg ctgcttctgg ctgtgctggg ggtgacggcg acggcgctccc cgctgctgg 120
tctgctgagc ctgctcacct ctggccaggg cgctctggat caagaggctc tgggaggcct 180
gttaaatacg ctggcggacc gtgtgactg caccaacggg ccgtgtggaa agtgctgtc 240
tgtggaggac gccctgggccc tgggcgagcc tgaggggtca gggctgcccc cgggcccgg 300
cctggaggcc aggtacgtcg ccgcctcag tgccgcgcgc gtectgtacc tcagcaacce 360
cgagggcacc tgtgaggaca ctcgggctgg cctctgggccc tctcatgcag accacctcct 420
ggccctgctc gagagcccca aggcctgac ccggggcctg agctggctgc tgcagaggat 480
gcaggcccgg gctgccggcc agaccccaa gacggcctgc gtagatatcc ctgagctgct 540
ggaggaggcg gtgggggcgg gggctccggg cagtgtggc ggcgtcctgg ctgccctgct 600
ggaccatgtc aggagcgggt cttgcttcca cgccttgccg agccctcagt acttcgtgga 660
ctttgtgttc cagcagcaca gcagcgaggt ccctatgacg ctggccgagc tgtcagcctt 720
gatgcagcgc ctgggggtgg gcaggaggcc ccacagtgc cacagtcac ggcaaggagg 780

```

```

agccagcagc cgggaccctg tgccctcat cagctccagc aacagctcca gtgtgtggga 840
cacggtatgc ctgagtcca gggacgtgat ggctgcatat aacaggctgg 900
ggtgaccccg gaggcctggg cccaactgag ccctgccctg ctccaacagc agctgagtg 960
agcctgcacc tcccagtcca gggccccctg ccaggaccag ctccagccagt cagagaggta 1020
tctgtacggc tccctggcca cgctgctcat ctgcctctgc gcggtctttg gcctcctgct 1080
gctgacctgc actggctgca ggggggtcac ccactacatc ctgcagacct tcctgagcct 1140
ggcagtgggt gcactcactg gggacgtgt cctgcatctg acgcccagg tgctggggct 1200
gcatacacac agcgaagagg gcctcagccc acagcccacc tggcgctcc tggctatgct 1260
ggcggggctc tacgccttct tcctgtttga gaacctctt aatctcctgc tgcccaggga 1320
cccggaggac ctggaggacg gggcctgcgg ccacagcagc catagccacg gggggccacag 1380
ccacggtgtg tccctgcagc tggcaccag cgagctccgg cagcccagg cccccacga 1440
gggctcccgc gcagacctgg tggcggagga gagcccgag ctgctgaacc ctgagcccag 1500
gagactgagc ccagagttga ggctactgcc ctatatgatc actctggcg acgctgtgca 1560
caacttcgcc gacgggctgg ccgtggggcg cgccttcgag tcctcctgga agaccgggct 1620
ggccacctgc ctggccgtgt tctgccacac gttgccacac gagctggggg acttcgccgc 1680
cttgcctgac gcggggctgt ccgtgcgcca agcactgctg ctgaacctgg cctccgcgt 1740
cacggccttc gctggctctc acgtggcact cgcggttgga gtcagcgagg agagcgaggc 1800
ctggatcctg gcagtggcca ccggcctgtt cctctacgta gcactctgcg acatgctccc 1860
ggcgatgttg aaagtacggg acccgcggcc ctggctcctc ttcctgctgc acaacgtggg 1920
cctgctgggc ggctggaccg tcctgctgct gctgtccctg tacgaggatg acatcacctt 1980
ctgataccct gccctagtcc ccacaccttg acttaagatc ccacacctca caaacctaca 2040
gccagaaac cagaagcccc tatagaggcc ccagtcctaa ctccagtaaa gacactcttg 2100
tccttggaag aaaaaaa 2117

```

<210> 54

<211> 1495

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1452856CB1

<400> 54

```

ctgaaatccc gcgaggatca accgagctcg ccgaaaggag ggaggaacgt atcccttctg 60
gaggctgtct cagggggcag agggaccgga ccggaagtga cgtgagcggg ttccggttgt 120
ctggagccca gcggcggtg tgagagtccg taaggagcag cttccaggat cctgagatcc 180
ggagcagccg gggctcggagc ggctcctcaa gagttactga tctatgaaat ggcagagaat 240
ggaaaaaatt gtgaccagag acgtgtagca atgaacaagg aacatcataa tggaaatttc 300
acagaccctt cttcagtga tgaagaag aggaggagc gggaagaaag gcagaatatt 360
gtcctgtgga gacagccgct cattaccttg cagtattttt ctctggaaat ccttgaatc 420
ttgaaggaat ggacctcaaa attatggcat cgtcaaagca ttgtggtgtc ttttttactg 480
ctgcttgctg tgcttatagc tacgtattat gttgaaggag tgcatacaaa gtatgtgcaa 540
cgtatagaga aacagtttct tttgtatgcc tactggatag gcttaggaat tttgtcttct 600
gttgggcttg gaacagggtt gcacaccttt ctgctttatc tgggtccaca tatagcctca 660
gttacattag ctgcttatga atgcaattca gttaattttc ccgaaccacc ctatcctgat 720
cagattattt gtccagatga agagggcact gaaggaacca tttctttgtg gagtatcatc 780
tcaaaagtta ggattgaagc ctgcatgtgg ggtatcggta cagcaatcgg agagctgcct 840
ccatatttca tggccagagc agctcgcctc tcagggtgctg aaccagatga tgaagagtat 900
caggaatttg aagagatgct ggaacatgca gagtctgcac aagactttgc ctcccgggcc 960
aaactggcag ttcaaaaact agtacagaaa gttggatttt ttggaatttt ggctgtgct 1020
tcaattccaa atccttttatt tgatctggct ggaataacgt gtggacactt tctggtacct 1080
ttttggacct tctttggtgc aaccctaatt ggaaaagcaa taataaaaat gcataatccag 1140
aaaatttttg ttataataac attcagcaag cacatagtgg agcaaattgg ggctttcatt 1200
ggtgctgtcc ccggcatagg tccatctctg cagaagccat ttcaggagta cctggagggt 1260
caacggcaga agcttcacca caaaagcgaa atgggcacac cacagggaga aaactggttg 1320
tcctgcatgt ttgaaaagtt ggtcgttgct atggtgtgtt acttcacctc atctatcct 1380
aactccatgg gcataagtta tgccaaacga atccagcagc ggttgaaactc agaggagaaa 1440
actaaataag tagagaaagt tttaaactgc agaaattgga gtggatgggt tctgc 1495

```

<210> 55

<211> 1747

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1562471CB1

<400> 55

accagcagaa	ggctgggagt	ctgtagtttg	ttcctgctgc	caggctccac	tgaggggaac	60
ggggacctgt	ctgaagagaa	gatgcccctg	ctgacactct	acctgtctct	cttctggctc	120
tcaggctact	ccattgccac	tcaaatacacc	ggtccaacaa	cagtgaatgg	cttggagcgg	180
ggctccttga	ccgtgcagtg	tgtttacaga	tcaggctggg	agacctactt	gaagtgggtg	240
tgctcaggag	ctatttggcg	tgactgcaag	atccttggtta	aaaccagtgg	gtcagagcag	300
gaggtgaaga	gggaccgggt	gtccatcaag	gacaatcaga	aaaaccgcac	gttactgtg	360
accatggagg	atctcatgaa	aactgatgct	gacacttact	gggtggaat	tgagaaaact	420
ggaaatgacc	ttgggggtcac	agttcaagtg	accattgacc	cagcaccagt	caccaagaa	480
gaaactagca	gctccccaac	tctgaccggc	caccacttgg	acaacaggca	caagctcctg	540
aaagctcagt	tcctcctgcc	cctcatcttc	accatattgc	tgctgctttt	ggtggccgcc	600
tcactcttgg	cttggaggat	gatgaagtac	cagcagaaag	cagccgggat	gtccccagag	660
caggtactgc	agcccctgga	gggagacctc	tgctatgcag	acctgacctt	gcagctggcc	720
ggaacctccc	cgcgaaaggc	taccacgaag	ctttcctctg	cccagggtga	ccagggtgaa	780
gtggaatatg	tcaccatggc	ttccttgccg	aaggaggaca	tttcttatgc	atctctgacc	840
ttgggtgctg	aggatcagga	accgacctac	tgcaacatgg	gccacctcag	tagccacctc	900
cccggcaggg	gccctgagga	gcccacggaa	tacagacca	tcagcaggcc	ttagcctgca	960
ctccaggctc	cttcttggac	cccaggctgt	gagcacactc	ctgcctcatc	gaccgtctgc	1020
cccctgctcc	cctcatcagg	accaaccccg	ggactgggtg	ctctgcctga	tcagccagca	1080
ttgcccctag	ctctgggttg	ggcttggggc	caagtctcag	ggggcttcta	ggagtggggg	1140
ttttctaaac	gtcccctcct	ctcctacata	gttgaggagg	gggctaggga	tatgctctgg	1200
ggctttcatg	ggaatgatga	agatgataat	gagaaaaatg	ttatcattat	tatcatgaag	1260
taccattatc	ataatacaat	gaacctttat	ttattgccta	ccacatgtta	tggtctgaat	1320
aatggccccc	aaagatatct	gtgtccta	cctcagaact	tgtgactgtt	accttctgtg	1380
gcagaaaggg	acagtgcaga	tgtatgtaag	ttaaggactt	tgagatagag	aggttattct	1440
tgctgattca	ggtgggcccc	aaatatcacc	acaagggtcc	tcataagaaa	gaggccagaa	1500
ggtcaaagag	gtagagacaa	agtgatgatg	gaagtggacg	tggtgtgac	gtgagcaggg	1560
gccatgaatg	ccgcagcctt	cagatgccag	aaagggaaag	gaatggattc	ccctgcctgg	1620
agcctccaaa	agaaaccagc	cctgcccacg	ccttgacttg	agccatttga	aactgatctt	1680
gagctcctgg	cctccagaat	tgcaggagaa	ttaatttgtg	ttgtttttaa	tgaaaaaaaa	1740
aaaaaaa						1747

<210> 56

<211> 1473

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1618158CB1

<400> 56

acagcgggac	agcccggcag	ccttgggctt	ccagcgtg	cttcatcgcc	acctccctgc	60
cttctccac	ttccgcattc	ttaggtgacg	gatgggtggca	ccatcaagca	aaagatcttc	120
accttcgacg	ccatgttctc	caccaactac	tcacacatgg	agaactaccg	caagcgagag	180
gacctggtgt	accagtccac	tgtgaggctg	cccagggtcc	ggatctcaga	caatgggtccc	240
tatgagtgcc	atgtgggcat	ctacgaccgc	gccaccaggg	agaagggtgt	cctggcatca	300
ggcaacatct	tcctcaacgt	catggctcct	cccacctcca	ttgaagtgtt	ggctgctgac	360
acaccagccc	ccttcagccg	ctaccaagcc	cagaacttca	cgctgggtctg	catcgtgtct	420
ggaggaaaaa	cagcaccatc	ggtttatttc	aaacagatg	gggaaccaat	cgacgcagtg	480
cccctatcag	agccaccagc	tgcgagctcc	ggccccctac	aggacagcag	gcccttccgc	540
agccttctgc	accgtgacct	ggatgacacc	aagatgcaga	agtcactgtc	cctcctggac	600
gccgagaacc	gggggtggcg	accctacacg	gagcgccctt	cccgtggcct	gacccagat	660
cccaacatcc	tcctccagcc	aaccacagag	aacataccag	agacgggtcgt	gagccgtgag	720
tttccccgct	gggtccacag	cgccgagccc	acctacttcc	tgcgccacag	ccgcaccccg	780
agcagtgcag	gcactgtgga	agtacgtgcc	ctgctcacct	ggaccctcaa	cccacagatc	840
gacaacgagg	ccctcttcag	ctgcgaggtc	aagcaccag	ctctgtcgat	gcccagtcag	900

```

gcagaggtca cgctggttgc ccccaaagga cccaaaattg tgatgacgcc cagcagagcc 960
cgggtagggg acacagttag gattctgttc catgggtttc agaacgaagt cttcccggag 1020
cccatgttca cgtggacgcg ggttgggagc cgcctcctgg acggcagcgc tgagtctgac 1080
gggaaggagc tgggtgctgga gcgggttccc gccgagctca atggctccat gtatcgctgc 1140
accgcccaga acccactggg ctccaccgac acgcacaccc ggctcatcgt gtttgaaaac 1200
ccaaatatcc caagaggaac ggaggactct aatggttcca ttggccccac tgggtgcccg 1260
ctcaccttgg tgctcgccct gacagtgatt ctggagctga cgtgaaggca cccgccccgg 1320
ccactccatc aggcactgac atctccacga ccgggttttc tttcttttct aaactatttc 1380
cagtcttggt cttagtctct ttccatctgt gtcttggtct cttcagtcgg ttttaattaa 1440
acaaacagaa caattttccc caaaaaaaaaa aaa 1473

```

<210> 57

<211> 1591

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1656935CB1

<400> 57

```

attaaacgac tcactatagg gaatttggcc ctcgaggcaa gaattcggca cgagggcttc 60
tataggagag ctttagaatg aagtcattta gaggagcagg cgaatcctaa ccgcatctct 120
ctcttttagc ggactgaacc caaacatgaa tgtcaacagc atggacatga ccggtggctt 180
gtcgggtgaag gacccatccc agtcccagtc acgcctcccc cagtggacgc accccaactc 240
catggataac ttgcccagtg ccgcttcccc cctggagcag aaccctagca agcatggtgc 300
tatccctgga ggtctaagca ttgggcctcc aggtaagtcc tccattgatg actcctatgg 360
ccggtacgat ttaatccaga acagtgagtc accagccagt cctcccgtag ctgttcccca 420
tagctgggtca cgtgccaaat ctgacagtga taaaatctca aatggctcta gcatcaactg 480
gccccagaa ttccatccgg gagttccatg gaaaggactg cagaatattg accctgagaa 540
tgacctgac gtcactcctg gcagtgtccc cactgggcct accatcaaca ccaccatcca 600
ggatgtcaac cgctacctcc tcaagagtgg agggctctcc ccgcatcatc ctcagaatgc 660
cacgctgect tcttcgagtg cctggccact cagtgcctcc ggctacagta gctctttcag 720
cagcattgca tccgcacctc gtgttgagg taaactgtca gacatcaaat cgacgtggtc 780
ctctggccct acctcccaca cgcaagcctc tctgtctcat gaactatgga aggtgcccag 840
aaacagtact gcacccacga ggccacctcc aggggttaacc aatcccaage cctcctccac 900
ctgggggtgcc agccccctcg gctggaccag ctctactccc tcgggttctg cctggagcac 960
cgacacctca ggaagaacca gcagctggct cgttcttcga aacctcactc ccagattga 1020
tgggttctaaa ctgaggacat tgtgtttgca acatgggcct cttatcacat tccacctgaa 1080
tctgactcaa ggcaatgctg tgggtccggtc cagctccaag gaggagggtc tgccaaaggc 1140
ccaggaagtc ctctgcacaa ttgtgctgcc ttgggagaca cttagccatt ccttgggccc 1200
gagtttccgc ttggttgga cgaaggaagt tgggaattcgg gtttccttta agccgccaga 1260
gggcccaggc cgcattggcc agagcacaat ttccaagggt ctggcacaat tccacgacca 1320
gaggggcggt tccaagctca caggacgtgg cgggattcac aggccacgcg ggaggggcaa 1380
ggcttctcac caacttgac atatgagaca ttgtgaatta acattttaag agcgttgcca 1440
ctaattggga caatcatcgg acaaggaccc tgagttcaat tttgtaaagt gccctgttat 1500
aaatgtttgc cgtgtagaat aaagggttct tccactata tccattgta tcgggatata 1560
tcacatcgta acagtgttat tttgggaaga a 1591

```

<210> 58

<211> 1858

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1859305CB1

<400> 58

```

gcaggcgcgg cgaaacttcc ctctacccgc ccggccccgc gcgcgcaccg ttggcgctgg 60
acgcttcttc cttggaagcg cctctccctc aggtctcaag aactgttttg agatgcagga 120
attcatctaa ttttactgac cgggcgaggt gtgagagccc tagcatctga aagtggtcga 180
cttgcgagtt gttatggaga aaacttgat agatgcactt cctcttacta tgaattcttc 240

```

agaaaagcaa	gagactgtat	gtattttttg	aactgggtgat	tttggaagat	cactgggatt	300
gaaaatgctc	cagtgtggtt	attctgttgt	ttttggaagt	cgaaaccccc	agaagaccac	360
cctactgccc	agtgggtgcag	aagtcttgag	ctattcagaa	gcagccaaga	agtctgaçat	420
cataatcata	gcaatccaca	gagagcatta	tgatttttctc	acagaattaa	ctgagggttct	480
caatggaaaa	atattggtag	acatcagcaa	caacctcaaa	atcaatcaat	atccagaatc	540
taatgcagag	taccttgctc	atttggtgcc	aggagcccac	gtggtaaaaag	catttaacac	600
catctcagcc	tgggctctcc	agtcaggagc	actggatgca	agtcggcagg	tgtttggtg	660
tggaaatgac	agcaaagcca	agcaaagagt	gatggatatt	gttcgtaatc	ttggacttac	720
tccaatggat	caaggatcac	tcatggcagc	caaagaaatt	gaaaagtacc	ccctgcagct	780
atttccaatg	tggagggttcc	ccttctattt	gtctgctgtg	ctgtgtgtct	tcttgttttt	840
ctattgtgtt	ataagagacg	taatctacce	ttatgtttat	gaaaagaaaag	ataatacatt	900
tcgtatggct	atttccattc	caaatcgtat	ctttccaata	acagcactta	cactgcttgc	960
tttggtttac	ctccctgggtg	ttattgctgc	cattctacaa	ctgtaccgag	gcacaaaata	1020
ccgtcgattc	ccagactggc	ttgaccactg	gatgctttgc	cgaaagcagc	ttggcttgggt	1080
agctctggga	tttgccctcc	ttcatgtcct	ctacacactt	gtgattccta	ttcgatatta	1140
tgtacgatgg	agattgggaa	acttaaccgt	taccagcgca	atactcaaga	aggagaatcc	1200
atttagcacc	tcctcagcct	ggctcagtg	ttcatatgtg	gctttgggaa	tacttgggtt	1260
ttttctgttt	gtactcttgg	gaatcacttc	tttgccatct	gttagcaatg	cagtcaactg	1320
gagagagttc	cgattttgtcc	agtcctaaact	gggttatttg	accctgatct	tggtgtacagc	1380
ccacaccctg	gtgtacgggtg	ggaagagatt	cctcagccct	tcaaattctca	gatggatctc	1440
tcctgcagcc	tacgtgttag	ggcttatcat	tccttgctact	gtgctgggtga	tcaagtttgt	1500
cctaactcatg	ccatgtgttag	acaacaccct	tacaaggatc	cgccagggct	gggaaaggaa	1560
ctcaaaacac	tagaaaaagc	attgaatgga	aatcaatat	ttgaaacaaa	gttcaattta	1620
gctggatttc	tgaactatgg	ttttgaatgt	ttaaagaaga	atgatgggta	cagttaggaa	1680
agtttttttc	ttacaccgtg	actgagggaa	acattgcttg	tctttgagaa	attgactgac	1740
atactggaag	agaacaccat	tttatctcag	gttagtgaag	aatcagtgca	ggtcctcgac	1800
tcttattttc	ccagaggcca	tggagctgag	attgagacta	gccttggtgt	tttcacac	1858

<210> 59

<211> 1454

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1949083CB1

<400> 59

caccctgggtg	tcttcccctg	tggccctgca	aggcatatca	tgettctgt	ctctaggacc	60
tgcttactag	aaagcagcac	ccgcctgaaa	cctcacgaag	cccagaacta	caggaagaag	120
gcattgtggg	tgctctgggt	ctccatcatt	gtcacccctg	ccctcgcggt	ggctgccttt	180
actgtctccg	ttatgaggta	cagcgctct	gcttttggtt	ttgcatttga	tgccatcctg	240
gacgtcctgt	catcggcgat	tgctctgtgg	cgttacagca	acgcggccgc	tggtgcactc	300
gccccatagg	agtacatagc	ctgtgtcctc	ttgggggtga	tattccttct	gtcatccata	360
tgtatagtgg	tcaaagccat	ccatgacctc	tcaactaggc	tgctcccaga	agtggacgat	420
ttcctgttca	gtgtctccat	tttaagtggg	attcttttga	gcacccctgg	cgtgttgaag	480
ttcatgctgg	ggaagggttct	gaccagtaga	gcactcataa	cagatgggtt	taactccctc	540
gtgggtggcg	tgatgggctt	ctccattctt	ctgagcgcg	aagtgttcaa	gcattgactcg	600
gcggtctgggt	acctggacgg	cagcataggc	gttctgatcg	gcctcaccat	atttgcctat	660
ggggtcaaac	tcctcatcga	catggtgcgg	aagggtgagg	agacacgtca	ctacgagatg	720
tttgagtga	gggggccagc	atccgcatga	gaccatcgag	atgaggagtt	cccacatagg	780
caaagggtgc	caatatttta	ctgaacatct	ggtttctttt	tggaaagttt	ctttcacatg	840
gtttgtcatt	acaagacaag	gtctgcccag	ccagggtggat	ctaccttgcc	cccatcacct	900
gccgccccca	tcaaacatgt	tgggacaatg	cccataggaa	tggacctcct	tccccgtctc	960
cagctggggac	tggcgttttt	tagtctctgg	agtatgatgg	ttctcatggg	taggatgaga	1020
tctttggcag	aaaggtcttc	ggtgggtgctc	tgagcctgcg	ctgcatagga	ctgacgagac	1080
ccacctctc	cagcttgggt	ggccctggcca	cccttggttc	caagtctctc	ctttcctggc	1140
aggtcttaag	ggaagattgt	acccctcacc	ctttacatac	ccagaatcat	cagtatgtca	1200
cttcctaatt	tctatcagtg	tatctcatta	tttcatactg	ttttactaat	cctaagtcta	1260
aacagatttg	ctcaaaagga	gaccattcta	ttttttaaag	tacttagtga	tacacgtata	1320
agctttgcat	ggacgaatta	aataagcaca	ttgacctttt	cttgtagatt	cagaacctga	1380
acatccatgt	gaaaactggg	tccatttttg	agagatgtga	aactacagtt	tatatgtaat	1440
aaataaatat	aata					1454

<210> 60
 <211> 2310
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1996357CB1

<400> 60
 ggccccggatg ttcgggtgcag ctgccagatc cgctgatcta gtgctttctcg aaaaaaacct 60
 tcaggcggcc catggctgtc gatattcaac cagcatgcct tggactttat tgtgggaaga 120
 ccttattatt taaaaatggc tcaactgaaa tatatggaga atgtggggta tgcccaagag 180
 gacagagAAC gaatgcacag aaatattgtc agccttgac agaactctct gaactttatg 240
 attggctcta tcttgattt atggcaatgc ttcctctggg ttacattgg ttcttcattg 300
 aatggctact ggggaaaaag agttccagcg cacttttcca acacatcact gcattatttg 360
 aatgcagcat ggcagctatt atcaccttac ttgtgagtga tccagttggg gttctttata 420
 ttcgttcatg tcgagtatt atgctttctg actggtacac gatgctttac aaccacaagt 480
 cagattacgt taccacagta cactgtactc atgaagccgt ctaccacta tataccattg 540
 tatttatcta ttacgcattc tgcttggtat taatgatgct gctccgacct cttctgggtga 600
 agaagattgc atgtgggtta gggaaatctg atcgatttaa aagtatttat gctgcacttt 660
 acttcttccc aattttaacc gtgcttcagg cagttgggtg aggcctttta tattacgcct 720
 tcccatacat tatattagt ttatctttgg ttactctggc tgtgtacatg tctgcttctg 780
 aaatagagaa ctgctatgat cttctgggtc gaaagaaaag acttattgtt ctcttcagcc 840
 actggttact tcatgcctat ggaataatct ccatttccag agtggataaa cttgagcaag 900
 atttgcccc tttggctttg gtacctacac cagccctttt ttacttggtc actgcaaaat 960
 ttaccgaacc ttcaaggata ctctcagaag gagccaatgg acactgagt tagacatgtg 1020
 aaatgccaaa aacctgagaa gtgctcctaa taaaaaagta aatcaatctt aacagtgtat 1080
 gagaactatt ctatcatata tgggaacaag attgtcagta tatcttaatg tttgggtttg 1140
 tctttgtttt gtttatggtt agacttacag acttgaaaa tgcaaaactc tgaataactc 1200
 tgttacacag gtaaatatta tctgctacac tggaaggccg ctaggagcc cttgcttctc 1260
 tcaacagttc agctgttctt tagggcaaaa tcatgttctt gtgtacctag caatgtgttc 1320
 ccattttatt aagaaaagct ttaacacgtg taatctgcag tcttaacag tggcgtaatt 1380
 gtacgtacct gttgtgtttc agtttgtttt tcacctataa tgaattgtaa aaacaaacat 1440
 acttggtggg tctgatagca aacatagaaa tgatgtatat tgttttttgt tatctattta 1500
 ttttcatcaa tacagtatt tgatgtattg caaaaataga taataattta tataacaggt 1560
 tttctgttta tagattggtt caagatttgt ttggattatt gttcctgtaa agaaaacaat 1620
 aataaaaaagc ttacctacat aaaatttcaa tgttttgaca ctttaattgtt gtttggcaca 1680
 atagtatgga agtaattcaa actggtaaat agtttctct catatctcgg gtatatatac 1740
 ataccatatt ttattgatcc agagatactt atttcacttt gtgacatctc tgaattagga 1800
 tgcattctac aactgatggc ttattaggtt taatgaaata cagaagatac acagtataaa 1860
 aagggttttc ctgtggttgg tttgtggttt gtgatagggt tctgtgatg tttatgcttt 1920
 gaaggcttta agactcatgg ttgcaaccat ggaagcaaaa tgaaattttt agctcttaac 1980
 ctaacaacct gaccatgttt atccattttt attgtttaga agtttattta ctgatacttg 2040
 gtggagggtg tgtgaattag ttaaatttta aatgtttaag acttctatta acagctgcaa 2100
 aatatgaaag taagtgcact cacttttccg gtatgtagt gtcttttgaa ttcacagcag 2160
 ttgtatcctt gagttacttt gttaatgtat ttttctcagt acatttaacc actgggaaat 2220
 gaacccttgt acgaatgtgt ttcttcttct ctgtaggaa aaaaaataaa tataaaaatt 2280
 ttatttgtat tgcacacaaa aaaaaaaaaa 2310

<210> 61
 <211> 744
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2061330CB1

<400> 61
 agtgggtgtg tacctgccga cagcataaag cgaggcaagg tccagccgtt ccgcccggca 60
 tacaagctta tggagcagcc ccctttgaag atctccaggt ggacttcaca gagatgtcaa 120
 agtgtagagg tgatcgagtg tggatcaaga actggaacgt agcctctttg tgtccactgt 180


```

ggaaaggacc ccagactgtc gttctgagca ctcccaccgc tgtgaaggta gaaggaatcc 240
cagcctggat ccaccacagc catgtaaaac ctgcagcgcc tgaaacctgg gaggcaagac 300
caagcccaga caaccctgc agagtgaccc tgaagaagac gacaagccct gctccagtca 360
caccgggaag ctgactggtc cagcgacggc cgaagcctga ggaagctcat catgagattc 420
atTTTTctta aatTTTggac ttatacagta agggcttcaa ctgatcttac tcaaactggg 480
gactgttccc agtgactca tcaggtcacc gaagtaggac agcaaattaa aacaatcttt 540
ctgttctata gttattatga atgtatggaa acaataaaag aaacttgttt gtataatgcc 600
actcagtaca aggtatgtag cccgagaaat gaccgacctg atgtgtgtta taacccatct 660
gagccccctg caccaccgtt tttgaaataa gaataagaac tggccttttc ctaggtgata 720
caagtaaaat aataactaga acag

```

<210> 62

<211> 1109

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2346947CB1

<220>

<221> unsure

<222> 30

<223> a, t, c, g, or other

<400> 62

```

gaagcagtgc agagaggaga gcgagcgcn agtgccgctg agcaaaggcc ttcaccatgg 60
ccgagtcccc cggctgctgc tccgtctggg cccgctgcct ccactgcctg tatagctgcc 120
actggaggaa atgccccaga gagaggatgc aaaccagcaa gtgcgactgt atctggtttg 180
gcctgctctt cctcaccttc ctcccttccc tgagctggct gtacatcggg ctcgctcttc 240
tcaatgacct gcacaacttc aatgaattcc tcttcgcccg ctggggacac tggatggact 300
ggtccctggc attcctgctg gtcactcttc tactggtcac atatgcatcc ttgctattgg 360
tccctggcct gctcctgcgg ctttgtagac agccccctgca tctgcacagc ctccacaagg 420
tgctgctgct cctcattatg ctgcttggtg cggctggcct tgtgggactg gacatccaat 480
ggcagcagga gtggcatagc ttgctgtgtg cactgcaggc cacagcccca ttccttcata 540
ttggagcagc cgctggaatt gccctcctgg cctggcctgt ggctgatacc ttctaccgta 600
tccaccgaag aggtcccaag attctgctac tgctcctatt ttttgagtt gtccctggta 660
tctacttggc cccctatgac atctcctcac cctgcatcat ggaaccacga gacttaccac 720
ccaagcctgg gctggtggga caccgagggg ccccatgct ggctcccag aacaccctga 780
tgtccttgcg gaagacagct gaatgcggac tactgtgttt gagactgatg tgatggctag 840
ctccgatggg gtcccttccc tcatgcatga tgagcacctc agcaggacca cgaatgtagc 900
ctctgtattc ccaaccgaa tcacagccca cagcagtgc ttctcctgga ctgaactgaa 960
gagactcaat gctggatcct ggttccctaga gaggcgacc ttctgggggg ccaaaccgct 1020
ggcaggccct gatcagaaag aggctgagag tcagtcggta ccagcattag aagagctatt 1080
ggaggaagct gcagccctca accctttca

```

<210> 63

<211> 2511

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2795577CB1

<400> 63

```

gcagccgtcg ccttcggagc gaagggtacc gaccgcggcag aagctcggag ctctcgggggt 60
atcgaggagg caggcccgcg ggcgcacggg cgagcgggcc gggagccgga gcggcggagg 120
agccggcagc agcggcgcgg cgggctccag gcgagggcgt cgacgctcct gaaaacttgc 180
gcgcgcgctc gcggcactgc gcccgagcgc atgaagatgg tcgcgccctg gacgcggttc 240
tactccaaca gctgctgctt gtgctgccat gtccgcaccg gcaccatcct gctcggcgctc 300
tggtatctga tcatcaatgc tgtggtactg ttgattttat tgagtgcctt ggctgacccg 360
gatcagtata acttttcaag ttctgaactg ggaggtgact ttgagttcat ggatgatgcc 420

```

```

aacatgtgca ttgccattgc gatttctctt ctcatgatcc tgatatgtgc tatgggtact 480
tacggagcgt acaagcaacg cgcagcctgg atcatcccat tcttctgtta ccagatcttt 540
gactttgccc tgaacatgtt gggttgcaatc actgtgctta tttatccaaa ctccattcag 600
gaatacatag ggcaactgcc tcctaatttt ccctacagag atgatgtcat gtcagtgaat 660
cctacctgtt tggctcctat tattctcttg tttattagca ttatcttgac ttttaagggg 720
tacttgatta gctgtgtttg gaactgctac cgatacatca atggtaggaa ctccctctgat 780
gtcctgggtt atgttaccag caatgacact acggtgctgc taccctcgta tgatgatgcc 840
actgtgaatg gtgctgccaa ggagccaccg ccaccttacg tgtctgccta agccttcaag 900
tgggcgaggc tgagggcagc agcttgactt tgcagacatc tgagcaatag ttctgttatt 960
tcacttttgc catgagcctc tctgagcttg tttgttgctg aaatgctact ttttaaaatt 1020
tagatgttag attgaaaact gtagttttca acatatgctt tgctggaaca ctgtgataga 1080
ttaactgtag aattcttctc gtacgattgg gtagataatg ggcttcacta acctcccta 1140
ggcattgaaa cttcccccaa atctgatgga cctagaaatc tgcttttgta cctgctgggc 1200
cccaaagttg ggcatttttc tctctgttcc ctctcttttg aaaatgtaaa ataaaacca 1260
aatagacaa ctttttcttc agccattcca gcatagagaa caaaacctta tggaaacagg 1320
aatgtcaatt gtgtaatcat tgttctaatt aggtaaatag aagtccttat gtatgtgtta 1380
caagaatttc cccacaaca tcctttatga ctgaagttca atgacagttt gtgtttgggtg 1440
gtaaaggatt ttctccatgg cctgaattaa gaccattaga aagcaccagg ccgtggggagc 1500
agtgaccatc tgctgactgt tcttggtggat cttgtgtcca gggacatggg gtgacatgcc 1560
tcgtatgtgt tagaggggtg aatggatgtg tttggcgctg catgggatct ggtgccctc 1620
ttctcctgga ttcacatccc caccagggc cgccttttac taagtgttct gccctagatt 1680
ggttcaagga ggtcatccaa ctgactttat caagtggat tggtatata ttgatatact 1740
tctgcctaac aacatggaaa agggttttct tttccctgca agctacatcc tactgctttg 1800
aacttccaag tatgtctagt cacttttaa aatgtaaa ttttcagaaa aatgaggatt 1860
gccttccttg tatgcgtttt ttaccttgac tacctgaatt gcaagggtat tttatatatt 1920
catatgttac aaagtcagca actctcctgt tggttcatta ttgaatgtgc tgtaaatata 1980
gtcgtttgca attaaaacaa ggtttgcccc caaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040
aaaaaaaaaa aaaaaaaaaa caatacgaac caacaagaac agcaagaatc aaaaagctta 2100
aaatcctcga gtgcacaatg aagaaaacag gggggcgccc ctctaggggt cccagtttcg 2160
gtcgggggtga ttggggggac tgagcccttc ctaggggac cccctaattc tgatttcagg 2220
ggcggggcggt tttttaaaaa ggcgggttaa cgggggaaac acccctgggg gttatcccc 2280
ctcttttaag cggcgtgttg gtgggaaaga caccgccctt ttttgcgccc aagtgggggc 2340
atatatctcc cgggtgaggg cgccctctac ccgatatctg gcctttcccc agaagttttg 2400
cccgcgcct ctactttggg gactttgggt gccgccacct ttattaaggg ggcgtttata 2460
ccagcggcgg gggtttgttg gcgtcctcgc gcgggaggcc cgctccacce t 2511

```

<210> 64

<211> 788

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3255825CB1

<400> 64

```

cccacgcgtc cgcagctgcg tcctctgggc cacgtccac ttgcccgtt ccacccggaa 60
agccccccag gctgagtgcg gcatgatctc catcaccgaa tggcagaaga ttggtgtggg 120
gatcacgggt ttccggcatct tcttcactct ctttggaaac ctctgtact ttgattccgt 180
gtcctctggc tttggaaacc tgctgttctt gacgggcctg tccctcatca ttggcctgag 240
gaagaccttt tggttcttct tccaacggca caaactcaag ggaaccagct tcctcctggg 300
gggtgtggtt atcgtgctcc tacgtggcc cctcctcggc atgttctctg aaacctacgg 360
attcttcagc ctctttaagg gctttttccc tgtgccttc ggttctctgg caatgtctgc 420
aacatcccc tctgggtgc gctgttcgg agacttcaag gcactagctc gatgggtctga 480
aaaacagaga tgagctcctt gaacttggat cattgggtga gggggctaga gggagaatgg 540
gaaccacccc ctcatcccc tgactgact cactccccga catatccgga cctccccagg 600
tccagaagga aggaatggag ctgagcaact gacgtcaaat cccaagtcg actcaagagg 660
ctgccaggaa gcagagatgc agaccccaag gagactgggc tggggctggt atcacacct 720
cactctatat ttatgggagg aaaagtgaag attaaattcc caagttgtgc gtgtgtctaa 780
aaaaaaaaa

```

788

<210> 65

<211> 1831

<212> DNA
<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3393430CB1

<400> 65

```
gccttataaa gtagcctctg catctgcctg cctcgggagc aggagggcta ccctggggct 60
gagagttcac ctgtctcagg aaccacctga gccacacagat cctgtgggca gcggccagg 120
cagccatggc ttgggcaagt aggctggggc tgctgctggc actgctgctg cccgtggctg 180
gtgcctccac gccaggcacc gtggtccgac tcaacaaggc agcattgagc tacgtgtctg 240
aaattgggaa agccctctc cagcggggcc tgcaggtcac tgtccctcat ttcctggact 300
ggagtggaga ggcgcttcag cccaccagga tccggattct gaatgtccat gtgccccgcc 360
tccacctgaa attcattgct ggtttcggag tgcgcctgct ggcagcagct aattttactt 420
tcaaggctct tgcgccccca gagccctgg agctgacgct gcctgtggaa ctgctggctg 480
acacccgcgt gaccagagc tccatcagga cccctgtggc cagcatctct gcctgtctct 540
tattctcggg ccacgccaac gagtttgatg gcagtaacag cacctccac gcgctgctgg 600
tcctggtgca gaagcacatt aaagctgtct tgagtaacaa gctgtgcctg agcatctcca 660
acctggtgca ggggtgtcaat gtccacctgg gcaccttaat tggcctcaac cccgtgggtc 720
ctgagtcaca gatccgctat tccatgggta gtgtgcccac tgtcaccagt gactacattt 780
ccctggaaagt caatgctgtt ctcttcctgc tgggcaagcc catcatcctg cccacggatg 840
ccaccccttt tgtgttgcca aggcattgtg gtaccgaggc ctccatggcc accgtggggc 900
tctcccagca gctgtttgac tctgcgctcc tgctgctgca gaaggccggt gccctcaacc 960
tggacatcac agggcagctg aggtcggatg acaacctgct gaacacctct gctctggggc 1020
ggctcatccc ggaggtggcc cgccagtttc ccgagcccat gcctgtgggt ctcaagggtc 1080
ggctgggtgc cacacctgtg gccatgctcc acacaaacaa cgccacctg cggctgcagc 1140
ccttcgtgga ggtcctggcc acagcctcca actcggcttt ccagtccctc ttctccctgg 1200
atgtggtagt gaacttgaga ctccagctct ctgtgtccaa ggtgaagctt caggggacca 1260
cgtctgtgct gggggatgtc cagctcacgg tggectctc caacgtgggc ttcatgata 1320
cagatcaggt gcgcacactg atgggcaccg tttttgagaa gccctgctg gaccatctca 1380
atgctctctt ggccatggga attgcccctc ctgggtgtgt caacctccac tatgtcgccc 1440
ctgagatctt tgtctatgag ggctacgtg tgatatccag tggactcttc taccagagt 1500
gaggcaagac cactgggagg cctgagagtg ggccagctcg ctgctcaggc gaatttctca 1560
tttcaagcca ctggggaaac tgaggcaaaa ccatacttag tcatcaccaa caagctggac 1620
tgcttagctg ggctgtttta tcttccctga gtgcctgggt ctccctccct cacttctgcc 1680
ctttcccttc ctccctctct tctcctccct ctccctccat ctcccccctc ctccctctgc 1740
cccaccccag gggggagcag actgctcctc caggctgtat agacctgccc tcttgcat 1800
aacaacttct cttgagctgc aaaaaaaaaa a 1831
```

<210> 66

<211> 1499

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3490990CB1

<400> 66

```
tgtttgagtt ggggtgaagaa ggcgggaagt taatcctctt ccaatcagct ccccatctcg 60
ccctctggcg gctccggacc acaggccgag ttactctgat taccgccatg ctccaagacc 120
cgcactgaca gtccccacct cccgttctct gtcatcggtt cctcaatgg agtccacatg 180
tttgggcaga atctggaggt gcagctgagc tctgcgagga ccgagaacac gactgtgggt 240
tggaaaagct tccatgacag catcacctc attgttctgt catctgaggt gggcatctct 300
gagctgagggc tggagagact actccaaatg gtgtttggag ccattggtcct tcttgtggga 360
cttgaagaac tgaccaatat ccgcaacgtg gagagactga agaaggactt gagggccagt 420
tattgcctca tcgacagctt cctgggggac tcggagctca tcggggacct gaccagtgt 480
gtggactgcg tgattcctcc agaggggtcc ctcttgacag aagccctctc cgggttcgct 540
gaggccgcgg gcacgacctt cgctcagtct gtggtgtccg gccgggtggg ggcagcaaca 600
gagggttggg ggcggctggg gacgcccag gccgtgctgc tcccctggct ggtgggggtc 660
ctgccgccgc agaccgctcg cgactaccgg gtgtacctac cgcacgggag cccacgggtc 720
ccacaccggc tctgacctt gactctgctg ccgagcctgg agctgtgtct actctgcggg 780
```

```

ccgagcccac ccctcagcca gttgtatcca cagcttcttg agcgctggtg gcagccactg 840
ctggaccgct tgccggcctg tctgccgttg ggaccccggg cgctgcccag tggcttcccc 900
cttcacacag acatcctcgg gctgctgctc ctccacctgg aactgaagcg ctgcctcttc 960
accgtggagc ccttggggga taaagagcct tcaccagaac agcgccggcg cctcctccga 1020
aacttctata ccctggtcac ctccacgcac tccccaccag agccagggcc accagagaag 1080
acagaagatg aggtctacca ggcccagctg cccagagctt gctacctggt gttggggact 1140
gaggaaccag gcacaggagt gcgtctggtg gccttgcaac tggggcttcg gcggctgctg 1200
ctgctgctgt ctcccagag tcccacccat gggtgcgaa gcctggccac ccacactctg 1260
catgcctca cccacttct ttgactacct agcagtgggt gatggacaca gacatggggc 1320
tgtagcgtc tctctgttta ttcgctcaca ataatacaca gccctggat ggggagggg 1380
taggaggggc tacaacaggg tggggtggga ggggaggaga catccacttc cctggccct 1440
ctcccctctg tgcttggggg ggaaagggag ggagggggac tcccctaac ccccagaa 1499

```

<210> 67

<211> 365

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3635154CB1

<400> 67

```

gtttttttgc actaacttca ggagccagct cgtgatctca ggatgtatgg aaaaataatc 60
tttgtattac tattgtcaga aattgtgagc atatcagcat caagtaccac tgggtgtggca 120
atgcacactt caacctcttc ttcagtcaca aagagttaca tctcatcaca gacaaatgga 180
gaaacgggac aacttgtcca tcgtttcact gtaccagctc ctgtagtgat aatactcatt 240
attttgtgtg tgatggctgg tattattgga acgatcctct tattttctta cagttttcgc 300
cgactgataa agggctgagg gtgtagcctg catgctgccg atcttgctct gaaccggccg 360
cattg
365

```

<210> 68

<211> 1102

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4374347CB1

<400> 68

```

gaggacagag caggcagcag agaccatggg gccccctca gcttgtcccc acagagaatg 60
catcccctgg caggggctct tgctcacagc ctactttta actttctgga acgcacccac 120
cactgcctgg ctctttattg catcagcgcc ctttgaagtt gctgaagggg agaatgttca 180
tctctctgtg gtttatctgc ccgagaatct ttacagctat ggctgggtaca aagggaacac 240
ggtggagccc aaccagctaa tcgcagcata tgtaatagac actcacgtta ggactccagg 300
gcctgcatac agcggtcgag agacaatatc acccagtggg gatctgcatt tccagaacgt 360
caccctagag gacacgggat actacaacct acaagtcaca tacagaaatt ctcatattga 420
acaggcatct caccatctcc gtgtatacga gtcagtggct cagccctcca tccaagccag 480
cagcaccaca gtcacagaga agggctccgt ggtcctgacc tgccacacaa ataacttg 540
aacctctttc cagtggattt tcaacaacca gcgtctgcag gtcacgaaga ggatgaagct 600
gtcctggttt aaccatgtgc tcaccataga ccccatcagg caggaggacg ctggggagta 660
tcagtgtgag gtctccaacc cagtcagctc caacaggagc gacccctca agctgactgt 720
aaaatatgac aacactctag gcatcctgat cggggtcctg gttgggagtc ttctggtggc 780
tgcaactgtg tgtttcctgc tctccgaaa aactggcagg gccagcgatc agagtgactt 840
cagggagcag cagccccagc cctccacccc cggccatgga ccctctgaca gctctgacag 900
ctccatctcc taggaattgc tacactctga cacaaacatt tactgctgga tcgaccacaa 960
agcagatgtg gcttcttagg ttctcttggg agctgctcct gtgggttgat ggagcgctcc 1020
tgaagcccc agccctgggg atggggaagg acatggagcc tgagccagag aaccagctct 1080
gagtcctgag gagacacagg cc
1102

```

<210> 69

<211> 2546

<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4596747CB1

<400> 69
 cccgggggagc gcccaccgag cgcgcatccat gttcgacacc acaccccact ctggccggag 60
 caccgcaagc agctcccat cgctccggaa acggctgcag ctcccgcccc caagccggcc 120
 cccacctgag ccagaaccag gcacatggt ggagaagga tcagatagct cctcagagaa 180
 ggggtgggtg cctgggaccc ccagcaccga gagcctaggc agccggaact tcatccgcaa 240
 cagcaagaag atgcagagct ggtacagtat gctgagcccc acttataagc agcgtaatga 300
 ggacttccgg aaactgttca gcaaaactccc cgaagcagaa cgctcattg tggattactc 360
 ctgcgccctg cagcgtgaga tcctgctcca gggccgcctc tacctctctg agaactggat 420
 ctgcttctac agcaacatct tccgctggga gaccacgac tccatccagc tgaagggaag 480
 gacatgtctg aagaaggaaa agacggccaa gctgatcccc aacgccatcc agatctgcac 540
 ggagagcgag aagcatttct tcaacttctt tggggcccg gaccgctgct tcctcctcat 600
 cttccgcctc tggcagaatg cactgcttga aaagacgctg agtccccgag agctctggca 660
 cctgggtgcat cagtgcacg gctcagagct gggcctcacc agtgaggatg aggactatgt 720
 ctcccccttg cagctgaacg gtctggggag ccccaaggaa gtgggagatg tgatcgccct 780
 gagecgacatc acctcctcgg ggcagctga ccgcagccag gagccaagcc cagtgggttc 840
 gcgcggtggc catgtcacg ccaaccttct ccgagccagc agcgacgcag acctggggc 900
 agagcaggagc aaggaggagc aggtagacag ccagccagac gcctcctcca gccagacagt 960
 gaccccggtg gctgaacccc cgagcacaga gcccaccag cctgacgggc ccaccaccct 1020
 gggccccctg gatctgctgc ccagtggaga gctattgaca gacacaagta actcctcttc 1080
 atccactggg gaggaagcgg acttggtgc cctgcttccc gacctctccg gccgcctcct 1140
 catcaactct gtcttccatc tgggcgctga gcggctccag cagatgctct tctcgactc 1200
 gccctcctc cagggtctcc tacagcagt caagttcaca gacgtgacgc tgagccctg 1260
 gagtggggac agcaagtgcc accagcgccg ggtgctgacg tacaccatcc ccatcagcaa 1320
 cccactgggc cccaagagcg cctccgtggt ggagacacag acgctgttcc ggcgcggccc 1380
 ccaggccggc ggggtgtgtg tggactccga ggtgctgacg cagggcaccc cctaccagga 1440
 ctactttctac actgcccacc gctactgcat cctgggtctg gcccggaaac aggcgcggct 1500
 ccgagtgtct tctgagatcc gctaccgaaa gcagccgtg agcctggtga agtcgctcat 1560
 tgagaagaac tctgtggagc gcattgaaga ctatttccac catctggagc gagagctcgc 1620
 caaggctgag aagctgtctc tggaggaagg cgggaaggat gcccggggct tgctatccgg 1680
 cctgcggcgg cggaagcggc ccctgagctg gcgggctcac ggggacgggc cccagcacc 1740
 agatcctgac ccctgtgccc gggccggcat tcacacctcg ggctccctca gctcccgctt 1800
 ctccgaacca tctgtggacc agggcccccg ggcaggcatc ccagtgccc tgggtctcat 1860
 cagcattgtg atctgtgtga gccttatcat cctcatcgcc ctcaacgtcc tgctcttcta 1920
 ccgctctctg tccctggaaa ggacagccca cacctttgag tcctggcaca gcctggccct 1980
 ggccaagggc aagttccccc agacggccac agagtgggac gagatcctgg cgctgcagaa 2040
 gcaattccac agcgtggagg tgcacaagtg gaggcagatc ctgcgggcct ccgtggagct 2100
 cctggatgag atgaagttct cgctggagaa gctgcacaa ggcacacag tctcagaccc 2160
 tcccttttag acccagcccc ggcccgatga cagcttttcc tgaggacccc ggccacgcag 2220
 ctgttccccc acatggacag atggacacac agagcctcgg cggccactgc tggcacggtg 2280
 tgagcgccag gcatctccca cccgccccctc ccgacggccc aaccaggggc tgtgcagacg 2340
 tggggaccac ggaaccgaga tgcactttag accagggagc tggcccggcc tctggcaggc 2400
 cccccactaa cttattttgc ccggtgagg ttgtgggggg cgctcctg ggtgcacgat 2460
 tccctcagct ctgggtttta tgtattatat ttatttgggg ccgacagtgc cccaataaag 2520
 ggtcagaagt gaaaaataaa aaaaaa 2546

<210> 70
<211> 1845
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5052680CB1

<400> 70
 ctctggcgga ctgcctggcg gaagcgggaa cgctcgcatcc tgaggtaaag gtgcacggca 60

```

tctctgggaca tgtagtcttg cccggggctcg gacgccccct cggatgaatg ggaccgaagc 120
tgacttgcgaa ctacagcttc ttggcagcgt cgggtgttggc cgcgggagaa ggggagaccg 180
cggcgggcccc cagtgaagagc ggctttccag gacggtgcga tgtgctgcgc agcgaagagg 240
caggaggccg gcttctctgg gtagcgggtac aggcggggcgc ttactctgtg cgcttgcttc 300
cccaaccctg caccggccat gcgcccggcc ttggcgggtgg gcctgggtgt cgcaggctgc 360
tgcagtaacg tgatcttcct agagctcctg gcccggaagc atccaggatg tgggaacatt 420
gtgacatttg cacaattttt atttattgct gtggaaggct tcctctttga agctgatttg 480
ggaaggaagc caccagctat cccaataagg tactatgcca taatggtgac catgttcttc 540
accgtgagcg tgggaacaa ctatgccctg aatctcaaca ttgccatgcc cctgcatatg 600
atatttagat ccggttctct aattgccaac atgattctag gaattatcat tttgaagaaa 660
agatacagta tattcaataa tacctccatt gccctggtgt ctgtggggat atttatttgc 720
acttttatgt cagcaaagca ggtgacttcc cagtccagct tgagtgaaga tgatggattc 780
caggcatttg tgtggtggtt actaggtatt ggggcattga cttttgtctt tctgatgtca 840
gcaaggatgg ggaatttcca agagactctc tacaacgat ttgggaaaca ctccaaggag 900
gctttgtttt ataatacagc ccttccactt ccgggtttcg tcttcttggc ttctgatatt 960
tatgaccatg cagttctatt caataagtct gagttatatg aaattcccgt catcggagtg 1020
accctgcccc tcatgtggtt ctacctctc atgaacatca tcactcagta cgtgtgcatt 1080
cggggtgtgt ttatcctcac cacggaatgc gcctccctca ccgtcacgct cgtcgtgacc 1140
ctacgcaaat ttgtgagcct catcttttcc atcttgtact tccagaacct cttcaccctg 1200
tggcactggc tgggcacctt gtttgtcttc attgggacct taatgtacac agagggtgtg 1260
aacaacctag ggaccacaaa aagtgagcct cagaaggaca gcaagaagaa ctgaggcctg 1320
tctggagtac gtagaccagt gtcgtcgtga ggggtggacc ctgtgaaggc ctgaccaccg 1380
tttcgctttt gttaatgccg agctacccgc agtgctgagc cagccgtgca aaaggaaatc 1440
ttcaggaggg gacttctcac gttgctcaga ctgacacatg tagactaaat agaaaccctt 1500
cagccctaaa atagaaaaaa gaacaggctt agactattga acaggcgatt catttatattc 1560
tgttttgttt accgacatat tcagttattt tcttgttttc tgaattctga gtctcctgaa 1620
caaaaggctt actatccatg gtcttggaaa agattgtccc ttctcttggc gttgtaatgc 1680
atctctgcta cagcagtcac tgctcatcct tgaccttacc ccttttgtct cccatcacct 1740
tttccttctt cagcagcacc tcacagtcag tcagacgctg caggcaccac tatcaccagg 1800
agtggccaag taaaccgaga caccagtaac agcagggtgg acttc 1845

```

<210> 71

<211> 1940

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5373575CB1

<400> 71

```

gtctcctgtc aaaagccatg ctccggcaggc ctgggtaccg ggcgctgccc ctgggtgatt 60
ttgaccgctt ccagcagtcg agcttcggct ttctgggctc gcagaagggc tgcttgctcc 120
cggagcgggg cggcgtgggg acagggggcc atgtacccca gagctggccc tctgctctcc 180
gtcatggcct catcagtttc ctggggttct tctgctgtt ttgtcaccttc cccatttctg 240
gctggtttgc cctgaagatt gtgcccacct acgagcggat gattgtgttc cgctggggcc 300
ggatccgcac cccccaggga cctggcatgg ttctgtctct gcccttcatt gactcctttc 360
agagggtgga tctgaggaca cgagccttca acgtccctcc ctgcaagctg gcctctaagg 420
acggggctgt gctgtccgtg ggagccgatg tccagtttcc catctgggac ccggtgctgt 480
cgggtgatgc tgtgaaagac ctgaacacag ccacacgcac gacagcccag aacgccatga 540
ccaaggccct gctcaagagg ccgctgcggg agatccagat ggagaagctc aagatcagcg 600
accagcttct gctggagatc aacgatgtga ccagggcctg ggggctggag gtagaccgcg 660
tggagctggc agtggaggcc gtgctccagc cgcgccagga cagcccagct gggcccaacc 720
tggacagcac cctccagcag ctggccctgc acttccctgg aggaagcatg aactcaatgg 780
caggaggtgc cccgtccccg gggccagcag acaccgtgga gatggtgagt gaagttgagc 840
cacctgcccc tcaagttggt gccaggtcca gtcgaagca gcctctggcg gaggggctac 900
tgactgtctc acagcccttc ctgtctgagg cctgtgtcag ccaagtccgg gcctgtacc 960
agttcaatgt cgtcctgccc agcggcaccc aaagcgcta ctctctggac ctactacag 1020
gacgaggaag agtgggacac ggggtgcctg atggcatccc tgatgtgggt gtggagatgg 1080
ccgaggcaga cctgcggggc ctgctatgca gagagctgcg gccctgggg gcctacatga 1140
gtggacggct gaaggtgaag ggcgacctgg ctatggccat gaagctggag gctgtcctca 1200
gggccttgaa gtagcagcct tggctgactt tccagagccc agtcccaagc ctggcaccaa 1260
gcccaggggg cctcttggag gaggaggtgt tcactctgcac cacagagagt tgaggcccta 1320

```

acaaatttca	ggcccagcca	agagcccatg	aatggaggct	gcaggaggct	gagtcaggct	1380
gccatgcacg	tctcccctac	agtgggtctc	tggacaaggc	tttgtccatc	ccggtcccca	1440
gctgagtgcc	cagcgctgag	ctgggtgcac	ggtgtgattc	caggaggaga	gccaggcctg	1500
ccctgccctg	ctggcttcct	gactggagag	acaggacca	cagaaacagc	ctgacagcag	1560
ctggtttggg	ccttgtgtga	gggaccaagc	atgtggccca	ggctctaagc	tctgcgggga	1620
ttggagaggg	atggggaggg	aagggaaggc	agctccaaga	agaggccct	gtggcgaagt	1680
tacctgggga	tcctggctgg	cccaccttcc	tggctgcagt	ccaggcccgt	gctggcgagg	1740
ttgggcatgg	gaaggagcag	ggcctgctgc	ttccctggcg	ctgctcccaa	agatttctga	1800
ctcatctgcc	agctctgtcc	tgcctgcctg	gcgagctggg	gcccagggca	gcataagga	1860
gagcctgcg	ttctgtgctt	cttaccagag	gtttgcaagc	ctcagacaaa	taaattgtgt	1920
gtttacaatg	taaaaaaaaa					1940

<210> 72

<211> 880

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5524468CB1

<400> 72

gcgaagccga	agggagcgcg	gctaagagtg	ccgcaccgcc	tcacaacctg	ggaaccggag	60
agtagggggc	gtcggctggc	agaagccgc	cgtgcctcct	cggcaagggc	catccgggtg	120
cacccatgtc	gcactagagc	agaagagggg	gagtcctgga	actgcaacct	gcacagagct	180
gctctgtact	gtccctgggt	gtcgccgcca	tgacctggtt	ggtgctgctg	gggacactgc	240
tctgcatgct	gcgcgttggg	ttaggcaccc	cggactccga	gggtttcccg	ccccgtgcgc	300
tcacaaactg	cccctacaaa	tgtatctgcg	ctgccgacct	gctaagctgc	actggcctag	360
ggctgcagga	cgtgccagcc	gagttacctg	ccgctactgc	ggacctcgac	ctgagccaca	420
acgcgctcca	gcgcctgcgc	cccggctggg	tggcgccctt	cttccagctg	cgcgccttgc	480
acctagacca	caacgaacta	gatgcgctgg	gtcgcggcgt	cttcgtcaac	gccagcggcc	540
tgaggctgct	cgatctatca	tctaacacgt	tgcggggcgt	tggccgccac	gacctcgacg	600
ggctgggggc	gctggagaag	ctgcttctgt	tcaataaccg	cttggtgcac	ttggacgagc	660
atgccttcca	cggcctgcgc	gcgctcagcc	atctctacct	gggctgcaac	gaactcgctt	720
cgttctcctt	cgaccacctg	caaggctctg	gcgccacca	cctgcttact	ctggacctct	780
cctccaaccg	gctgggacac	atctccgtac	ctgagctggc	cgcgctgccg	gccttctctc	840
agaacggcct	ctacctgcac	gacaacacat	aaaaaaaaaa			880

<210> 73

<211> 2403

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5944279CB1

<400> 73

agccacctgc	tggccaggta	cccctccctt	acctggggca	gtgtctgcct	ggtggccact	60
agagacagcc	cagcctgggc	catggaagaa	aaccgcacct	tggaaatcaga	agcctggggc	120
tcctctaggg	ggtggctggc	cccccgagg	gccagaggag	gcccctcgct	gtcttctgtg	180
ctgaacgagc	tgcccagtgc	tgccaccctt	cggtaaccgag	accctggggg	gctgccttgg	240
ggggcgctgg	aggaggagga	ggaggatgga	ggaaggagca	gaaaggcctt	cacagaagtc	300
acccagacag	agctgcagga	ccctcaccct	tcccgggaac	tggcctggcc	catgcaggcc	360
agacgggcac	acaggcaaag	aaatgccagc	agggaccagg	tggctctatg	ctctgggaact	420
aagacggacc	gatgggcgcg	gctacttcgg	aggtccaagg	agaaaacaaa	ggaaggcttg	480
cgaagcctgc	agccctgggc	gtggacactg	aagaggatcg	ggggccaggt	tggcgccggc	540
acggagtcct	acttctccct	gtgcgcttcc	ctgctccttc	ttaacgtgct	ggcctctgtg	600
ctcatggcct	gcatgacgct	gctgcccacc	tgggtgggag	gcgctcccc	agggcctccc	660
ggccccgaca	tctctctgcc	ctgcggctcc	tataaccccc	actcccaggg	cctggtcacc	720
tttgccaccc	agctcttcaa	cttgctctcg	ggtgagggtt	acctggaatg	gtccccctct	780
ttctatggct	tctacccgcc	ccgcccacgc	ctggcggtca	cctacctgtg	ctgggccttt	840
gccgttggcc	tcactctgct	cctgctcctc	ctgcctcgct	cgggtgtctg	gctgaagcag	900

```

acactgctgg cggagtcgga ggctctgacc agctacagcc accgggtgtt ctcggcctgg 960
gacttcggtc tctgcgggga cgtccacgtg cggctgcgcc agcgcatcat cttgtacgaa 1020
ttaaagggtg agctggagga gacagtgggtg cggcgccagg ctgcggtgcg gacgctgggc 1080
cagcaagcca ggggttgggtt ggtgcgggtg ctgctcaacc tgctgggtgg cgcgctcctg 1140
ggggcagcct tctatggcgt ctactgggct acggggtgca ccgtggagct gcaggagatg 1200
ccccctgtcc aggagttgcc actgctgaag cttgggggtga attaccttc gtccatcttc 1260
atcgctgggg tcaattttgt gctgccgccc gtgttcaagc tcattgctcc actggagggc 1320
tacactcgga gtcgccagat cgtttttatc ctgctcagga ccgtgtttct tcgcctcgcc 1380
tccctgggtg tcctgctctt ctctctctgg aatcagatca cttgtggggg cgactccgag 1440
gctgaggact gcaaaacctg tggctacaat tacaacaac ttccgtgctg ggagactgtc 1500
ctgggcccagg aaatgtacaa acttctgctc tttgatctgc tgactgtctt ggcagtcgcg 1560
ctgctcatcc agtttcctag aaagctcctc tgtggcctct gtccctggggc gctgggtcgt 1620
ctggcgggga cccaggagtt ccaggtgccc gacgaggtgc tggggctcat ctacgcgcag 1680
acgggtggct ggggtgggga ttttttctgc cctgtcttaa cacggtcaag 1740
ttcctgctgc tttctacct gaagaagctt accctcttct ccacctgctc cccggctgcc 1800
cgcaccttcc gggcctccgc ggcgaatttc tttttccctt tggctcttct cctgggtctg 1860
gccatctcca gcgttcccc gctttacagc atcttctga tcccgccttc taagctttgt 1920
ggtccattcc gggggcagtc gtccatctgg gccagatcc ctgagttctat ttccagcctc 1980
cctgagacca cccagaattt cctcttcttc ctggggagcc aggccttttg tgtgccccct 2040
ctgctgatct ccagcatcct gatggcgta acgtgtggctc tggctaactc ctacggagac 2100
ctcatctctg agctcaaacg tcagagacag acggagcgcg agaataaagt cttcctggca 2160
cggcgcgctg tggcgctgac ctccaccaa cgggtcttct gacccccgca gccacgtcc 2220
cgctttcaga ccccaggccc attgtaagcc taggtcacia catctgtaaa ctaggagaa 2280
tggagaagac tccacgcctt tccagctttg gtatctggag atttccaggg cccctcgccg 2340
ccacgtccct gactctcggg tgatcttctt tgtatcaata aatacagccg aggttgggtg 2400
gcc 2403

```

<210> 74

<211> 2850

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6114480CB1

<400> 74

```

cggctcgagc ggctcgagtg aagagcctct ccacggctcc tgcgcctgag acagctggcc 60
tgacctccaa atcatccatc caccctgtct gtcattctgtt ttcattagtgt gagatcaacc 120
cacaggaata tccatggctt ttgtgctcat tttgggtctc agtttctacg agctgggtgc 180
aggacagtgg caagtcactg gaccgggcaa gtttgtccag gccttgggtgg gggaggagcg 240
cgtgttctcc tgctccctct ttcctgagac cagtgcagag gctatggaag tgcgggttctt 300
caggaatcag ttccatgctg tggccacct ctacagagat ggggaagact ggggaatctaa 360
gcagatgccca cagtatcgag ggagaactga gtttgtgaag gactccattg caggggggcg 420
tgtctctcta aggctaaaaa acatcactcc ctoggacatc ggctgtatg ggtgtggtt 480
cagttcccag atttacgatg aggaggccac ctgggagctg cgggtggcag cactgggctc 540
acttctctct atttccatcg tgggatatgt tgacggaggt atccagttac tctgcctgtc 600
ctcaggctgg ttccccagc ccacagccaa gtggaaaggt ccacaaggac aggatattgtc 660
ttcagactcc agagcaaag cagatgggta cagcctgtat gatgtggaga tctccattat 720
agtccaggaa aatgctggga gcatattgtg ttccatccac cttgctgagc agagtcattg 780
ggtggaatcc aaggtattga taggagagac gtttttccag ccctcacctt ggcgcctggc 840
ttctatttta ctcggttac tctgtggtgc cctgtgtggt gttgtcatgg ggatgataat 900
tgttttcttc aaatccaaag ggaaaatcca ggcggaactg gactggagaa gaaagcacgg 960
acaggcagaa ttgagagacg cccggaaaca cgcagtggag gtgactctgg atccagagac 1020
ggctcacccc aagctctgcg tttctgatct gaaaactgta acccatagaa aagctcccca 1080
ggaggtgcct cactctgaga agagatttac aaggaagagt gtggtggctt ctacgggttt 1140
ccaagcaggg agacattact gggagggtga cgtgggacaa aatgtagggt ggtatgtggg 1200
agtgtgtcgg gatgacgtag acagggggaa gaacaatgtg actttgtctc ccaacaatgg 1260
gtattgggtc ctcagactga caacagaaca tttgtatttc acattcaatc cccattttat 1320
cagcctcccc cccagcacc ctcctacacg agtaggggtc ttccctggact atgaggggtg 1380
gaccatctcc ttcttcaata caaatgacca gtcccttatt tataccctgc tgacatgtca 1440
gtttgaaggc ttgttgagac cctatatcca gcatgcgatg tatgacgagg aaaaggggac 1500
tcccatattc atatgtccag tgcctgggg atgagacaga gaagaccctg cttaaaggcg 1560

```


cccacaccac	agacccagac	acagccaagg	gagagtgtc	ccgacaggtg	gccccagctt	1620
cctctccgga	gcctgcgcac	agagagtcac	gccccccact	ctcctttagg	gagctgaggt	1680
tcttctgccc	tgagccctgc	agcagcgga	gtcacagctt	ccagatgagg	ggggattggc	1740
ctgacctgt	gggagtcaga	agccatggct	gccctgaagt	ggggacggaa	tagactcaca	1800
ttaggtttag	tttgtgaaaa	ctccatccag	ctaagcgatc	ttgaacaagt	cacaacctcc	1860
caggctcctc	atttgctagt	cacggacagt	gattcctgcc	tcacaggtga	agattaaaga	1920
gacaacgaat	gtgaatcatg	cttgcaagg	tgagggcaca	gtgtttgcta	atgatgtgtt	1980
tttatattat	acattttccc	accataaact	ctgtttgtct	attccacatt	aatttacttt	2040
tctctatacc	aatcaccca	tggaaatagt	attgaacacc	tgctttgtga	ggctcaaaga	2100
ataaagagga	ggtaggattt	ttcactgatt	ctataagccc	agcattacct	gataccaaaa	2160
ccaggcaaag	aaaacagaag	aagaggaagg	aaaactacag	gtccatatcc	ctcattaaca	2220
cagacacaaa	aattctaaat	aaaattttta	caaattaaac	taaacaatat	atttaaagat	2280
gatatataac	tactcagtgt	ggtttgtccc	acaaatgcag	agttggttta	atatttaaat	2340
atcaaccagt	gtaattcagc	acattaatag	agtaggaaag	aaaccatagg	aaggagagac	2400
gaaggaaaga	gaaaaggggg	cgccccaaact	agttagctgg	caccccgggg	attttccggc	2460
gcggactctc	aggggggtag	cggggtatcga	agcagctttg	aagacggaga	ctcggggggg	2520
gccgggacca	aattcccca	gaggggcggt	tatcggcgcg	cgctggcctc	ggtgttaaac	2580
gccggagtgg	gaaccaggg	gtaccaagtt	agggcttgag	ccgaccgtat	gttgggcttc	2640
cagctagaga	ctgcaaggaa	agaagaagga	ggctagaggc	caggatagtg	gagagagaac	2700
agacgcgaga	acaaggggaa	atgaggacgc	gaggggagag	atgaggagag	atagagatgt	2760
gacgagcagc	gaggagagcc	gggaccggag	aggcaacgga	gaagtgcga	acgaagaaag	2820
acaagggagc	gaaggaacgc	agaagtggaa				2850